

TRANSLATIONAL SCIENCES



Cell Surface Platelet Tissue Factor Expression: Regulation by P2Y₁₂ and Link to Residual Platelet Reactivity

Marta Brambilla¹, Alessia Becchetti¹, Gian Enrico Rovati¹, Nicola Cosentino, Maria Conti¹, Paola Canzano, Peter L.A. Giesen, Alessia Loffreda¹, Alice Bonomi, Marco Cattaneo¹, Erica De Candia¹, Gian Marco Podda¹, Daniela Trabattoni¹, Pablo José Werba¹, Jeness Campodonico¹, Christian Pinna¹, Giancarlo Marenzi¹, Elena Tremoli¹, Marina Camera¹

BACKGROUND: ADP-induced platelet activation leads to cell surface expression of several proteins, including TF (tissue factor). The role of ADP receptors in platelet TF modulation is still unknown. We aimed to assess the (1) involvement of P2Y₁ and P2Y₁₂ receptors in ADP-induced TF exposure; (2) modulation of TF^{pos}-platelets in anti-P2Y₁₂-treated patients with coronary artery disease. Based on the obtained results, we revisited the intracellular localization of TF in platelets.

METHODS: The effects of P2Y₁ or P2Y₁₂ antagonists on ADP-induced TF expression and activity were analyzed in vitro by flow cytometry and thrombin generation assay in blood from healthy subjects, P2Y₁₂^{-/-}, and patients with gray platelet syndrome.

Ex vivo, P2Y₁₂ inhibition of TF expression by clopidogrel/prasugrel/ticagrelor, assessed by VASP (vasodilator-stimulated phosphoprotein) platelet reactivity index, was investigated in coronary artery disease (n=238). Inhibition of open canalicular system externalization and electron microscopy (TEM) were used for TF localization.

RESULTS: In blood from healthy subjects, stimulated in vitro by ADP, the percentage of TF^{pos}-platelets (17.3±5.5%) was significantly reduced in a concentration-dependent manner by P2Y₁₂ inhibition only (-81.7±9.5% with 100 nM AR-C69931MX). In coronary artery disease, inhibition of P2Y₁₂ is paralleled by reduction of ADP-induced platelet TF expression (VASP platelet reactivity index: 17.9±11%, 20.9±11.3%, 40.3±13%; TF^{pos}-platelets: 10.5±4.8%, 9.8±5.9%, 13.6±6.3%, in prasugrel/ticagrelor/clopidogrel-treated patients, respectively). Despite this, 15% of clopidogrel good responders had a level of TF^{pos}-platelets similar to the poor-responder group. Indeed, a stronger P2Y₁₂ inhibition (130-fold) is required to inhibit TF than VASP. Thus, a VASP platelet reactivity index <20% (as in prasugrel/ticagrelor-treated patients) identifies patients with TF^{pos}-platelets <20% (92% sensitivity). Finally, colchicine impaired in vitro ADP-induced TF expression but not α-granule release, suggesting that TF is open canalicular system stored as confirmed by TEM and platelet analysis of patients with gray platelet syndrome.

CONCLUSIONS: Data show that TF expression is regulated by P2Y₁₂ and not P2Y₁; P2Y₁₂ antagonists downregulate the percentage of TF^{pos}-platelets. In clopidogrel good-responder patients, assessment of TF^{pos}-platelets highlights those with residual platelet reactivity. TF is stored in open canalicular system, and its membrane exposure upon activation is prevented by colchicine.

GRAPHIC ABSTRACT: A graphic abstract is available for this article.

Key Words: blood platelets ■ coronary artery disease ■ P2Y₁₂ antagonists ■ P2Y₁₂ receptor ■ thromboplastin

The prothrombotic properties of platelets are the result of their ability to expose phosphatidylserine, which promotes the assembly of coagulation factors

and TF (tissue factor). TF is the key protein triggering the coagulation cascade mostly localized at perivascular level where it provides the hemostatic envelope to prevent

Correspondence to: Prof Marina Camera, Department of Pharmaceutical Sciences, Università degli Studi di Milano, via Balzaretti, 9, Milan 20133, Italy. Email marina.camera@unimi.it

Supplemental Material is available at <https://www.ahajournals.org/doi/suppl/10.1161/ATVBAHA.123.319099>.

For Sources of Funding and Disclosures, see page 2055.

© 2023 The Authors. *Arteriosclerosis, Thrombosis, and Vascular Biology* is published on behalf of the American Heart Association, Inc., by Wolters Kluwer Health, Inc. This is an open access article under the terms of the [Creative Commons Attribution Non-Commercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use, distribution, and reproduction in any medium, provided that the original work is properly cited, the use is noncommercial, and no modifications or adaptations are made.

Arterioscler Thromb Vasc Biol is available at www.ahajournals.org/journal/atvb

Nonstandard Abbreviations and Acronyms

aGPIIb/IIIa	activated glycoprotein IIb/IIIa
CAD	coronary artery disease
GPIIb/IIIa	glycoprotein IIb/IIIa
GPS	gray platelet syndrome
HS	healthy subject
OCS	open canalicular system
PDI	protein disulphide isomerase
PI3	phosphoinositide 3
PLC	phospholipase C
PRI	platelet reactivity index
TF	tissue factor
TG	thrombin generation
VASP	vasodilator-stimulated phosphoprotein

bleeding.¹ Upon activation with ADP, TXA₂ (thromboxane A₂), or thrombin, platelets expose on their surface a functionally active TF, which binds factor VIIa thereby promoting generation of subnanomolar concentration of thrombin.² Thrombin in turn causes further platelet activation via protease-activated receptors that amplifies thrombin formation to achieve hemostasis.^{3,4}

Over the years, increase in the number of prothrombotic TF^{pos}-platelets in different thrombophilic conditions has been extensively described,^{5–9} including in SARS-CoV-2 infection, where we recently showed that platelet-associated TF expression was controlled by antiplatelet drugs, especially clopidogrel.¹⁰

Clopidogrel, prasugrel, and ticagrelor, all used in the treatment of patients with coronary artery disease (CAD), antagonize ADP, one of the several platelet agonists. ADP is stored at high concentrations in platelet dense granules, and, when secreted, it enables platelet activation and aggregation by binding to 2 purinergic receptors on the platelet surface: P2Y₁, which triggers platelet activation, and P2Y₁₂, which amplifies the process potentiating secretion of molecules stored in platelet granules such as P-selectin.^{11,12} Since secreted ADP enhances platelet responses to many physiological agonists including TXA₂, collagen, and thrombin, mainly through P2Y₁₂,¹³ this receptor has become a major pharmacological target, together with cyclooxygenase-1, for the management of diseases associated with arterial thrombosis.¹⁴

Interestingly, unlike the regulation of platelet adhesion and aggregation, the 2 P2Y receptors are differently involved in the prothrombotic activity of platelets. Indeed, inhibition of P2Y₁₂ only, and not of P2Y₁, significantly reduces phosphatidylserine exposure.¹⁵ Whether this occurs also for platelet-associated TF is currently unknown. Since previous studies showing that TF is localized both in α -granules and in the open canalicular system (OCS) of resting platelets,^{16,17} it can be hypothesized

Highlights

- ADP-induced TF (tissue factor) expression on the platelet surface is regulated by P2Y₁₂ receptor activation only.
- In patients with coronary artery disease, P2Y₁₂ antagonists hamper ADP-induced cell surface exposure of open canalicular system–stored TF to a different extent (prasugrel=ticagrelor>clopidogrel).
- Reduction of ADP-induced platelet-associated TF exposure occurs with a P2Y₁₂ inhibition much greater than that required to reduce VASP (vasodilator-stimulated phosphoprotein) platelet reactivity index, P-selectin, and GPIIb/IIIa (glycoprotein IIb/IIIa).
- Cell surface platelet TF expression identifies patients with coronary artery disease who, despite a good clopidogrel response, show a residual platelet reactivity in terms of prothrombotic potential.

that both P2Y₁ and P2Y₁₂ receptors regulate TF expression similarly to the α -granule–stored P-selectin.¹⁵ To date, however, this evidence is lacking.

Based on this rationale, the aim of the study was to assess the involvement of P2Y₁ and P2Y₁₂ receptors in ADP-induced TF expression on the platelet surface. For this purpose, we also took advantage of the analysis of platelets from a P2Y₁₂-deficient patient.

Since our data showed the unique involvement of P2Y₁₂ in platelet-associated TF expression, 2 additional aims were pursued in this study. First, we assessed in vivo TF by P2Y₁₂ antagonists in platelets of patients with CAD, establishing the relationship, if any, between TF expression and the residual platelet reactivity, a condition known to be significantly associated with recurrent cardiovascular events.¹⁸ Second, we reanalyzed the cytoplasmic localization of TF to shed light on how a protein reported to be stored both in the α -granules and in OCS could be regulated by P2Y₁₂ only. This was accomplished by means of a pharmacological approach using cytoskeleton blockers such as cytochalasin D and colchicine and taking advantage from analysis of platelets from patients with gray platelet syndrome (GPS). TF intracellular localization was finally evaluated by immune gold transmission electron microscopy.

METHODS

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Study Population

In vitro experiments were performed on 10 healthy subjects (HS; n=5 men and n=5 women; mean age, 39±6 years) who did not take antiplatelet drugs within 10 days before blood

withdrawal. The effect of clopidogrel on platelet-associated TF expression was assessed on 193 patients with CAD with non-ST-segment-elevation myocardial infarction admitted to Centro Cardiologico Monzino IRCCS from January 2017 to December 2019. Clinical characteristics of the enrolled patients are reported in [Table S1](#). All patients were on aspirin (100 mg/die) and clopidogrel (75 mg/die) and blood was withdrawn between 4 and 30 days from the index event. Exclusion criteria were age >80 years, valvular heart disease, atrial fibrillation, thyrotoxicosis, history of hemorrhagic diathesis, platelet disorder or thrombocytopenia, and thrombolytic treatment. Consecutive patients on prasugrel treatment (n=34; 10 mg/die for at least 4 days) or on ticagrelor treatment (n=31; 90 mg/bis in die for at least 4 days) were recruited for comparison. Patients with CAD not treated with P2Y₁₂ antagonists (n=39) were enrolled as controls. Two patients with GPS (males, aged 31 and 15 years), previously confirmed by mutational screening of NBEAL2,^{19,20} and a subject with inherited severe P2Y₁₂ deficiency (male, aged 65 years)²¹ were also studied.

The study was approved by the ethical committee of the institution, and written informed consent was obtained from all participants according to the principles of the Declaration of Helsinki.

Blood Collection and Platelet Isolation

Whole blood was drawn with a 19-gauge needle without venous stasis into sodium citrate (0.129 M, 1/10 volume/volume)-containing tubes (Vacutainer, catalog number 367704; Becton Dickinson), discarding the first 4 mL. Blood samples were processed within 15 minutes of blood drawing. Please see the [Supplemental Material](#) for extended experimental procedures.

Flow Cytometry Analysis

TF, P-selectin, activated GPIIb/IIIa (glycoprotein IIb/IIIa; aGPIIb/IIIa), CD36, CD40L, P2Y₁₂, and P2Y₁ expression was evaluated by whole blood flow cytometry. Please see the [Supplemental Material](#) for extended experimental procedures.

Thrombin Generation and FXa Generation

TF activity was assessed in thrombin generation (TG) assay, performed by calibrated automated thrombogram. TG was run without adding any exogenous TF but using the platelet-associated TF as the only trigger, as described previously.²² FXa (factor Xa) generation was also assessed. Please see the [Supplemental Appendix](#) for extended experimental procedures.

Electron Microscopy

Platelet TF intracellular localization was analyzed by immunogold electron microscopy. Please see the [Supplemental Material](#) for extended experimental procedures.

In Vitro Studies

Whole blood was incubated with AR-C69931MX (Cangrelor, 1 pM to 100 nM, catalog number 5720/1; Bio-Techne srl) or MRS-2500 (1 pM to 100 nM, catalog number 2159/1; Bio-Techne srl) for 30 minutes at room temperature (timing that, in preliminary time course experiments, provided

maximal and repeatable inhibition). Platelets (1×10^6) were then stimulated with ADP (10 μ M, catalog number A2754; Sigma-Aldrich), U46619 (1 μ M, catalog number 538944; Sigma-Aldrich), or TRAP-6 (thrombin receptor-activating peptide; 10 μ M, catalog number 4017752; Bachem) for 15 minutes at room temperature. Concentration-response curves of TF, P-selectin, and aGPIIb/IIIa expression and VASP (vasodilator-stimulated phosphoprotein) platelet reactivity index (PRI) were obtained by flow cytometry analysis after stimulation of human platelets with ADP (10^{-8} – 10^{-4} M, for 15 minutes), with or without a 30-minute preincubation with AR-C69931MX (10^{-12} – 10^{-5} M).

To assess the involvement of actin and tubulin polymerization in TF and P-selectin exposure, platelets were preincubated for 15 minutes at 37 °C with cytochalasin D (10 μ M, catalog number C8273; Sigma-Aldrich) and colchicine (20 nM to 100 μ M, catalog number C9754; Sigma-Aldrich), respectively, and then stimulated for 15 minutes at room temperature with ADP (10 μ M). Treatment of platelets with *Bordetella pertussis* toxin (5 μ g/mL for 30 minutes, catalog number P7208; Sigma-Aldrich) or with the specific PLC (phospholipase C) inhibitor U73122 (1 mM for 30 minutes, catalog number U6756; Sigma-Aldrich) was performed to characterize the Gi signaling involved in ADP-induced TF and P-selectin exposure. Preincubation with LY294002 (10 μ M for 30 minutes, catalog number 440202; Sigma-Aldrich) was used to assess PI3 (phosphoinositide 3) kinase pathway involvement.

Statistical Analysis

Sample Size Calculation

Assuming that patients with CAD have a mean value of TF^{pos}-platelets of $19.4 \pm 4.5\%$ and that the ratio between good responder and poor responder to clopidogrel treatment is 4:1, a sample of 190 patients (152 good responders and 38 poor responders) was estimated to provide 80% power to deem as significant ($\alpha=0.05$) a group difference of 2.3% of TF^{pos}-platelets, with an SD of 4.5. The sample size of prasugrel- and ticagrelor-treated patients' groups was estimated based on the mean difference value of platelet-associated TF expression in patients treated or not with clopidogrel. A sample of 65 patients was estimated to provide 90% power to deem as significant ($\alpha=0.05$) a difference between groups. Thus, 34 consecutive patients treated with prasugrel and 31 patients treated with ticagrelor were enrolled.

Statistical Analysis

Results are expressed as mean \pm SD and were analyzed by the Student paired *t* test or Mann-Whitney *U* test. Differences between patient groups were performed using 1-way ANOVA followed by Dunnett post hoc analysis. The receiver operating characteristic curve was used to determine the optimal cut-off value of VASP-PRI to detect a percentage of TF-positive platelets $\leq 20\%$. A *P* value of 0.05 was considered statistically significant. Analyses were performed using SPSS statistical package (v9.4). Concentration-response curves were evaluated using GraphPad Prism 8. pA₂s were calculated accordingly to the equation set as described in Prism 8. Parameter errors are expressed as percentage coefficient of variation and calculated by analysis of 4 to 11 different independent experiments. All curves shown are computer generated.

RESULTS

P2Y₁₂ Receptor Is Critical for ADP-Induced Platelet-Associated TF Expression and TG

To assess the contribution of the 2 purinergic receptors to the surface expression of TF, whole blood from HS was stimulated *in vitro* with ADP in the presence or absence of a selective P2Y₁ (MRS-2500) or P2Y₁₂ (AR-C69931MX) antagonist. ADP induced an almost 7-fold increase in the number of TF-positive platelets (Figure 1A) and a 2-fold increase in the expression of the protein/platelet (mean fluorescence intensity; Figure 1B).

Platelet treatment with MRS-2500, at any tested concentration, did not affect either TF exposure (Figure 1A and 1B) or its activity measured as TG capacity (Table 1), indicating that P2Y₁ is not involved in the regulation of platelet-associated TF expression. Conversely, AR-C69931MX concentration dependently prevented ADP-induced TF expression, both in terms of the number of TF-positive cells and mean protein expression, with a 60% inhibition with 10 nM becoming almost complete with 100 nM (Figure 1C and 1D). Similarly, the P2Y₁₂ antagonist concentration dependently delayed ADP-induced TG time, which was significantly impaired by 10 and 100 nM AR-C69931MX (Figure 1E). To demonstrate the direct contribution of platelet TF to TG, the test was performed after preincubation of platelets with a neutralizing α TF antibody. Under these conditions, thrombin generated by platelets was impaired. Of note, the effect of the antibody was the greatest in ADP-stimulated samples, and thus in the presence of a greater number of TF^{pos}-platelets, and was progressively reduced in a concentration-dependent manner in the presence of AR-C69931MX, as the expression of TF decreased (Figure S1).

Of note, the P2Y₁₂ antagonism did not significantly affect the amount of thrombin produced (endogenous thrombin potential; Table 1) when assessed in the presence of an excess of exogenously added phospholipids—to make TF the only rate-limiting contributor of thrombin formation—thus supporting the role of TF primarily in the early kinetic phase of the TG. Conversely, when platelet TG was assessed in a more physiological setting, that is, without any addition of phospholipids—thus making it dependent on both the cell surface platelet TF and the phospholipid surface—inhibition of P2Y₁₂ receptor resulted in both a decrease in the kinetic of thrombin formation (lag time: 31.28±2.43 versus 35.22±3.66 minutes, in control and AR-C69931MX-treated samples, respectively; *P*=0.05) and a reduction in the concentration of generated thrombin (peak: 76.31±11.7 versus 59.13±5.21 nmol/L and endogenous thrombin potential: 1402±93 versus 1312±40.6 nMxmin, in control and AR-C69931MX-treated samples, respectively; *P*=0.03 for both; Figure S2). FXa generation experiments,

performed as an alternative approach to evaluate the modulation of platelet TF by AR-C69931MX, confirmed the reduction of TF activity when platelets were treated with the P2Y₁₂ antagonist but not with the P2Y₁ inhibitor MRS-2500 (−19.5% and +1%, with the P2Y₁₂ and P2Y₁ antagonists, respectively). The specific contribution of TF to FXa generation was evaluated performing the assay in the presence of a neutralizing α TF antibody. Under these conditions, FXa generation was reduced by 20%.

Modulation of TF expression can occur through multiple pathways, such as those induced by thrombin and thromboxane, as shown previously.^{2,23} On this regard, TF expression induced by TRAP-6 (+12-fold) and U46619 (+13-fold) was also partially, although significantly, affected by AR-C69931MX, highlighting the enhancing effect of ADP on the platelet response to PAR-1 agonist peptide thrombin and the thromboxane analog U46619 (Figure 1F and 1G), as previously reported.²²

To confirm the exclusive involvement of P2Y₁₂ in TF expression, we analyzed the blood of a P2Y₁₂-deficient patient.²¹ As expected, P2Y₁₂ expression, assessed by flow cytometry analysis, was lower in the proband than in HS, while P2Y₁ expression was comparable (Figure 1H and 1I). Of note, upon stimulation with ADP, no increase of TF expression on the platelet surface was observed in the P2Y₁₂-deficient patient (Figure 1J and 1K). Conversely, it was readily detectable after stimulation with TRAP-6 (+5-fold) or U46619 (+4-fold). As expected, based on the *in vitro* data, inhibition of the P2Y₁ receptor, which unlike P2Y₁₂ is normally expressed in this subject's platelets, did not affect TF expression by any of the tested agonists (Figure 1J and 1K). Finally, intraplatelet TF levels were similar in the P2Y₁₂-deficient patient and in HS (Figure 1L and 1M) sustaining a comparable TG (Figure 1N). The delay exerted by the neutralizing anti-TF antibody was also comparable between samples from the P2Y₁₂-deficient patient and HS (+3.9 and +3.8±2.7 minutes, respectively).

Altogether, these results highlight that P2Y₁₂, and not P2Y₁, is involved in the regulation of ADP-induced TF exposure on the platelet surface, and this event is prevented by selective pharmacological inhibition of the P2Y₁₂ receptor.

P2Y₁₂ Antagonists Reduce Platelet-Associated TF Expression in Patients With CAD

Since the *in vitro* data showed that AR-C69931MX concentration dependently inhibited TF expression, it can be expected that a similar inhibition of cell surface platelet-associated TF expression *in vivo* is displayed by anti-P2Y₁₂ drugs that are used in the clinical setting. To test this hypothesis, we first analyzed the association between cell surface platelet-associated TF expression and the degree of P2Y₁₂ inhibition, measured by the VASP phosphorylation assay, in clopidogrel-treated




Figure 1. Effect of P2Y receptor inhibition on platelet-associated TF (tissue factor) expression and activity in healthy subjects (HS) and in a P2Y₁₂-deficient patient.

Platelet-associated TF expression was analyzed by whole blood flow cytometry while TF-dependent thrombin generation was measured in isolated platelets by calibrated automated thrombogram (CAT) assay, in HS (**A–G**) and in a P2Y₁₂-deficient patient (**H–N**). Effect of the P2Y₁ antagonist MRS-2500 and of P2Y₁₂ antagonist AR-C69931MX (blue bars), incubated 30 minutes before ADP stimulation, on platelet-associated TF expression (**A** and **C**, percentage of positive platelets; **B** and **D**, mean fluorescence intensity [MFI]) and on the lag time of thrombin generation (**E**) in HS (n=6). The specificity of TF contribution to thrombin generation was evaluated by preincubating resting and ADP-stimulated platelets with a neutralizing anti-TF antibody. AR-C69931MX and MRS-2500 effects on TF expression were (*Continued*)

patients with CAD. The degree of VASP phosphorylation is indeed associated with levels of clopidogrel active metabolite,²⁴ allowing the classification of patients as good responders when the platelet reactivity index is <60%.²⁵ Results show that the percentage of TF-positive platelets was significantly inhibited (−30% versus ADP-stimulated samples) in good responders only, whereas in the poor responders, it was similar to that measured in patients with CAD not on clopidogrel (Figure 2A and 2B; Table S2). Comparable results were obtained when phosphatidylserine exposure on the platelet membrane was assessed under resting conditions in a subgroup of patients (Figure S3). Interestingly, while inhibition of aGPIIb/IIIa showed a trend similar to that of TF, with a ≈40% inhibition in good responder (Figure 2C; Table S2), clopidogrel significantly inhibited P-selectin expression to the same extent in all patients with CAD (−55%), irrespective of the degree of VASP phosphorylation (Figure 2D; Table S2).

These results are consistent with the evidence that the estimated in vitro AR-C69931MX potency (pA2) for inhibition of P-selectin expression is ≈130× greater than that needed to inhibit TF. Inhibition of aGPIIb/IIIa and VASP occurs at the same potency, which is 4-fold smaller than that for P-selectin and 40-fold greater than that for TF (8.6±0.3% CV for TF, 10.2±0.1% CV for aGPIIb/IIIa, 10.3±0.3% CV for VASP, and 10.8±0.3% CV for P-selectin; Figure 2E through 2H).

Thus, to inhibit cell surface TF expression and, therefore, the platelet prothrombotic potential, drug concentration has to be much higher than that needed to inhibit the exposure of the classical platelet activation markers such as P-selectin and aGPIIb/IIIa, as well as VASP.

Of interest, 23 of 144 good-responder patients, accounting for 15% of them, had a percentage of TF-positive platelets greater than the mean value (20.4±8.4%) measured in the poor-responder group (Figure 2A, green dots). This percentage even doubles (reaching 38%) if the mean amount of protein expressed on the platelet surface is considered (Figure 2B, green dots). This finding highlights the presence of a subgroup of patients with a potentially increased TG capacity despite an apparent good pharmacological response to clopidogrel according to the VASP-PRI (although it was significantly higher in the subgroup compared with the good responder patients, 45.9±10.9% and 39.0±13.1%, respectively; *P*=0.0192) and P-selectin and aGPIIb/IIIa expression (Figure S4).

To gain insight into this finding, we evaluated the expression of cell surface platelet-associated TF in patients treated with prasugrel, an irreversible P2Y₁₂ receptor inhibitor, or with ticagrelor, a reversibly binding P2Y₁₂ antagonist. All enrolled patients were good responders with a mean VASP-PRI value significantly lower than that of clopidogrel good-responder patients (17.9±11%, 20.9±11.3% versus 40.3±13%, in prasugrel-, ticagrelor-, and clopidogrel-treated patients, respectively; Figure 3A). After stimulation with ADP, the percentage of TF^{pos}-platelets was significantly lower than that measured in patients treated with clopidogrel (10.5±4.8%, 9.8±5.9%, and 13.6±6.3% in prasugrel-, ticagrelor-, and clopidogrel-treated patients, respectively) and, unlike what was observed in the clopidogrel group, no subject showed levels of TF^{pos}-platelets higher than the mean value (20.4±8.4%) of the clopidogrel poor-responder group. In addition, significantly lower levels of P-selectin and aGPIIb/IIIa were also observed in these patients compared with those of clopidogrel-treated patients (Figure 3B through 3D), highlighting an overall higher degree of inhibition of platelet activation by prasugrel and ticagrelor than by clopidogrel.

Based on these observations, receiver operating characteristic curve was used to determine the optimal cutoff value of VASP-PRI to detect a percentage of TF^{pos}-platelets <20%, the mean value of TF^{pos}-platelets measured in poor-responder patients. This relationship was evaluated both on platelets treated in vitro with AR-C69931MX and on platelets from clopidogrel-treated patients. While in vitro a VASP-PRI of 60% was identified as the matched point to have TF^{pos}-platelet levels lower than 20% (area under the curve, 0.986 [95% CI, 0.959–1.000]; *P*<0.0001; sensitivity, 95%; specificity, 95%), in vivo only a VASP-PRI <20% identifies with a 92% sensitivity (and 23% specificity) patients with a TF^{pos}-platelet level <20%. As observed in the enrolled patients, such a low VASP-PRI value is better obtained with ticagrelor- and prasugrel-treated patients.

TF Is Localized in the OCS

The finding that P2Y₁₂ only is involved in the regulation of ADP-induced TF exposure on the platelet surface, together with the evidence of the different degrees of inhibition of P-selectin and TF expression by P2Y₁₂ inhibition, led us to hypothesize that different pathways or cellular localization are involved in the membrane

Figure 1 Continued. evaluated also upon TRAP-6 (thrombin receptor-activating peptide) and U46619 stimulation (**F** and **G**; n=3). Platelet purinergic receptor expression was analyzed by flow cytometry in a P2Y₁₂-deficient patient (red dot) in comparison with a reference group of HS (**H**: percentage of positive platelets; **I**, MFI). Platelet-associated TF expression upon stimulation with ADP, TRAP-6, and U46619 in the presence or absence of the P2Y₁ receptor antagonist MRS-2500 in P2Y₁₂-deficient patient (**J** and **K**). TF intracellular expression (**L**, percentage of positive platelets; **M**, MFI; red dot, P2Y₁₂-deficient patient). Thrombin generation of untreated platelets from the P2Y₁₂-deficient patient (red lines) was assessed in comparison to that of n=3 HS (**N**). Curves generated by preincubating platelets with a neutralizing αTF antibody to evaluate TF-dependent thrombin formation are also reported. Data are expressed as mean±SD from the indicated numbers of independent experiments and were analyzed by Student paired *t* test or Mann-Whitney *U* test as appropriate. IC indicates intracellular; and PLT, platelets.

Table 1. Effect of AR-C69931MX and MRS-2500 on ADP-Induced Thrombin Generation by Platelets

ADP	–	+	+	+
MRS-2500	–	–	+	–
AR-C69931MX	–	–	–	+
Lag time, min	27.9±9.3	24.4±6.5*	18.1±1	32.4±8.8†
ETP, nM·min	1182.5±267.8	1263.7±303.1	1579.7±101.2	1025.9±343.2
ttPeak, min	33.6±8.7	27.9±7.9‡	26.9±1.7	40.5±7.4§
Velocity index, nM/min	15.1±2.2	17.8±4.5	12.7±2.1	13.1±7.1

Values are mean±SD of 5 independent experiments performed in triplicate. ETP indicates endogenous thrombin potential; and ttPeak, time to peak.

**P*=0.049 vs resting platelets;

†*P*=0.0073 vs ADP-stimulated platelets;

‡*P*=0.050 vs resting platelets;

§*P*=0.0067 vs ADP-stimulated platelets.

exposure of the 2 proteins. Indeed, unlike what was observed for TF, the membrane exposure of P-selectin, a protein stored in platelet α -granules, was significantly modulated by both P2Y₁ and P2Y₁₂ (Figure S5). Engagement of P2Y₁₂ receptor implies the activation of Gi-mediated intracellular signaling pathways. Treatment with *Bordetella pertussis* toxin, by inhibiting the Gi α -mediated signaling (Figure S6A and S6B), or with LY294002, affecting PI3 kinase signaling, downstream Gi $\beta\gamma$ activation (Figure S6E and S6F), reduced ADP-dependent TF expression. Under these experimental settings, P-selectin expression was not affected (Figure S6C, S6D, S6G, and S6H), further highlighting P2Y₁ involvement. As expected, U73112-mediated inhibition of PLC, the common enzyme to both the P2Y₁ and P2Y₁₂, triggered signal mediating intracellular calcium increase, led to a reduction of both TF and P-selectin exposure in response to ADP stimulation (Figure S6I through S6L). Altogether, these results confirm that the mechanism underpinning the membrane expression of TF and P-selectin, upon platelet activation, operates through different signaling pathways according to the receptors involved; they also suggest that TF is not stored in the α -granules.

We tested this hypothesis in patients with GPS, a disease characterized by large platelets with reduced number of α -granules. As expected, indeed, flow cytometry analysis showed that levels of P-selectin, exposed on the platelet surface after stimulation with ADP, TRAP-6, or U46619, as well as its intracellular levels, were considerably reduced in patients with GPS compared with HS (Figure 4A and 4B). As a direct functional consequence, the number of platelet-monocyte and platelet-granulocyte aggregates was also reduced in these patients compared with HS (Figure 4C and 4D). Conversely, of note, cell surface expression of platelet-associated TF in patients with GPS, both in resting conditions and upon stimulation with classical platelet agonists, was comparable, despite the α -granule defect, to that of HS. Not only the number of platelets exposing TF on the cell

surface (Figure 4E) but also the number of TF-positive platelets by intracellular staining (Figure 4F) was similar in patients with GPS and in HS. Moreover, cell surface TF expression was also regulated by the same P2Y₁₂ downstream pathway, being the preincubation with pertussis toxin and LY294002 able to prevent ADP-induced TF expression (Figure 4G and 4H) while not affecting the already low P-selectin expression (Figure 4I and 4J).

This compelling evidence that TF is not stored in α -granules led us to verify its association with OCS, a further proposed intracellular platelet storage site. Immunogold labeling and electron microscopy analysis confirmed that in resting platelets from HS, TF is associated with the OCS rather than with α -granules (Figure 5). Of note, cell surface TF expression upon platelet stimulation is the result of OCS externalization. Preincubation of platelets with cytochalasin D, which blocks actin polymerization, resulted indeed in a statistically significant reduction of ADP-induced TF expression (Figure 6A and 6B), without affecting its intracellular levels (Figure 6C and 6D). Similar data were obtained upon stimulation of platelets by other stimuli, such as TRAP-6 or U46619 (data not shown). Of note, cytochalasin D treatment did not prevent P-selectin (Figure 6E and 6F) and CD36 (Figure 6G and 6H) exposure, which are stored within α -granules, nor that of CD40L, a cytoplasmic protein²⁶ (Figure 6I and 6J), while it reduced the surface recruitment of aGPIIb/IIIa (Figure 6K and 6L) as published previously by Addo et al.²⁷ Also microtubule disruption by colchicine concentration dependently prevented ADP-induced TF externalization (Figure 7A and 7B). Similarly to what observed using cytochalasin D, colchicine did not affect P-selectin exposure, inhibiting α -granule release only at the highest concentration tested (Figure 7C and 7D), while reducing aGPIIb/IIIa membrane expression (Figure 7E and 7F). The same results were also obtained when we analyzed ADP-stimulated platelets from patients with GPS (Figure S7) or TRAP-stimulated platelets from P2Y₁₂-deficient patients (Figure S8).

Altogether, these results suggest that in our experimental conditions, cell surface TF expression is not associated with platelet degranulation but rather with OCS externalization.

DISCUSSION

The present study, by using in vitro and ex vivo experiments combining a cell biology, pharmacological, and electron microscopy approach, provides new insights into the regulation and cellular localization of platelet-associated TF expression, thus expanding the characterization of the platelet prothrombotic properties. These findings, relevant also for patients with CAD treated with P2Y₁₂ antagonists, were supported by 2 human models of platelet defects such as P2Y₁₂ deficiency and GPS. Results show for the first time that ADP-induced TF




Figure 2. Platelet activation marker expression in clopidogrel-treated patients with coronary artery disease (CAD).

Clopidogrel (75 mg/die)-treated patients were divided according to their platelet reactivity index (PRI) measured by VASP (vasodilator-stimulated phosphoprotein) phosphorylation assay by flow cytometry: patients with PRI <60% were defined as responders (n=144; red bars), and those with PRI >60% were defined as poor responders (n=49; blue bars). Flow cytometry analysis of platelet (PLT)-associated TF (tissue factor) expression (**A**, percentage of positive platelets; **B**, mean fluorescence intensity [MFI]), activated GPIIb/IIIa (glycoprotein IIb/IIIa; aGPIIb/IIIa; **C**), and P-selectin (**D**) assessment upon ADP stimulation (10 μ M, 15 min) was reported for the 2 groups of patients and compared with a group of CAD patients (n=39) free of anti P2Y₁₂ treatment (no P2Y₁₂ antagonist; gray bar). Green dots highlight patients who, despite a good clopidogrel response, show a percentage of TF^{pos}-platelets higher than the median value measured in the poor responder group. Dot line in **A** and **B** indicates the mean value measured in healthy subject as reference. Results are shown as mean \pm SD of the percentage and MFI of TF^{pos}-platelets and as percentage of aGPIIb/IIIa-positive and P-selectin-positive platelets. Data were analyzed by 1-way ANOVA followed by Dunnet post hoc analysis. Concentration-response curves of platelet TF (**E**), platelet-aGPIIb/IIIa (**F**), platelet-Psel (**G**) and VASP (**H**) induced by ADP (black squares) and upon AR-C69931MX treatment (open triangles). Agonist EC₅₀ was calculated using a 4-parameter logistic model while antagonist pA_{2s} were calculated as described in Materials and Methods. Results are presented as percentage of protein expression over baseline (s/b; n=4–11). All curves shown were computer generated.




Figure 3. TF (tissue factor) expression in prasugrel-, ticagrelor-, and clopidogrel-treated patients with coronary artery disease (CAD).

Flow cytometry analysis of VASP (vasodilator-stimulated phosphoprotein) platelet reactivity index (PRI; **A**) and platelet-associated TF (**B**), Psel (P-selectin; **C**), and activated GPIIb/IIIa (glycoprotein IIb/IIIa; aGPIIb/IIIa; **D**) expression upon ADP stimulation (10 μ M, 15 min) was analyzed in prasugrel- ($n=34$) and ticagrelor-treated ($n=31$) patients and compared with that measured in clopidogrel-treated patients. Results are shown as mean \pm SD of the percentage of antigen-positive platelets (PLT) and were analyzed by 1-way ANOVA followed by Dunnet post hoc analysis.

expression on the platelet surface is regulated by P2Y₁₂ receptor activation only, being P2Y₁ engagement effectless. Indeed, only P2Y₁₂ antagonism prevented in a concentration-dependent manner TF exposure on the cell surface, in turn affecting the kinetic of TG. This effect was observed both in in vitro experiments and ex vivo in platelets from clopidogrel-, prasugrel-, and ticagrelor-treated patients with CAD.

This study also shows for the first time that downregulation of ADP-stimulated platelet-associated TF exposure occurs with a P2Y₁₂ inhibition much greater—requiring a higher drug concentration—than that needed to reduce aGPIIb/IIIa, VASP, and P-selectin expression. From a clinical point of view, this finding is of relevance since assessment of TF levels and activity allows to highlight patients who, despite a good clopidogrel response according to the VASP-PRI, the gold standard method for the assessment of P2Y₁₂ antagonism, show a residual platelet reactivity in terms of prothrombotic potential.

Finally, we also provide the evidence that TF is stored in the platelet OCS; its appearance on the cell membrane upon activation occurs following OCS externalization, and this event is prevented by cytoskeleton blockers, such as cytochalasin D and, more clinically relevant, colchicine.

P2Y₁ and P2Y₁₂ are the 2 ADP receptors that mediate the amplification of platelet response within the hemostatic and thrombotic process. It is well established that platelet activation (ie, granule secretion and cell aggregation) requires concomitant activation of P2Y₁ and P2Y₁₂

signaling.²⁸ Conversely, regulation of the prothrombotic properties of platelets, essential in secondary hemostasis, appears to be a P2Y₁₂ prerogative. Indeed, it has been previously shown that inhibition of P2Y₁₂ only, and not of P2Y₁, is able to modify the platelet prothrombotic phenotype by decreasing platelet phosphatidylserine exposure.¹⁵ Furthermore, P2Y₁₂ antagonism strongly delayed the kinetic of thrombin formation in PRP (platelet-rich plasma), a process that is mainly TF dependent.^{15,29} Our data complement these findings by showing that TF expression on the platelet membrane following ADP stimulation is also a P2Y₁₂-dependent mechanism since (1) only the P2Y₁₂ antagonist, and not the P2Y₁ antagonist, was able in vitro to inhibit it and (2) no increase of TF expression on the platelet surface was observed in a P2Y₁₂-deficient patient upon ADP activation, whereas the protein was detectable following TRAP-6 or U46619 stimulation. Within this context, our findings add new insights into the regulation of the prothrombotic activity of platelets highlighting that both phosphatidylserine and TF exposure are P2Y₁₂-dependent events. To further complement this scenario, it is worth mentioning that the secretion of platelet PDI (protein disulphide isomerase)—a key enzyme involved in the regulation of TF prothrombotic activity³⁰—is also controlled by P2Y₁₂ thus emphasizing an overall consistent and uniform regulation of the prothrombotic activity of platelets.³¹ Clearly, the platelet-associated prothrombotic response may be elicited through multiple pathways including, for instance, activation of the




Figure 4. Platelet activation marker expression in patients with gray platelet syndrome (GPS).

Platelet-Psel (P-selectin; **A** and **B**) and TF (tissue factor; **E** and **F**) expression under resting conditions and upon ADP (10 µM, 15 min), TRAP-6 (thrombin receptor-activating peptide; 10 µM, 15 min), and U46619 (1 µM, 15 min) stimulation was assessed by flow cytometry and reported as surface (**A** and **E**) and intracellular (IC; **B** and **F**) percentage of positive cells in 2 patients with GPS and in 2 healthy subjects studied in parallel. Percentage of monocyte-platelet aggregates (mon-plt aggre; **C**) and of granulocyte-platelet aggregates (gran-plt; **D**) was also reported. The effect of the Gαi inhibitor *Bordetella pertussis* toxin (PTx, 5 µg/mL; **G** and **I**) and the PI3 (phosphoinositide 3) kinase inhibitor LY294002 (10 µM; **H** and **J**) was analyzed upon ADP (10 µM) stimulation as percentage of TF-positive and Psel-positive platelets. Data are expressed as mean±SD.

thrombin receptors PAR-1 and PAR-4.^{32,33} The possible regulation of platelet-associated TF expression by these other signaling pathways was not analyzed being out of the scope of the study.

Increased levels of TF^{pos}-platelets have been reported in several pathological conditions characterized by a

prothrombotic phenotype, including thrombocytopenia,⁵ cancer,⁷ antiphospholipid syndrome,⁹ viral infections,^{8,10} and cardiovascular diseases.^{6,34} The clinical relevance of these findings lays on the significant correlation between platelet-associated TF expression and its functional activity, measured as TG capacity, to residual cardiovascular




Figure 5. Localization of TF (tissue factor) in human platelets.

Ultrathin Tokuyasu cryosections were single immunolabeled for TF (10 nm gold). Only a fraction of the platelet population expresses TF (A); when expressed, TF localizes in the open canalicular system (OCS; B–D).

risk.^{10,22} Indeed, the translational implications of TF regulation by P2Y₁₂ receptor are extremely relevant in this setting, considering that treatment with P2Y₁₂ antagonists is, together with aspirin, the gold standard therapy for secondary prevention of cardiovascular diseases.³⁵

We report here that in CAD patients good responders to P2Y₁₂ antagonists, based on VASP-PRI <60%, the highest is P2Y₁₂ inhibition, as occurs in prasugrel- and ticagrelor-treated patients, and the lowest is the cell surface TF expression. Interestingly indeed, the receiver operating characteristic analysis showed that in vivo levels of TF^{pos}-platelet <20% (ie, below the levels found in poor-responder patients) are predicted by VASP-PRI ≤20%, value that is consistently obtained in prasugrel- and ticagrelor-treated patients only and not in clopidogrel-treated ones.

Several data in the literature, also supported by our present results, have already established that prasugrel and ticagrelor are able, compared with clopidogrel, to achieve a more pronounced inhibition of platelet activation in terms of P-selectin or aGPIIb/IIIa expression and release of microvesicles.^{36–38} Interestingly, Braun et al³⁹ reported that prasugrel treatment also results in a better inhibition of prothrombotic anionic phospholipids exposure than clopidogrel. In agreement, our study shows that the platelet prothrombotic potential, measured as percentage of circulating TF^{pos}-platelets, was

significantly lower in patients receiving prasugrel or ticagrelor compared with clopidogrel-treated ones. This overall greater inhibition of platelet prothrombotic capacity could, therefore, account for the observed reduction in the incidence of MACE (major adverse cardiovascular events) in patients with ACS (acute coronary syndrome) who received prasugrel or ticagrelor compared with clopidogrel.⁴⁰ Our findings may also help to explain why a positive impact of a personalized antiplatelet therapy has been observed only in trials where patients with high-on-treatment platelet reactivity on clopidogrel (based on aggregometry-based tests) were shifted, to achieve a better platelet inhibition, to ticagrelor or prasugrel and not to high-dose clopidogrel.^{41–47} Indeed, a clinically relevant aspect highlighted in this study is the finding that 15% of clopidogrel good-responder patients show levels of cell surface TF^{pos}-platelets similar to those found in poor-responder patients, thus carrying a residual prothrombotic potential. This finding led us to calculate the potency of AR-C69931MX in inhibiting TF and VASP, as well as other classical markers of platelet activation such as P-selectin and aGPIIb/IIIa. Results showed that the degree of P2Y₁₂ inhibition needed to downregulate ADP-induced TF expression is greater than that needed to inhibit VASP and aGPIIb/IIIa (40-fold). Furthermore, the potency of the P2Y₁₂ antagonists to inhibit P-selectin exposure is even greater. This accounts for our evidence




Figure 6. Involvement of actin polymerization in platelet activation marker expression.

Whole blood was incubated with cytochalasin D (CytD; 10 $\mu\text{mol/L}$, 15 min). Surface expression of platelet-associated TF (tissue factor; **A** and **B**), P-selectin (**E** and **F**), CD36 (**G** and **H**), CD40L (**I** and **J**) and activated GPIIb/IIIa (glycoprotein IIb/IIIa; aGPIIb/IIIa; **K** and **L**) was measured by flow cytometry upon stimulation with ADP (10 μM , 15 min). Intracellular (IC) expression of platelet-associated TF was also assessed (**C** and **D**). Data are expressed as mean \pm SD ($n=3-8$) of percentage of positive platelets (PLT; **A**, **C**, **E**, **G**, **I**, and **K**) or mean fluorescence intensity (MFI; **B**, **D**, **F**, **H**, **J**, and **L**). Results were analyzed by Student paired t test or Mann-Whitney U test as appropriate.

that P-selectin does not indeed differ between good and poor responders, in agreement with published data showing that ADP-induced P-selectin inhibition occurs already at low levels of the P2Y₁₂ blockade.⁴⁸ Although aGPIIb/IIIa evaluation seems to better distinguish the responsiveness to clopidogrel, aggregation-based tests

fail to provide an accurate stratification of patients' thrombotic risk. Indeed, data from different clinical studies, such as the ARCTIC and GRAVITAS trials, showed that, depending on the cutoff value used to define platelet hyperreactivity, the degree of aGPIIb/IIIa inhibition might not be able to fully predict major adverse cardiovascular events in stable patients with CAD treated with clopidogrel.^{44,49-51} By contrast, a subanalysis of the PREPARE POST-STENTING study reported that, in a subgroup of patients with post-PCI (percutaneous coronary intervention) ischemic events—despite optimal inhibition by aspirin and clopidogrel—high-platelet fibrin clot strength and prothrombotic potential, involving elevated TF expression phenotype, are driving factors for ischemic events.^{52,53}

Since recurrent thrombotic events occur in about 1 to 10 patients in the first year after CAD despite treatment with aspirin and the most potent P2Y₁₂ antagonists,⁵⁴ the efficacy of the use of anticoagulants on top of antiplatelet treatment has been explored in the clinical setting. Interestingly, the ATLAS ACS and the COMPASS study reported the benefit of very-low-dose rivaroxaban in reducing MACE when combined with antiplatelet therapy.^{55,56} A remaining challenge is the identification of the biomarkers of platelet-dependent TG that can help to identify patients who would have a clinical benefit from this dual pathway inhibition. In this context, platelet-associated TF could be a good candidate because it might be a potential driver toward adverse cardiovascular outcomes. The predictive value of platelet-associated TF is indeed matter of our currently ongoing investigations. Although $<5\%$ of total thrombin is formed during the initiation step,³ it is important to consider that this is critical in the development of the prothrombotic response.⁵⁷ In the absence of platelet contribution to thrombin formation, thrombin indeed would be rapidly inhibited by circulating plasma antithrombin in <1 minute.⁵⁸ By contrast, the TF-dependent initiation phase could support the continuous generation of thrombin that can persist for months after the index event.⁵⁹

A further important clinical implication of this study results from the evidence that in resting platelets, the pool of intracellular TF is associated with OCS rather than with α -granules, as proposed previously.^{16,17} This evidence, highlighted by pharmacological data and by immune gold labeling and transmission electron microscopy images, was confirmed by the finding that patients with GPS, deficient in α -granules, have normal levels of platelet-associated TF. Cytochalasin D or colchicine treatment—inhibiting actin or tubulin polymerization, respectively, and thus OCS externalization—prevented TF and GPIIb/IIIa exposure on the cell membrane. In this view, in resting platelets, OCS appears to be a cryptic site where proteins such as TF and GPIIb/IIIa are not accessible to their physiological agonists (ie, FVIIa [factor VIIa] to promote TG and fibrinogen to sustain platelet aggregation, respectively). It could be speculated that in this compartment, TF can




Figure 7. Involvement of microtubule disruption in platelet activation marker expression.

Whole blood was incubated with colchicine (20 nM to 100 μM, 15 min) and platelet-associated TF (tissue factor; **A** and **B**), Psel (P-selectin; **C** and **D**), and activated GPIIb/IIIa (glycoprotein IIb/IIIa; aGPIIb/IIIa; **E** and **F**) expression was measured by flow cytometry upon stimulation with ADP (10 μM, 15 min). Data are expressed as mean±SD (n=4–7) of percentage of positive cells (**A**, **C**, and **E**) or mean fluorescence intensity (MFI; **B**, **D**, and **F**) and were analyzed by Student paired *t* test or Mann-Whitney *U* test as appropriate.

be retained in its inactive form. Upon platelet activation, rapid events including platelet shape change, cytoskeleton reorganization, and OCS externalization—bringing these proteins to the cell membrane—occur. Then they can switch to their active form due to the redox enzymatic activity of PDI and become competent for triggering coagulation and platelet plug formation.³⁰

Interestingly, low-dose colchicine on top of standard antiplatelet therapy has been shown to reduce major adverse cardiovascular events either in stable patients^{60,61} or, albeit with less consistent results,^{62,63} in patients with acute cardiovascular disease.^{64–66} Cimmino et al⁶⁷ speculated that colchicine, by modulating platelet aggregation, may be effective in reducing the risk of developing future acute cardiovascular events. Although some evidences indicates that colchicine in the therapeutic range, as that used in this study, is not able to inhibit in vivo platelet aggregation,^{68,69} it could prevent TF exposure on cell surface. Thus, the effect of colchicine, besides its well-established anti-inflammatory properties, could be also related to inhibition of the platelet prothrombotic potential.

CONCLUSIONS

Overall, our findings provide a step forward in the characterization of platelet-associated TF regulation and its pharmacological modulation and support the hypothesis that the therapeutic benefits of the antiplatelet therapy with P2Y₁₂ antagonists are also the result of the inhibition of the cell surface TF expression. Of note, active drug concentrations that adequately inhibit the most studied platelet activation markers may not completely turn off the residual platelet reactivity, and this is particularly relevant in clopidogrel-treated patients. This may explain the adverse events recorded in trials despite an adequate inhibition of platelet aggregation. Within this context, concomitant assessment of both VASP-PRI and TF in P2Y₁₂ antagonist-treated patients could improve the evaluation of the residual platelet TF-dependent prothrombotic potential.

ARTICLE INFORMATION

Received April 21, 2023; accepted July 26, 2023.

Affiliations

Centro Cardiologico Monzino IRCCS, Milan, Italy (M.B., A. Becchetti, N.C., M. Conti, P.C., A. Bonomi, D.T., P.J.W., J.C., G.M., M. Camera). Department of Pharmaceutical Sciences (G.E.R., C.P., M. Camera) and Unità di Medicina II, ASST Santi Paolo e Carlo, Department of Scienze della Salute (M. Cattaneo, G.M.P.), Università degli Studi di Milano, Italy. CoagScope BV, Maastricht, the Netherlands (P.L.A.G.). Experimental Imaging Center, San Raffaele Scientific Institute, Milan, Italy (A.L.). Department of Translational Medicine and Surgery, Università Cattolica del Sacro Cuore, Rome, Italy (E.D.C.). Maria Cecilia Hospital, Cotignola, Italy (E.T.).

Acknowledgments

The authors would like to thank Dr Federica Fumoso for the helpful discussions on the manuscript. M. Camera designed the research and provided financial resources; M. Brambilla and M. Camera interpreted the data and wrote the manuscript; P. Canzano, M. Conti, A. Becchetti, and A. Loffreda performed experiments

and analyzed data; A. Bonomi analyzed the data; N. Cosentino, E. De Candia, G.M. Podda, M. Cattaneo, D. Trabattoni, and J. Campodonico enrolled patients; G.E. Rovati, P.L.A. Giesen, C. Pinna, E. Tremoli, and M. Camera contributed to manuscript preparation; and all authors reviewed and approved the manuscript before submission.

Sources of Funding

This study was supported by research funding from Fondazione Monzino and from the Italian Ministry of Health (RC2017 Bio014 to M. Camera).

Disclosures

M. Camera, M. Brambilla, and P. Canzano are inventors of a patent application covering the work described in the study. The other authors report no conflicts.

Supplemental Material

Expanded Materials & Methods
Tables S1 and S2
Figures S1–S8

REFERENCES

- Drake TA, Morrissey JH, Edgington TS. Selective cellular expression of tissue factor in human tissues. Implications for disorders of hemostasis and thrombosis. *Am J Pathol*. 1989;134:1087–1097.
- Camera M, Frigerio M, Toschi V, Brambilla M, Rossi F, Cottell DC, Maderna P, Parolari A, Bonzi R, De Vincenti O, et al. Platelet activation induces cell-surface immunoreactive tissue factor expression, which is modulated differently by antiplatelet drugs. *Arterioscler Thromb Vasc Biol*. 2003;23:1690–1696. doi: 10.1161/01.ATV.0000085629.23209.AA
- Mann KG, Brummel K, Butenas S. What is all that thrombin for? *J Thromb Haemost*. 2003;1:1504–1514. doi: 10.1046/j.1538-7836.2003.00298.x
- Brummel KE, Paradis SG, Butenas S, Mann KG. Thrombin functions during tissue factor-induced blood coagulation. *Blood*. 2002;100:148–152. doi: 10.1182/blood.v100.1.148
- Falanga A, Marchetti M, Vignoli A, Balducci D, Russo L, Guerini V, Barbui T. V617f jak-2 mutation in patients with essential thrombocythemia: relation to platelet, granulocyte, and plasma hemostatic and inflammatory molecules. *Exp Hematol*. 2007;35:702–711. doi: 10.1016/j.exphem.2007.01.053
- Brambilla M, Camera M, Colnago D, Marenzi G, De Metrio M, Giesen PL, Balduini A, Veglia F, Gertow K, Biglioli P, et al. Tissue factor in patients with acute coronary syndromes: expression in platelets, leukocytes, and platelet-leukocyte aggregates. *Arterioscler Thromb Vasc Biol*. 2008;28:947–953. doi: 10.1161/ATVBAHA.107.161471
- Tilley RE, Holscher T, Belani R, Nieva J, Mackman N. Tissue factor activity is increased in a combined platelet and microparticle sample from cancer patients. *Thromb Res*. 2008;122:604–609. doi: 10.1016/j.thromres.2007.12.023
- Mayne E, Funderburg NT, Sieg SF, Asaad R, Kalinowska M, Rodriguez B, Schmaier AH, Stevens W, Lederman MM. Increased platelet and microparticle activation in HIV infection: upregulation of p-selectin and tissue factor expression. *J Acquir Immune Defic Syndr*. 2012;59:340–346. doi: 10.1097/QAI.0b013e3182439355
- Capozzi A, Manganelli V, Riitano G, Recalchi S, Truglia S, Alessandri C, Longo A, Garofalo T, Misasi R, Valesini G, et al. Tissue factor over-expression in platelets of patients with anti-phospholipid syndrome: induction role of anti-beta2-GPI antibodies. *Clin Exp Immunol*. 2019;196:59–66. doi: 10.1111/cei.13248
- Canzano P, Brambilla M, Porro B, Cosentino N, Tortorici E, Vicini S, Poggio P, Cascella A, Pengo MF, Veglia F, et al. Platelet and endothelial activation as potential mechanisms behind the thrombotic complications of COVID-19 patients. *JACC Basic Transl Sci*. 2021;6:202–218. doi: 10.1016/j.jaccbts.2020.12.009
- Gachet C, Hechler B, Leon C, Vial C, Leray C, Ohlmann P, Cazenave JP. Activation of ADP receptors and platelet function. *Thromb Haemost*. 1997;78:271–275.
- Dangelmaier C, Jin J, Smith JB, Kunapuli SP. Potentiation of thromboxane a₂-induced platelet secretion by G1 signaling through the phosphoinositide-3 kinase pathway. *Thromb Haemost*. 2001;85:341–348.
- Hechler B, Cattaneo M, Gachet C. The p2 receptors in platelet function. *Semin Thromb Hemost*. 2005;31:150–161. doi: 10.1055/s-2005-869520
- Hollopeter G, Jantzen HM, Vincent D, Li G, England L, Ramakrishnan V, Yang RB, Nurden P, Nurden A, Julius D, et al. Identification of the platelet ADP receptor targeted by anti-thrombotic drugs. *Nature*. 2001;409:202–207. doi: 10.1038/35051599

15. Leon C, Ravanat C, Freund M, Cazenave JP, Gachet C. Differential involvement of the p2y1 and p2y12 receptors in platelet procoagulant activity. *Arterioscler Thromb Vasc Biol*. 2003;23:1941–1947. doi: 10.1161/01.ATV.00000092127.16125.E6
16. Muller I, Klocke A, Alex M, Kotsch M, Luther T, Morgenstern E, Zieseniss S, Zahler S, Preissner K, Engelmann B. Intravascular tissue factor initiates coagulation via circulating microvesicles and platelets. *FASEB J*. 2003;17:476–478. doi: 10.1096/fj.02-0574fj
17. Leon C, Alex M, Klocke A, Morgenstern E, Moosbauer C, Eckly A, Spannagl M, Gachet C, Engelmann B. Platelet adp receptors contribute to the initiation of intravascular coagulation. *Blood*. 2004;103:594–600. doi: 10.1182/blood-2003-05-1385
18. Parodi G, Marcucci R, Valenti R, Gori AM, Migliorini A, Giusti B, Buonamici P, Gensini GF, Abbate R, Antoniucci D. High residual platelet reactivity after clopidogrel loading and long-term cardiovascular events among patients with acute coronary syndromes undergoing pci. *JAMA*. 2011;306:1215–1223. doi: 10.1001/jama.2011.1332
19. Larocca LM, Heller PG, Podda G, Pujol-Moix N, Glembotsky AC, Pecci A, Alberelli MA, Balduini CL, Landolfi R, Cattaneo M, et al. Megakaryocytic emperipolesis and platelet function abnormalities in five patients with gray platelet syndrome. *Platelets*. 2015;26:751–757. doi: 10.3109/09537104.2014.994093
20. De Candia E, Pecci A, Ciabattini G, De Cristofaro R, Rutella S, Yao-Wu Z, Lazzareschi I, Landolfi R, Coughlin S, Balduini CL. Defective platelet responsiveness to thrombin and protease-activated receptors agonists in a novel case of gray platelet syndrome: correlation between the platelet defect and the alpha-granule content in the patient and four relatives. *J Thromb Haemost*. 2007;7:551–559. doi: 10.1111/j.1538-7836.2007.02329.x
21. Cattaneo M, Lecchi A, Randi AM, McGregor JL, Mannucci PM. Identification of a new congenital defect of platelet function characterized by severe impairment of platelet responses to adenosine diphosphate. *Blood*. 1992;80:2787–2796.
22. Brambilla M, Rossetti L, Zara C, Canzano P, Giesen PLA, Tremoli E, Camera M. Do methodological differences account for the current controversy on tissue factor expression in platelets? *Platelets*. 2018;29:406–414. doi: 10.1080/09537104.2017.1327653
23. Camera M, Brambilla M, Toschi V, Tremoli E. Tissue factor expression on platelets is a dynamic event. *Blood*. 2010;116:5076–5077. doi: 10.1182/blood-2010-09-307306
24. Liang Y, Johnston M, Hirsh J, Pare G, Li C, Mehta S, Teo KK, Sloane D, Yi Q, Zhu J, et al. Relation between clopidogrel active metabolite levels and different platelet aggregation methods in patients receiving clopidogrel and aspirin. *J Thromb Thrombolysis*. 2012;34:429–436. doi: 10.1007/s11239-012-0762-2
25. Jeong YH, Bliden KP, Tantry US, Gurbel PA. High on-treatment platelet reactivity assessed by various platelet function tests: is the consensus-defined cut-off of vasp-p platelet reactivity index too low? *J Thromb Haemost*. 2012;10:487–489. doi: 10.1111/j.1538-7836.2011.04604.x
26. Hermann A, Rauch BH, Braun M, Schror K, Weber AA. Platelet CD40 ligand (CD40L)—subcellular localization, regulation of expression, and inhibition by clopidogrel. *Platelets*. 2001;12:74–82. doi: 10.1080/09537100020031207
27. Addo JB, Bray PF, Grigoryev D, Faraday N, Goldschmidt-Clermont PJ. Surface recruitment but not activation of integrin alpha IIb beta 3 (GPIIb-IIIa) requires a functional actin cytoskeleton. *Arterioscler Thromb Vasc Biol*. 1995;15:1466–1473. doi: 10.1161/01.atv.15.9.1466
28. Quinton TM, Murugappan S, Kim S, Jin J, Kunapuli SP. Different g protein-coupled signaling pathways are involved in alpha granule release from human platelets. *J Thromb Haemost*. 2004;2:978–984. doi: 10.1111/j.1538-7836.2004.00741.x
29. Ollivier V, Wang J, Manly D, Machlus KR, Wolberg AS, Jandrot-Perrus M, Mackman N. Detection of endogenous tissue factor levels in plasma using the calibrated automated thrombogram assay. *Thromb Res*. 2010;125:90–96. doi: 10.1016/j.thromres.2009.03.003
30. Versteeg HH, Ruf W. Tissue factor coagulant function is enhanced by protein-disulfide isomerase independent of oxidoreductase activity. *J Biol Chem*. 2007;282:25416–25424. doi: 10.1074/jbc.M702410200
31. Popielarski M, Ponamarczuk H, Stasiak M, Gdula A, Bednarek R, Wolska N, Swiatkowska M. P2y12 receptor antagonists and AR receptor agonists regulates protein disulfide isomerase secretion from platelets and endothelial cells. *Biochem Biophys Res Commun*. 2020;526:756–763. doi: 10.1016/j.bbrc.2020.03.143
32. Heemskerck JW, Siljander PR, Bevers EM, Farndale RW, Lindhout T. Receptors and signalling mechanisms in the procoagulant response of platelets. *Platelets*. 2000;11:301–306. doi: 10.1080/09537100050144704
33. Dorsam RT, Tuluc M, Kunapuli SP. Role of protease-activated and ADP receptor subtypes in thrombin generation on human platelets. *J Thromb Haemost*. 2004;2:804–812. doi: 10.1111/j.1538-7836.2004.00692.x
34. Trabattini D, Brambilla M, Canzano P, Becchetti A, Teruzzi G, Porro B, Fiorelli S, Muratori M, Tedesco CC, Veglia F, et al. Migraine in patients undergoing PFO closure: characterization of a platelet-associated pathophysiological mechanism: the learner study. *JACC Basic Transl Sci*. 2022;7:525–540. doi: 10.1016/j.jaccbts.2022.02.002
35. Benjamin EJ, Muntner P, Alonso A, Bittencourt MS, Callaway CW, Carson AP, Chamberlain AM, Chang AR, Cheng S, Das SR, et al; American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics-2019 update—a report from the American Heart Association. *Circulation*. 2019;139:e56–e528. doi: 10.1161/CIR.0000000000000659
36. Judge HM, Buckland RJ, Sugidachi A, Jakubowski JA, Storey RF. The active metabolite of prasugrel effectively blocks the platelet p2y12 receptor and inhibits procoagulant and pro-inflammatory platelet responses. *Platelets*. 2008;19:125–133. doi: 10.1080/09537100701694144
37. Chyrchel B, Drozd A, Dlugosz D, Stepień EL, Surdacki A. Platelet reactivity and circulating platelet-derived microvesicles are differently affected by p2y(12) receptor antagonists. *Int J Med Sci*. 2019;16:264–275. doi: 10.7150/ijms.28580
38. Gasecka A, Nieuwland R, Budnik M, Dignat-George F, Eyleten C, Harrison P, Lacroix R, Leroyer A, Opolski G, Pluta K, et al. Ticagrelor attenuates the increase of extracellular vesicle concentrations in plasma after acute myocardial infarction compared to clopidogrel. *J Thromb Haemost*. 2020;16:609–623. doi: 10.1111/jth.14689
39. Braun OO, Johnell M, Varenhorst C, James S, Brandt JT, Jakubowski JA, Winters KJ, Wallentin L, Erlinge D, Siegbahn A. Greater reduction of platelet activation markers and platelet-monocyte aggregates by prasugrel compared to clopidogrel in stable coronary artery disease. *Thromb Haemost*. 2008;100:626–633.
40. Wiviott SD, Braunwald E, McCabe CH, Montalescot G, Ruzyllo W, Gottlieb S, Neumann F-J, Ardissino D, De Servi S, Murphy SA, et al; TRITON-TIMI 38 Investigators. Prasugrel versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med*. 2007;357:2001–2015. doi: 10.1056/NEJMoa0706482
41. Gurbel PA, Bliden KP, Butler K, Tantry US, Gesheff T, Wei C, Teng R, Antonino MJ, Patil SB, Karunakaran A, et al. Randomized double-blind assessment of the onset and offset of the antiplatelet effects of ticagrelor versus clopidogrel in patients with stable coronary artery disease: the onset/offset study. *Circulation*. 2009;120:2577–2585. doi: 10.1161/CIRCULATIONAHA.109.912550
42. Aradi D, Tornóyos A, Pinter T, Vorobcsuk A, Konyi A, Falukozy J, Veress G, Magyar B, Horváth IG, Komócsi A. Optimizing p2y12 receptor inhibition in patients with acute coronary syndrome on the basis of platelet function testing: impact of prasugrel and high-dose clopidogrel. *J Am Coll Cardiol*. 2014;63:1061–1070. doi: 10.1016/j.jacc.2013.12.023
43. Mayer K, Schulz S, Bernlochner I, Morath T, Braun S, Hausleiter J, Massberg S, Schunkert H, Laugwitz KL, Kastrati A, et al. A comparative cohort study on personalised antiplatelet therapy in pci-treated patients with high on-clopidogrel platelet reactivity. Results of the ISAR-HPR registry. *Thromb Haemost*. 2014;112:342–351. doi: 10.1160/TH13-10-0874
44. Price MJ, Berger PB, Teirstein PS, Tanguay JF, Angiolillo DJ, Spriggs D, Puri S, Robbins M, Garratt KN, Bertrand OF, et al; GRAVITAS Investigators. Standard- vs high-dose clopidogrel based on platelet function testing after percutaneous coronary intervention: the gravitas randomized trial. *JAMA*. 2011;305:1097–1105. doi: 10.1001/jama.2011.290
45. Collet JP, Cuisset T, Range G, Cayla G, Elhadad S, Pouillot C, Henry P, Motreff P, Carrié D, Boueri Z, et al; ARCTIC Investigators. Bedside monitoring to adjust antiplatelet therapy for coronary stenting. *N Engl J Med*. 2012;367:2100–2109. doi: 10.1056/NEJMoa1209979
46. Wallentin L, Becker RC, Budaj A, Cannon CP, Emanuelsson H, Held C, Horrow J, Husted S, James S, Katus H, et al; PLATO Investigators. Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med*. 2009;361:1045–1057. doi: 10.1056/NEJMoa0904327
47. Kotsia A, Brilakis ES, Held C, Cannon C, Steg GP, Meier B, Cools F, Claeys MJ, Cornel JH, Aylward P, et al. Extent of coronary artery disease and outcomes after ticagrelor administration in patients with an acute coronary syndrome: insights from the platelet inhibition and patient outcomes (PLATO) trial. *Am Heart J*. 2014;168:68–75.e2. doi: 10.1016/j.ahj.2014.04.001
48. Judge HM, Buckland RJ, Sugidachi A, Jakubowski JA, Storey RF. Relationship between degree of p2y12 receptor blockade and inhibition of p2y12-mediated platelet function. *Thromb Haemost*. 2010;103:1210–1217. doi: 10.1160/TH09-11-0770

49. Reny JL, Berdague P, Poncet A, Barazer I, Nolli S, Fabbro-Peray P, Schved J-F, Bounameaux H, Mach F, de Moerloose P, et al; Antiplatelet Drug Resistances and Ischemic Events (ADRIE) Study Group. Antiplatelet drug response status does not predict recurrent ischemic events in stable cardiovascular patients: results of the antiplatelet drug resistances and ischemic events study. *Circulation*. 2012;125:3201–3210. doi: 10.1161/CIRCULATIONAHA.111.085464
50. Trenk D, Stone GW, Gawaz M, Kastrati A, Angiolillo DJ, Muller U, Richardt G, Jakubowski JA, Neumann FJ. A randomized trial of prasugrel versus clopidogrel in patients with high platelet reactivity on clopidogrel after elective percutaneous coronary intervention with implantation of drug-eluting stents: results of the trigger-PCI (testing platelet reactivity in patients undergoing elective stent placement on clopidogrel to guide alternative therapy with prasugrel) study. *J Am Coll Cardiol*. 2012;59:2159–2164. doi: 10.1016/j.jacc.2012.02.026
51. Montalescot G, Range G, Silvain J, Bonnet JL, Boueri Z, Barthelemy O, Cayla G, Belle L, Van Belle E, Cuisset T, et al; ARCTIC Investigators. High on-treatment platelet reactivity as a risk factor for secondary prevention after coronary stent revascularization: a landmark analysis of the ARCTIC study. *Circulation*. 2014;129:2136–2143. doi: 10.1161/CIRCULATIONAHA.113.007524
52. Gurbel PA, Bliden KP, Guyer K, Cho PW, Zaman KA, Kreutz RP, Bassi AK, Tantry US. Platelet reactivity in patients and recurrent events post-stenting: results of the prepare post-stenting study. *J Am Coll Cardiol*. 2005;46:1820–1826. doi: 10.1016/j.jacc.2005.07.041
53. Gurbel PA, Bliden KP, Navickas I, Cohen E, Tantry US. Post-coronary intervention recurrent ischemia in the presence of adequate platelet inhibition by dual antiplatelet therapy: what are we overlooking? *J Thromb Haemost*. 2007;5:2300–2301. doi: 10.1111/j.1538-7836.2007.02749.x
54. Capodanno D, Alfonso F, Levine GN, Valgimigli M, Angiolillo DJ. ACC/AHA versus ESC guidelines on dual antiplatelet therapy: JACC guideline comparison. *J Am Coll Cardiol*. 2018;72:2915–2931. doi: 10.1016/j.jacc.2018.09.057
55. Mega JL, Braunwald E, Mohanavelu S, Burton P, Poulter R, Misselwitz F, Hricak V, Barnathan ES, Bordes P, Witkowski A, et al; ATLAS ACS-TIMI 46 Study Group. Rivaroxaban versus placebo in patients with acute coronary syndromes (ATLAS ACS-TIMI 46): a randomised, double-blind, phase II trial. *Lancet*. 2009;374:29–38. doi: 10.1016/S0140-6736(09)60738-8
56. Eikelboom JW, Connolly SJ, Bosch J, Dagenais GR, Hart RG, Shestakovska O, Diaz R, Alings M, Lonn EM, Anand SS, et al; COMPASS Investigators. Rivaroxaban with or without aspirin in stable cardiovascular disease. *N Engl J Med*. 2017;377:1319–1330. doi: 10.1056/NEJMoa1709118
57. Pieters J, Lindhout T, Hemker HC. In situ-generated thrombin is the only enzyme that effectively activates factor VIII and factor V in thromboplastin-activated plasma. *Blood*. 1989;74:1021–1024.
58. Monroe DM, Hoffman M, Roberts HR. Platelets and thrombin generation. *Arterioscler Thromb Vasc Biol*. 2002;22:1381–1389. doi: 10.1161/01.atv.0000031340.68494.34
59. Merlini PA, Bauer KA, Oltrona L, Ardissino D, Cattaneo M, Belli C, Mannucci PM, Rosenberg RD. Persistent activation of coagulation mechanism in unstable angina and myocardial infarction. *Circulation*. 1994;90:61–68. doi: 10.1161/01.cir.90.1.61
60. Nidorf SM, Fiolet ATL, Mosterd A, Eikelboom JW, Schut A, Opstal TSJ, The SHK, Xu X-F, Ireland MA, Lenderink T, et al; LoDoCo2 Trial Investigators. Colchicine in patients with chronic coronary disease. *N Engl J Med*. 2020;383:1838–1847. doi: 10.1056/NEJMoa2021372
61. Opstal TSJ, Fiolet ATL, van Broekhoven A, Mosterd A, Eikelboom JW, Nidorf SM, Thompson PL, Duyvendak M, van Eck JWM, van Beek EA, et al; LoDoCo2 Trial Investigators. Colchicine in patients with chronic coronary disease in relation to prior acute coronary syndrome. *J Am Coll Cardiol*. 2021;78:859–866. doi: 10.1016/j.jacc.2021.06.037
62. Tong DC, Quinn S, Nasis A, Hiew C, Roberts-Thomson P, Adams H, Sriamaseswaran R, Htun NM, Wilson W, Stub D, et al. Colchicine in patients with acute coronary syndrome: the Australian Cops Randomized Clinical Trial. *Circulation*. 2020;142:1890–1900. doi: 10.1161/CIRCULATIONAHA.120.050771
63. Bresson D, Roubille F, Prieur C, Biere L, Ivanov F, Bouletti C, Dubreuil O, Rioufol G, Boutitie F, Sideris G, et al. Colchicine for left ventricular infarct size reduction in acute myocardial infarction: a phase II, multicenter, randomized, double-blinded, placebo-controlled study protocol - the COVERT-MI study. *Cardiology*. 2021;146:151–160. doi: 10.1159/000512772
64. Hennessy T, Soh L, Bowman M, Kurup R, Schultz C, Patel S, Hillis GS. The low dose colchicine after myocardial infarction (LoDoCo-MI) study: a pilot randomized placebo controlled trial of colchicine following acute myocardial infarction. *Am Heart J*. 2019;215:62–69. doi: 10.1016/j.ahj.2019.06.003
65. Tardif JC, Kouz S, Waters DD, Bertrand OF, Diaz R, Maggioni AP, Pinto FJ, Ibrahim R, Gamra H, Kiwan GS, et al. Efficacy and safety of low-dose colchicine after myocardial infarction. *N Engl J Med*. 2019;381:2497–2505. doi: 10.1056/NEJMoa1912388
66. Tong DC, Bloom JE, Quinn S, Nasis A, Hiew C, Roberts-Thomson P, Adams H, Sriamaseswaran R, Htun NM, Wilson W, et al. Colchicine in patients with acute coronary syndrome: two-year follow-up of the Australian Cops Randomized Clinical Trial. *Circulation*. 2021;144:1584–1586. doi: 10.1161/CIRCULATIONAHA.121.054610
67. Cimmino G, Tarallo R, Conte S, Morello A, Pellegrino G, Loffredo FS, Cali G, De Luca N, Golino P, Trimarco B, et al. Colchicine reduces platelet aggregation by modulating cytoskeleton rearrangement via inhibition of cofilin and lim domain kinase 1. *Vascul Pharmacol*. 2018;111:62–70. doi: 10.1016/j.vph.2018.09.004
68. Shah B, Allen N, Harchandani B, Pillinger M, Katz S, Sedlis SP, Echagarraga C, Samuels SK, Morina P, Singh P, et al. Effect of colchicine on platelet-platelet and platelet-leukocyte interactions: a pilot study in healthy subjects. *Inflammation*. 2016;39:182–189. doi: 10.1007/s10753-015-0237-7
69. Raju NC, Yi Q, Nidorf M, Fagel ND, Hiralal R, Eikelboom JW. Effect of colchicine compared with placebo on high sensitivity C-reactive protein in patients with acute coronary syndrome or acute stroke: a pilot randomized controlled trial. *J Thromb Thrombolysis*. 2012;33:88–94. doi: 10.1007/s11239-011-0637-y