

Original Research Article

Association between AXL, Hippo transducers and survival outcomes in male breast cancer.[†]

Running head: AXL, Hippo and male breast cancer

Anna Di Benedetto^{1,*}, Marcella Mottolise^{1,*}, Francesca Sperati², Cristiana Ercolani¹, Luigi Di Lauro³, Laura Pizzuti³, Patrizia Vici³, Irene Terrenato², Abeer M. Shaaban⁴, Matthew P. Humphries⁵, Sreekumar Sundara-Rajan⁵, Maddalena Barba^{3,6}, Valerie Speirs^{5,*}, Ruggero De Maria⁷, and Marcello Maugeri-Saccà^{3,6,*}

¹ Department of Pathology, “Regina Elena” National Cancer Institute, Via Elio Chianesi 53, 00144, Rome, Italy

² Biostatistics-Scientific Direction, “Regina Elena” National Cancer Institute, Via Elio Chianesi 53, 00144, Rome, Italy

³ Division of Medical Oncology 2, “Regina Elena” National Cancer Institute, Via Elio Chianesi 53, 00144, Rome, Italy

⁴ Queen Elizabeth Hospital Birmingham and University of Birmingham, Department of Histopathology, Edgbaston, Birmingham B15 2GW, UK

⁵ Leeds Institute of Cancer and Pathology, Wellcome Trust Brenner Building, University of Leeds, Leeds LS9 7TF, UK.

⁶ Scientific Direction, “Regina Elena” National Cancer Institute, Via Elio Chianesi 53, 00144, Rome, Italy

⁷ Institute of General Pathology, Catholic University of the Sacred Heart, Largo Agostino Gemelli, 10, 00168, Rome

* **Corresponding author:** Dr. Marcello Maugeri-Saccà, Division of Medical Oncology B and Scientific Direction, “Regina Elena” National Cancer Institute, Via Elio Chianesi 53, 00144 Rome, Italy. Tel. +39-06-52662724, E-mail: maugeri.marcello@gmail.com; Prof Valerie Speirs, Leeds Institute of Cancer and Pathology, Wellcome Trust Brenner Building, University of Leeds, Leeds LS9 7TF, UK. Tel. 0113 343 8633, e-mail: v.speirs@leeds.ac.uk

[†]This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/jcp.25745]

Received 7 November 2016; Revised 5 December 2016; Accepted 15 December 2016

Journal of Cellular Physiology

This article is protected by copyright. All rights reserved

DOI 10.1002/jcp.25745

Keywords:

- Male Breast Cancer
- AXL
- TAZ
- YAP
- Hippo Pathway

Funding

This study was supported by the “Associazione Italiana per la Ricerca sul Cancro” (AIRC IG Grant N:13431 to RDM), by Breast Cancer Now (formerly Breast Cancer Campaign; grant 2007MayPR02 to VS and AMS), and Yorkshire Cancer Research (grant L378 to VS).

Abstract

Male breast cancer (MBC) is an uncommon malignancy. We have previously reported that the expression of the Hippo transducers TAZ/YAP and their target CTGF was associated with inferior survival in MBC patients. Preclinical evidence demonstrated that Axl is a transcriptional target of TAZ/YAP. Thus, we herein assessed AXL expression to further investigate the significance of active TAZ/YAP-driven transcription in MBC. For this study, 255 MBC samples represented in tissue microarrays were screened for AXL expression, and 116 patients were included. The association between categorical variables was verified by the Pearson's Chi-squared test of independence (2-tailed) or the Fisher Exact test. The relationship between continuous variables was tested with the Pearson's correlation coefficient. The Kaplan-Meier method was used for estimating survival curves, which were compared by log-rank test. Factors potentially impacting 10-year and overall survival were verified in Cox proportional regression models. AXL was positively associated with the TAZ/CTGF and YAP/CTGF phenotypes ($p=0.001$ and $p=0.002$, respectively). Patients with TAZ/CTGF/AXL- or YAP/CTGF/AXL-expressing tumors had inferior survival compared with non-triple-positive patients (log rank $p=0.042$ and $p=0.048$, respectively). The variables TAZ/CTGF/AXL and YAP/CTGF/AXL were adverse factors for 10-year survival in the multivariate Cox models (HR 2.31, 95%CI:1.02-5.22, $p=0.045$, and HR 2.27, 95%CI:1.00-5.13, $p=0.050$). Nearly comparable results were obtained from multivariate analyses of overall survival. The expression pattern of AXL corroborates the idea of the detrimental role of TAZ/YAP activation in MBC. Overall, Hippo-linked biomarkers deserve increased attention in this rare disease. This article is protected by copyright. All rights reserved

INTRODUCTION

Male breast cancer (MBC) is a rare disease often arising in elderly men (Ruddy and Winer, 2013). Despite some recent molecular characterization efforts, its biology remains understudied (Piscuoglio, 2016; Johansson 2013; Kornegoor, 2012; Johansson, 2011; Callari, 2011; Johansson, 2012; Fassan, 2009). Indeed, apart from hormone receptors that are frequently expressed (Anderson 2010; Shaaban, 2012), and whose targeting represents the mainstay of medical treatment (Doyen, 2010; Eggemann, 2013; Zagouri, 2013a; Zagouri, 2013b; Di Lauro, 2013; Di Lauro, 2014; Di Lauro, 2015), little is known about the nature of deregulated molecular networks underlying the disease. Coherent with the paucity of data available so far, both in terms of biological processes and therapeutic targets, the European Organization for Research and Treatment (EORTC) has launched an initiative dedicated toward obtaining a better understanding of the disease (EORTC 10085, available at: <http://www.eortc.org/research-groups/breast-cancer-group/ongoing-and-future-projects/>).

Over the past years, an Italy-UK collaboration was set off with the aim of investigating specific oncogenic pathways in MBC, and their relationship with clinical-pathological features and survival outcomes. On the basis of our preclinical experience, the focus was placed on signals connected with cancer stem cells (CSCs), a rare cellular subset whose hallmarks include self-renewal, ability to propagate the tumor in animal models, and resistance to current anticancer therapies (Beck and Blanpain, 2013). The evolutionary conserved Hippo signaling pathway was chosen for proof-of-principle studies, considering that both our group and others demonstrated that TAZ/YAP-mediated gene transcription is associated with the retention/acquisition of stem-like features, self-renewal, epithelial-mesenchymal transition, chemoresistance and metastatic dissemination in breast CSC (BCSC) models (Bartucci, 2015; Cordenonsi, 2011; Xiang, 2014; Chang, 2015; Nandy, 2015). Moreover, our proof-of-concept clinical studies connected the expression of

TAZ/YAP to inferior therapeutic and survival outcomes in female breast cancer patients who received neoadjuvant therapy (Vici, 2014; Vici, 2016).

Hippo is a “two face” pathway, composed by two modules that, in a neoplastic background, elicit opposite biological outcomes (Johnson and Halder, 2014). While core kinases (MST1/2 and LATS1/2) and adaptors (SAV1 and MOB1A/1B) act as tumor suppressors by mediating nuclear exclusion, cytoplasmic retention and proteasomal degradation of the Hippo transducers TAZ/YAP, these, together with their DNA binding platform chiefly consisting in TEAD factors, have been extensively connected with tumor-promoting functions (Johnson and Halder, 2014). Many tumors exhibit defective activation of core Hippo kinases/adaptors, and/or activation of upstream or lateral signals that, albeit not canonically placed within the pathway architecture, can directly activate TAZ/YAP (Piccolo, 2014).

In MBC, we have previously reported that the expression of TAZ/YAP and their established target CTGF conferred inferior survival outcomes (Di Benedetto, 2016a). To provide further insights into the oncogenic role of TAZ/YAP in such an uncommon tumor, we herein assessed AXL, a member of the TAM family of receptor tyrosine kinases (RTKs) (Graham, 2014), as the *Axl* gene is a direct target of the Hippo transcriptional machinery (Azzolin, 2014; Zanconato, 2015). In turn, compelling evidence demonstrated that AXL mediates a variety of oncogenic functions, spanning from cancer cell survival to therapeutic resistance (Graham, 2014), and AXL expression has been reported in a number of hematological and epithelial malignancies (Wu, 2014). Overall, the present study was designed with the goals of i) describing the expression pattern of AXL, and ii) providing further ground on the clinical potential of Hippo-related biomarkers in MBC.

MATERIALS AND METHODS

Study Participants and procedures

For this retrospective analysis samples from 255 histologically confirmed, non-metastatic MBC patients, represented in tissue microarrays (TMAs) (Shaaban, 2012), were immunostained for evaluating the expression of AXL. Eligibility was assessed based on the following criteria: i) complete data on AXL, ii) complete data on TAZ, YAP and CTGF, iii) complete data on the following routine clinical-pathological features: histology, tumor grade, hormone receptors (ER, PgR) and Ki-67, and iv) availability of survival data. On this basis, 116 patients were included. Nodal status was not considered among the inclusion criteria, given that this information was available for 87 patients only. As already specified, tamoxifen was the most commonly administered agent in the adjuvant setting, albeit complete information pertinent to post-surgical therapy were not available (Shaaban, 2012). This study was conducted in accordance with the Declaration of Helsinki and approved by the Institutional Review Board of the “Regina Elena” National Cancer Institute of Rome and by the Leeds (East) Research Ethics Committee (06/Q1205/156). Samples were anonymised to the research team and informed consent was not required (Shaaban, 2012). This study adheres to the REMARKS guidelines (McShane; 2006).

Immunohistochemistry

TMAs were built from formalin-fixed paraffin-embedded (FFPE) material (Shaaban, 2012). The immunohistochemical assessment of AXL was performed using the polyclonal antibody anti-AXL (HPA037422, Sigma-Aldrich, St. Louis, USA) at the dilution of 1:200. AXL expression was determined both in terms of staining intensity on a four-grade scale (0: negative, 1+: weak, 2+: moderate, 3+: strong) and percentage of tumor-expressing cells (0-100%). Samples were considered positive if $\geq 30\%$ of neoplastic cells exhibited a distinct

membranous/cytoplasmic immunoreactivity of any intensity. This cutoff reflected the lowest percentage of AXL-expressing cells observed in this cohort (range 30-90%). Two investigators (ADB and MM) evaluated immunoreactivity. The modalities used for assessing TAZ, YAP, CTGF, hormone receptors and Ki-67 levels were detailed elsewhere (Shaaban, 2012; Di Benedetto, 2016a).

Statistical analysis

The relationship between categorical variables was assessed with the Pearson's Chi-squared test of independence (2-tailed) or the Fisher Exact test. The Pearson's correlation coefficient was used to establish the correlation between continuous variables. Survival curves were estimated and compared with the Kaplan-Meier method and log-rank test, respectively. Overall survival (OS) was calculated as the time from diagnosis to death due to any cause. For 10-year survival analysis, patients who experienced the outcome of interest after this time point were censored. Potential predictors of 10-year and overall survival were verified in uni- and multivariate Cox proportional regression models. The final multivariate models were built by adjusting for standard clinical-molecular variables (invasive ductal carcinoma (IDC)/invasive lobular carcinoma (ILC) vs other histotypes, G3 vs G1-2, ER+/PgR+ vs other, Ki-67 high vs Ki-67 low) independently on whether these factors were significant at the univariate assessment. Statistical significance was set at $p < 0.05$. Statistical analyses were carried out using SPSS software (SPSS version 21, SPSS Inc., Chicago, IL, USA).

RESULTS

Association between AXL expression and clinical-pathological factors

For this study, 255 MBC samples were screened for the expression of AXL and, based on the aforementioned eligibility criteria, 116 patients were included in the final analysis.

Representative immunohistochemical staining is presented in Figure 1. Baseline characteristics of the patients who met the eligibility criteria are summarized in Table 1. As shown in Table 2, we did not observe any significant association between AXL and standard clinical-pathological factors, namely histology, nodal status, tumor grade, hormone receptor status, Ki-67. Conversely, AXL was positively associated with the TAZ/CTGF and YAP/CTGF phenotypes (χ^2 $p=0.001$ and $p=0.002$, respectively, as reported in Table 2). Moreover, a significant positive correlation, in terms of percentage of tumor-expressing cells, was seen between AXL and CTGF (Pearson's correlation coefficient=0.285; $p=0.002$, data available upon request). Overall, these data support our hypothesis of active TAZ/YAP-driven gene transcription in a subset of MBC patients.

Relationship between the TAZ/CTGF/AXL and YAP/CTGF/AXL phenotypes and clinical outcomes

Patients whose tumors had a triple-positive phenotype (TAZ/CTGF/AXL or YAP/CTGF/AXL) experienced shorter survival compared with their negative counterparts (log rank $p=0.042$ and $p=0.048$, respectively. Figure 2, panel A and B), analogously to the TAZ/CTGF and YAP/CTGF models that we previously connected with inferior survival outcomes. Results from the univariate regression analyses, presented in Table 3, performed for identifying variables impacting 10-year survival confirmed that the TAZ/CTGF/AXL and YAP/CTGF/AXL phenotypes are adverse prognostic indicators (HR 2.37, 95%CI: 1.05-5.34, $p=0.038$, and HR 2.31, 95%CI:1.02-5.22, $p=0.044$, respectively). The multivariate Cox models (Table 3) confirmed these data (TAZ/CTGF/AXL: HR 2.31, 95%CI: 1.02-5.22, $p=0.045$. YAP/CTGF/AXL: HR 2.27, 95%CI:1.00-5.13, $p=0.050$). Nearly comparable results, even though with a trend toward statistical significance, were obtained in the

Accepted Article

multivariate Cox models for overall survival (TAZ/CTGF/AXL: HR 2.19, 95%CI: 0.97-4.94, p=0.058. YAP/CTGF/AXL: HR 2.16, 95%CI:0.96-4.87, p=0.063) (Table 4).

DISCUSSION

The present study aimed at providing further evidence supporting the negative prognostic significance of active TAZ/YAP-mediated gene transcription in MBC patients. To this end, 116 MBC samples, previously characterized for the expression of TAZ, YAP and CTGF (Di Benedetto, 2016a), were immunostained for AXL, another established target of the Hippo transducers. The message conveyed by the present study is that: i) AXL expression is significantly associated with the TAZ/CTGF and YAP/CTGF phenotypes, and positively correlated with CTGF, an established TAZ/YAP target gene, ii) patients whose tumors harbored the TAZ/CTGF/AXL and YAP/CTGF/AXL molecular backgrounds had adverse survival outcomes compared with patients whose tumors did not show molecular evidence of TAZ/YAP activation, and iii) the prognostic significance of the triple-positive model herein investigated was comparable to that of the previously described TAZ/CTGF and YAP/CTGF models, further enforcing the hypothesis that deregulation of the Hippo machinery may represent an important source of oncogenic stimuli in MBC.

We acknowledge that, being retrospective in nature, our study has some limitations, chiefly the fact that we were unable to consider cancer-specific events owing to the lack of complete information related to the cause of death, as discussed elsewhere (Di Benedetto, 2016a). Nevertheless, ten-year survival is a suitable endpoint to overcome this drawback, when considering that MBC is a disease of elderly men and with a long natural history, comparable to that of luminal-type breast cancer arising in post-menopausal women (Ruddy and Winer, 2013). Overall, our analyses at an intermediate time point (10-year survival) conveyed the

message that deregulated Hippo pathway is involved in the genesis of MBC, analogously to earlier preclinical and clinical reports in female breast cancer (Maugeri-Saccà, 2015).

Next, our interest towards the Hippo pathway in MBC extends beyond the analysis of key pathway components. Indeed, we are currently analyzing other pathways intersecting the Hippo cascade at various levels. Bearing in mind the number of molecular signals interacting with the Hippo cascade (Piccolo, 2014), we focused our attention on the following: i) the mevalonate pathway and, ii) the DNA damage repair (DDR) machinery. Evidence of metabolic control of the Hippo pathway stemmed from a high-throughput screening of Food and Drug Administration (FDA)-approved compounds performed in the attempt of identifying TAZ/YAP modulators through a drug repositioning approach. This screening identified both hydrophilic and lipophilic statins, a class of widely prescribed cholesterol-lowering medications, as the most potent TAZ/YAP modulators. Mechanistically, it was demonstrated that the mevalonate pathway promotes TAZ/YAP activation via the production of geranylgeranyl pyrophosphate (GGPP), which is essential for correct membrane localization, and consequently activation, of Rho GTPases (Sorrentino, 2014). These, in turn, inhibit TAZ/YAP phosphorylation independently on the activity of core Hippo pathway kinases (LATS1/2). Interestingly, the effects of statins were reproduced with the administration of nitrogen-containing bisphosphonates, such as zoledronic acid, a class of compounds that inhibit farnesyl pyrophosphate synthase, another key enzyme in the mevalonate cascade (Sorrentino, 2014). Overall, the preclinically documented metabolic control of TAZ/YAP operated by the mevalonate pathway, together with the association between obesity and MBC, makes the mevalonate pathway an attractive therapeutic target in MBC (Humphries, 2015). Our results indicates that 3-Hydroxy-3-methylglutharylcoenzyme A reductase (HMG-CoAR), the rate-limiting enzyme of the mevalonate cascade and the

Accepted Article

target of statins (Buhaescu and Izzedine, 2007), is significantly co-expressed with Hippo transducers, suggesting that the mevalonate pathway regulates the activity of TAZ/YAP (Di Benedetto, 2016b). When reanalyzing this association in light of the data on AXL herein presented, we confirmed that HMG-CoAR is positively associated with both the TAZ/CTGF/AXL and YAP/CTGF/AXL phenotypes (Fisher Exact test $p=0.024$ and $p=0.014$, respectively). Regarding the DDR, a wave of preclinical evidence connected key orchestrators of the DDR network, namely ATM and ATR, with the Hippo pathway (Pefani DE and O'Neill, 2016). In a greater detail, Hippo pathway kinases are targeted by the ATM/Chk2 and ATR/Chk1 pathways, thus participating in an array of interconnected biological processes spanning from cell-cycle checkpoint activation and replication fork stability to DNA repair and apoptosis (Pefani DE and O'Neill, 2016). The molecular cooperation between the Hippo pathway and the molecular network safeguarding genome integrity is further enforced when considering the interactions between Hippo pathway components and p53. Indeed, it was demonstrated that LATS2 stabilizes p53 (Ganem NJ, 2014), and that LATS1/2 silencing shifted p53 function toward an oncogenic (gain-of-function), mutant-like state (Furth N, 2015). Moreover, YAP physically interacts with mutant p53 proteins promoting the aberrant expression of cell cycle-related genes (Di Agostino S, 2016). Our group initiated characterization of the ATM-Chk2 and ATR-Chk1 axes in MBC, with the aim of providing novel information on DNA repair-linked biomarkers and their connection with the Hippo machinery and metabolic avenues.

We believe that our efforts toward achieving a better comprehension of the molecular basis of MBC potentially hold important therapeutic implications. Indeed, inhibition of the Hippo transducers TAZ/YAP was preclinically obtained with various Food and Drug Administration- (FDA) approved compounds, such as statins, bisphosphonates, verteporfin, dobutamine, metformin, and dasatinib (Sorrentino, 2014; Frangou, 2014; Rosenbluh, 2012;

Liu-Chittenden, 2012; Yu, 2014; Bao, 2011; Wang, 2014; DeRan, 2014). In our opinion, agents targeting the mevalonate pathway (e.g. statins and nitrogen-containing bisphosphonates) and glucose-lowering compounds (e.g. metformin) are those for which it is plausible gathering retrospective, hypothesis-generating data. Indeed, a non-negligible fraction of MBC patients supposedly received these treatments for co-existing medical conditions, such as elevated cholesterol levels, osteoporosis and diabetes, since these comorbidities are fairly common in elderly men. Reconsidering survival outcomes of MBC patients in light of the use of these agents, target expression, and activation of the TAZ/YAP-mediated oncogenic program is a strategy to be pursued.

In summary, by analyzing AXL, an established target of TAZ/YAP widely exploited in the preclinical setting as a proxy of their activation, we herein provided further evidence to the detrimental role of activated Hippo transducers in MBC. Furthermore, ongoing investigations will shortly help elucidate the connection between Hippo and other oncogenic signals.

Acknowledgements

We thank members of the Male Breast Cancer Consortium for kindly contributing cases. We thank Tania Merlino for technical assistance.

Conflicts of interest

All authors disclose no conflict of interest.

Author contributions

VS, RDM and MM-S conceived the study and designed the study. ADB, MM, CE, AMS carried out molecular pathology analyses. FS, IT and MB performed statistical analyses. LDL, LP, PV, MPH and SS-R acquired the data related to clinical-pathological features and therapeutic outcomes. MM-S wrote the final version of the manuscript. All authors read and approved the final version of the manuscript and agree to be accountable for all aspects of the work.

References

Anderson WF, Jatoi I, Tse J, Rosenberg PS. 2010. Male breast cancer: a population-based comparison with female breast cancer. *J Clin Oncol* 28:232-239

Azzolin L, Panciera T, Soligo S, Enzo E, Bicciato S, Dupont S, Bresolin S, Frasson C, Basso G, Guzzardo V, Fassina A, Cordenonsi M, Piccolo S. 2014. YAP/TAZ incorporation in the β -catenin destruction complex orchestrates the Wnt response. *Cell* 158:157-170

Bao Y, Nakagawa K, Yang Z, Ikeda M, Withanage K, Ishigami-Yuasa M, Okuno Y, Hata S, Nishina H, Hata Y. 2011. A cell-based assay to screen stimulators of the Hippo pathway reveals the inhibitory effect of dobutamine on the YAP-dependent gene transcription. *J Biochem* 150:199-208

Bartucci M, Dattilo R, Moriconi C, Pagliuca A, Mottolise M, Federici G, Benedetto AD, Todaro M, Stassi G, Sperati F, Amabile MI, Pillozzi E, Patrizii M, Biffoni M, Maugeri-Saccà M, Piccolo S, De Maria R. 2015. TAZ is required for metastatic activity and chemoresistance of breast cancer stem cells. *Oncogene* 34:681-690

Beck B, Blanpain C. 2013. Unravelling cancer stem cell potential. *Nat Rev Cancer* 13:727-738

Buhaescu I, Izzedine H. 2007. Mevalonate pathway: a review of clinical and therapeutical implications. *Clin Biochem* 40:575-584

Callari M, Cappelletti V, De Cecco L, Musella V, Miodini P, Veneroni S, Gariboldi M, Pierotti MA, Daidone MG. 2011. Gene expression analysis reveals a different transcriptomic landscape in female and male breast cancer. *Breast Cancer Res Treat* 127: 601-610

Chang C, Goel HL, Gao H, Pursell B, Shultz LD, Greiner DL, Ingerpau S, Patarroyo M, Cao S, Lim E, Mao J, McKee KK, Yurchenco PD, Mercurio AM. 2015. A laminin 511 matrix is

regulated by TAZ and functions as the ligand for the $\alpha 6\beta 1$ integrin to sustain breast cancer stem cells. *Genes Dev* 29:1-6

Cordenonsi M, Zanconato F, Azzolin L, Forcato M, Rosato A, Frasson C, Inui M, Montagner M, Parenti AR, Poletti A, Daidone MG, Dupont S, Basso G, Bicciato S, Piccolo S. 2011. The Hippo transducer TAZ confers cancer stem cell-related traits on breast cancer cells. *Cell*. 147:759-772

DeRan M, Yang J, Shen CH, Peters EC, Fitamant J, Chan P, Hsieh M, Zhu S, Asara JM, Zheng B, Bardeesy N, Liu J, Wu X. 2014. Energy stress regulates hippo-YAP signaling involving AMPK-mediated regulation of angiotensin-like 1 protein. *Cell Rep* 9:495-503

Di Agostino S, Sorrentino G, Ingallina E, Valenti F, Ferraiuolo M, Bicciato S, Piazza S, Strano S, Del Sal G, Blandino G. 2016. YAP enhances the pro-proliferative transcriptional activity of mutant p53 proteins. *EMBO Rep*. 17:188-201.

Di Benedetto A, Mottolese M, Sperati F, Ercolani C, Di Lauro L, Pizzuti L, Vici P, Terrenato I, Sperduti I, Shaaban AM, Sundara-Rajan S, Barba M, Speirs V, De Maria R, Maugeri-Saccà M. 2016a. The Hippo transducers TAZ/YAP and their target CTGF in male breast cancer. *Oncotarget* May 27. doi: 10.18632/oncotarget.9668. [Epub ahead of print]

Di Benedetto A, Mottolese M, Sperati F, Ercolani C, Di Lauro L, Pizzuti L, Vici P, Terrenato I, Shaaban AM, Sundara-Rajan S, Humphries MP, Barba M, Speirs V, De Maria R, Maugeri-Saccà M. 2016b. HMG-CoAR expression in male breast cancer: relationship with hormone receptors, Hippo transducers and survival outcomes. *Sci Rep* 6:35121

Di Lauro L, Vici P, Del Medico P, Laudadio L, Tomao S, Giannarelli D, Pizzuti L, Sergi D, Barba M, Maugeri-Saccà M. 2013. Letrozole combined with gonadotropin-releasing hormone analog for metastatic male breast cancer. *Breast Cancer Res Treat* 141:119-123

Di Lauro L, Vici P, Barba M, Pizzuti L, Sergi D, Rinaldi M, Di Benedetto A, Sperduti I, Shaaban AM, Speirs V, Mottolese M, De Maria R, Maugeri-Saccà M. 2014. Antiandrogen therapy in metastatic male breast cancer: results from an updated analysis in an expanded case series. *Breast Cancer Res Treat* 148:73-80

Di Lauro L, Pizzuti L, Barba M, Sergi D, Sperduti I, Mottolese M, Amoreo CA, Belli F, Vici P, Speirs V, Santini D, De Maria R, Maugeri-Saccà M. 2015. Role of gonadotropin-releasing hormone analogues in metastatic male breast cancer: results from a pooled analysis. *J Hematol Oncol* 8:53

Doyen J, Italiano A, Largillier R, Ferrero JM, Fontana X, Thyss A. 2010. Aromatase inhibition in male breast cancer patients: biological and clinical implications. *Ann Oncol* 21:1243-1245

Eggemann H, Ignatov A, Smith BJ, Altmann U, von Minckwitz G, Röhl FW, Jahn M, Costa SD. 2013. Adjuvant therapy with tamoxifen compared to aromatase inhibitors for 257 male breast cancer patients. *Breast Cancer Res Treat* 137:465-470

Fassan M, Baffa R, Palazzo JP, Lloyd J, Crosariol M, Liu CG, Volinia S, Alder H, Rugge M, Croce CM, Rosenberg A. 2009. MicroRNA expression profiling of male breast cancer. *Breast Cancer Res* 11: R58

Frangou C, Li YW, Shen H, Yang N, Wilson KE, Blijlevens M, Guo J, Nowak NJ, Zhang J. 2014. Molecular profiling and computational network analysis of TAZ-mediated mammary tumorigenesis identifies actionable therapeutic targets. *Oncotarget* 5:12166-12176

Furth N, Bossel Ben-Moshe N, Pozniak Y, Porat Z, Geiger T, Domany E, Aylon Y, Oren M. 2015. Down-regulation of LATS kinases alters p53 to promote cell migration. *Genes Dev.* 29:2325-30.

Ganem NJ, Cornils H, Chiu SY, O'Rourke KP, Arnaud J, Yimlamai D, Théry M, Camargo FD, Pellman D. 2014. Cytokinesis failure triggers hippo tumor suppressor pathway activation. *Cell* 158:833-48.

Graham DK, DeRyckere D, Davies KD et al. 2014. The TAM family: phosphatidylinositol sensing receptor tyrosine kinases gone awry in cancer. *Nat Rev Cancer* 14:769-785

Humphries MP, Jordan VC, Speirs V. 2015. Obesity and male breast cancer: provocative parallels? *BMC Med* 13:134

Johansson I, Nilsson C, Berglund P, Strand C, Jönsson G, Staaf J, Ringnér M, Nevanlinna H, Barkardottir RB, Borg A, Olsson H, Luts L, Fjällskog ML, Hedenfalk I. 2011. High-resolution genomic profiling of male breast cancer reveals differences hidden behind the similarities with female breast cancer. *Breast Cancer Res Treat* 129:747-760

Johansson I, Nilsson C, Berglund P, Lauss M, Ringnér M, Olsson H, Luts L, Sim E, Thorstenson S, Fjällskog ML, Hedenfalk I. 2012. Gene expression profiling of primary male breast cancers reveals two unique subgroups and identifies N-acetyltransferase-1 (NAT1) as a novel prognostic biomarker. *Breast Cancer Res* 14:R31

Johansson I, Ringnér M, Hedenfalk I. 2013. The landscape of candidate driver genes differs between male and female breast cancer. *PLoS One* 8: e78299

Johnson R, Halder G. 2014. The two faces of Hippo: targeting the Hippo pathway for regenerative medicine and cancer treatment. *Nat Rev Drug Discov* 13:63-79

Kornegoor R, Moelans CB, Verschuur-Maes AH, Hogenes MC, de Bruin PC, Oudejans JJ, Marchionni L, van Diest PJ. 2012. Oncogene amplification in male breast cancer: analysis by multiplex ligation-dependent probe amplification. *Breast Cancer Res Treat* 135:49-58

Liu-Chittenden Y, Huang B, Shim JS, Chen Q, Lee SJ, Anders RA, Liu JO, Pan D. 2012. Genetic and pharmacological disruption of the TEAD-YAP complex suppresses the oncogenic activity of YAP. *Genes Dev* 26:1300-1305

Maugeri-Saccà M, Barba M, Pizzuti L, Vici P, Di Lauro L, Dattilo R, Vitale I, Bartucci M, Mottolese M, De Maria R. 2015. The Hippo transducers TAZ and YAP in breast cancer: oncogenic activities and clinical implications. *Expert Rev Mol Med* 17:e14

McShane LM, Altman DG, Sauerbrei W, Taube SE, Gion M, Clark GM. 2006. REporting recommendations for tumor MARKer prognostic studies (REMARK). *Breast Cancer Res Treat* 100:229-235

Nandy SB, Arumugam A, Subramani R, Pedroza D, Hernandez K, Saltzstein E, Lakshmanaswamy R. 2015. MicroRNA-125a influences breast cancer stem cells by targeting leukemia inhibitory factor receptor which regulates the Hippo signaling pathway. *Oncotarget* 6:17366-17378

Pefani DE, O'Neill E. 2016. Hippo pathway and protection of genome stability in response to DNA damage. *FEBS J* 283:1392-1403

Piccolo S, Dupont S, Cordenonsi M. 2014. The biology of YAP/TAZ: hippo signaling and beyond. *Physiol Rev* 94:1287-1312

Piscuoglio S, Ng CK, Murray MP, Guerini-Rocco E, Martelotto LG, Geyer FC, Bidard FC, Berman S, Fusco N, Sakr RA, Eberle CA, De Mattos-Arruda L, Macedo GS, Akram M, Baslan T, Hicks JB, King TA, Brogi E, Norton L, Weigelt B, Hudis CA, Reis-Filho JS. 2016. The Genomic Landscape of Male Breast Cancers. *Clin Cancer Res* 22:4045-4056

Rosenbluh J, Nijhawan D, Cox AG, Li X, Neal JT, Schafer EJ, Zack TI, Wang X, Tsherniak A, Schinzel AC, Shao DD, Schumacher SE, Weir BA, Vazquez F, Cowley GS, Root DE,

Mesirov JP, Beroukhir R, Kuo CJ, Goessling W, Hahn WC. 2012. β -Catenin-driven cancers require a YAP1 transcriptional complex for survival and tumorigenesis. *Cell* 151:1457-1473

Ruddy KJ, Winer EP. 2013. Male breast cancer: risk factors, biology, diagnosis, treatment, and survivorship. *Ann Oncol* 24:1434-1443

Shaaban AM, Ball GR, Brannan RA, Cserni G, Di Benedetto A, Dent J, Fulford L, Honarpisheh H, Jordan L, Jones JL, Kanthan R, Maraqa L, Litwiniuk M, Mottolese M, Pollock S, Provenzano E, Quinlan PR, Reall G, Shousha S, Stephens M, Verghese ET, Walker RA, Hanby AM, Speirs V. 2012. A comparative biomarker study of 514 matched cases of male and female breast cancer reveals gender-specific biological differences. *Breast Cancer Res Treat* 133:949-958

Sorrentino G, Ruggeri N, Specchia V, Cordenonsi M, Mano M, Dupont S, Manfrin A, Ingallina E, Sommaggio R, Piazza S, Rosato A, Piccolo S, Del Sal G. 2014. Metabolic control of YAP and TAZ by the mevalonate pathway. *Nat Cell Biol* 16:357-366

Vici P, Mottolese M, Pizzuti L, Barba M, Sperati F, Terrenato I, Di Benedetto A, Natoli C, Gamucci T, Angelucci D, Ramieri MT, Di Lauro L, Sergi D, Bartucci M, Dattilo R, Pagliuca A, De Maria R, Maugeri-Saccà M. 2014. The Hippo transducer TAZ as a biomarker of pathological complete response in HER2-positive breast cancer patients treated with trastuzumab-based neoadjuvant therapy. *Oncotarget* 5: 9619-9625

Vici P, Ercolani C, Di Benedetto A, Pizzuti L, Di Lauro L, Sperati F, Terrenato I, Gamucci T, Natoli C, Di Filippo F, Botti C, Barba M, Mottolese M, De Maria R, Maugeri-Saccà M. 2016. Topographic expression of the Hippo transducers TAZ and YAP in triple-negative breast cancer treated with neoadjuvant chemotherapy. *J Exp Clin Cancer Res* 35:62

- Wang Z, Wu Y, Wang H, Zhang Y, Mei L, Fang X, Zhang X, Zhang F, Chen H, Liu Y, Jiang Y, Sun S, Zheng Y, Li N, Huang L. 2014. Interplay of mevalonate and Hippo pathways regulates RHAMM transcription via YAP to modulate breast cancer cell motility. *Proc Natl Acad Sci USA* 111: E89-98
- Wu X, Liu X, Koul S, Lee CY, Zhang Z, Halmos B. 2014. AXL kinase as a novel target for cancer therapy. *Oncotarget* 5:9546-9563
- Xiang L, Gilkes DM, Hu H, Takano N, Luo W, Lu H, Bullen JW, Samanta D, Liang H, Semenza GL. 2014. Hypoxia-inducible factor 1 mediates TAZ expression and nuclear localization to induce the breast cancer stem cell phenotype. *Oncotarget* 5:12509-12527
- Yu FX, Luo J, Mo JS, Liu G, Kim YC, Meng Z, Zhao L, Peyman G, Ouyang H, Jiang W, Zhao J, Chen X, Zhang L, Wang CY, Bastian BC, Zhang K, Guan KL. 2014. Mutant Gq/11 promote uveal melanoma tumorigenesis by activating YAP. *Cancer Cell* 25: 822-830
- Zagouri F, Sergentanis TN, Koutoulidis V, Sparber C, Steger GG, Dubsky P, Zografos GC, Psaltopoulou T, Gnant M, Dimopoulos MA, Bartsch R. 2013a. Aromatase inhibitors with or without gonadotropin-releasing hormone analogue in metastatic male breast cancer: a case series. *Br J Cancer* 108:2259-2263
- Zagouri F, Sergentanis TN, Chrysikos D, Zografos E, Rudas M, Steger G, Zografos G, Bartsch R. 2013b. Fulvestrant and male breast cancer: a case series. *Ann Oncol* 24:265-266
- Zanconato F, Forcato M, Battilana G, Azzolin L, Quaranta E, Bodega B, Rosato A, Bicciato S, Cordenonsi M, Piccolo S. 2015. Genome-wide association between YAP/TAZ/TEAD and AP-1 at enhancers drives oncogenic growth. *Nat Cell Biol* 17:1218-1227

Figure Legends

Figure 1: Representative examples of immunohistochemical expression of AXL, TAZ, YAP and CTGF in MBC samples. Panels a-b-c show a sample with cytoplasmic AXL expression (a) cytoplasmic/nuclear TAZ (b) and cytoplasmic CTGF (c). In panels d-f, a tumor case with cytoplasmic AXL expression (d), cytoplasmic/nuclear YAP (e) and cytoplasmic CTGF (f). Scale bar 30 μ m. Magnification 40x. Inset magnification 10x.

Figure 2: Kaplan-Meier survival curves of overall survival regarding: (A) TAZ/CTGF/AXL and (B) YAP/CTGF/AXL (N=116).

Table 1: Baseline characteristics of MBC patients characterized for the expression of AXL (N=116)

Characteristics		N (%)
Age at diagnosis	Median (min-max) [IQ range]	67 (34-88) [59-75]
Histology	IDC/ILC	101 (87.1)
	Other	15 (12.9)
Grade	G1-2	57 (49.1)
	G3	59 (50.9)
Nodal status	Positive	50 (43.1)
	Negative	37 (31.9)
	Unknown	29 (25.0)
Hormone Receptors	Positive (ER+/PgR+)	99 (85.3)
	Other*	17 (14.7)
Ki-67	High ($\geq 14\%$)	51 (44.0)
	Low ($< 14\%$)	65 (56.0)
TAZ/CTGF	Positive	43 (37.1)
	Negative	73 (62.9)
YAP/CTGF	Positive	48 (41.4)
	Negative	68 (58.6)
AXL	Positive	37 (31.9)
	Negative	79 (68.1)

Abbreviations: ER: estrogen receptor, IDC: invasive ductal carcinoma, ILC: invasive lobular carcinoma, PgR: progesterone receptor.

* ER+/PgR- (N=15), ER-/PgR- (N=2)

Table 2: Association between AXL, clinical-pathological factors and expression of the TAZ/CTGF and YAP/CTGF phenotypes.

		AXL		Chi2 Test p-value
		Positive N(%)	Negative N(%)	
Histology	IDC/ILC	32 (31.7)	69 (68.3)	0.999*
	Other	5 (33.3)	10 (66.7)	
Grade	G1-2	14 (24.6)	43 (75.4)	0.096
	G3	23 (39.0)	36 (61.0)	
Nodal status	Positive	18 (36.0)	32 (64.0)	0.666
	Negative	15 (40.5)	22 (59.5)	
Hormone Receptors	ER+/PgR+	31 (31.3)	68 (68.7)	0.745
	Other	6 (35.3)	11 (64.7)	
Ki-67	High	17 (33.3)	34 (66.7)	0.769
	Low	20 (30.8)	45 (69.2)	
TAZ/CTGF	Positive	22 (51.2)	21 (48.8)	0.001
	Negative	15 (20.5)	58 (79.5)	
YAP/CTGF	Positive	23 (47.9)	25 (52.1)	0.002
	Negative	14 (20.6)	54 (79.4)	

Abbreviations: ER: estrogen receptor, IDC: invasive ductal carcinoma, ILC: invasive lobular carcinoma, PgR: progesterone receptor.

* Fisher's Exact Test

Table 3: Uni- and multivariate Cox regression models for 10-year survival (N=116)

		Univariate Cox regression model		Multivariate Cox regression model		Multivariate Cox regression model	
		HR (95%CI)	p-value	HR (95%CI)	p-value	HR (95%CI)	p-value
Histology	IDC/ILC vs other	1.20 (0.36-3.96)	0.762	1.00 (0.30-3.34)	0.994	1.02 (0.30-3.42)	0.977
Grade	G3 vs G1-2	1.84 (0.89-3.80)	0.099	1.69 (0.79-3.60)	0.175	1.70 (0.80-3.63)	0.169
Hormone Receptors	ER⁺/PgR⁺ vs other	0.66 (0.28-1.54)	0.335	0.71 (0.29-1.71)	0.447	0.71 (0.29-1.71)	0.448
Ki-67	High vs Low	1.18 (0.58-2.39)	0.655	1.13 (0.54-2.38)	0.737	1.13 (0.54-2.36)	0.753
TAZ/CTGF/AXL	TAZ/CTGF/AXL vs other	2.37 (1.05-5.34)	0.038	2.31 (1.02-5.22)	0.045		
YAP/CTGF/AXL	YAP/CTGF/AXL vs other	2.31 (1.02-5.22)	0.044			2.27 (1.00-5.13)	0.050

Table 4: Univariate and multivariate Cox regression models for overall survival (N=116).

		Univariate Cox regression model		Multivariate Cox regression model		Multivariate Cox regression model	
		HR (95%CI)	p-value	HR (95%CI)	p-value	HR (95%CI)	p-value
Histology	IDC/ILC vs other	1.30 (0.40-4.29)	0.662	1.11 (0.33-3.70)	0.871	1.13 (0.34-3.77)	0.848
Grade	G3 vs G1-2	1.94 (0.94-3.97)	0.071	1.85 (0.88-3.91)	0.106	1.86 (0.88-3.93)	0.103
Hormone Receptors	ER⁺/PgR⁺ vs other	0.76 (0.32-1.76)	0.519	0.85 (0.35-2.04)	0.718	0.85 (0.35-2.04)	0.719
Ki-67	High vs Low	1.09 (0.54-2.18)	0.805	1.01 (0.49-2.08)	0.982	1.00 (0.48-2.07)	0.995
TAZ/CTGF/AXL	TAZ/CTGF/AXL vs other	2.26 (1.01-5.08)	0.048	2.19 (0.97-4.94)	0.058		
YAP/CTGF/AXL	YAP/CTGF/AXL vs other	2.21 (0.98-4.97)	0.054			2.16 (0.96-4.87)	0.063




Figure 1




Figure 2