



Biology of Blood and Marrow Transplantation

journal homepage: www.bbmt.org



Reviews

The Microbiome and Hematopoietic Cell Transplantation: Past, Present, and Future



Tessa M. Andermann¹, Jonathan U. Peled^{2,3}, Christine Ho⁴, Pavan Reddy⁵, Marcie Riches⁶, Rainer Storb^{7,8}, Takanori Teshima⁹, Marcel R.M. van den Brink^{10,11}, Amin Alousi¹², Sophia Balderman¹³, Patrizia Chiusolo¹⁴, William B. Clark¹⁵, Ernst Holler¹⁶, Alan Howard¹⁷, Leslie S. Kean^{7,18,19}, Andrew Y. Koh²⁰, Philip L. McCarthy⁴, John M. McCarty²¹, Mohamad Mohty^{22,23,24}, Ryotaro Nakamura²⁵, Katy Rezvani^{26,27}, Brahm H. Segal^{28,29,30}, Bronwen E. Shaw³¹, Elizabeth J. Shpall³², Anthony D. Sung³³, Daniela Weber¹⁶, Jennifer Whangbo³⁴, John R. Wingard^{35,36}, William A. Wood³⁷, Miguel-Angel Perales^{2,3}, Robert R. Jenq^{38,*}, Ami S. Bhatt^{39,**} on behalf of the Blood and Marrow Transplant Clinical Trials Network

¹ Division of Infectious Diseases, Department of Medicine, Stanford University, Stanford, California

² Adult Bone Marrow Transplantation Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, New York

³ Department of Medicine, Weill Cornell Medical College, New York, New York

⁴ Blood and Marrow Transplantation, Department of Medicine, Roswell Park Comprehensive Cancer Center, Buffalo, New York

⁵ Department of Medicine, University of Michigan Cancer Center, Ann Arbor, Michigan

⁶ Division of Hematology/Oncology, Department of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina

⁷ Fred Hutchinson Cancer Research Center, Seattle, Washington

⁸ Department of Medicine, University of Washington School of Medicine, Seattle, Washington

⁹ Department of Hematology, Hokkaido University Faculty of Medicine, Sapporo, Japan

¹⁰ Immunology Program, Sloan Kettering Institute, New York, New York

¹¹ Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, New York

¹² Multidiscipline GVHD Clinic and Research Program, Department of Stem Cell Transplant and Cellular Therapies, University of Texas, MD Anderson Cancer Center, Houston, Texas

¹³ Department of Medicine, Roswell Park Comprehensive Cancer Center, Buffalo, New York

¹⁴ Hematology Department, Fondazione Policlinico Universitario A. Gemelli, Università Cattolica Sacro Cuore, Rome, Italy

¹⁵ Bone Marrow Transplant Program, Division of Hematology/Oncology and Palliative Care, Department of Internal Medicine, Virginia Commonwealth University, Richmond, Virginia

¹⁶ Department of Internal Medicine 3, University Medical Center, Regensburg, Germany

¹⁷ Center for International Blood and Marrow Transplant Research, Minneapolis, Minnesota

¹⁸ Department of Pediatrics, University of Washington School of Medicine, Seattle, Washington

¹⁹ Ben Towne Center for Childhood Cancer Research, Seattle Children's Research Institute, Seattle, Washington

²⁰ Divisions of Hematology/Oncology and Infectious Diseases, Departments of Pediatrics and Microbiology, University of Texas Southwestern Medical Center, Dallas, Texas

²¹ Bone Marrow Transplantation Program, Virginia Commonwealth University Massey Cancer, Richmond, Virginia

²² Clinical Hematology and Cellular Therapy Department, Hôpital Saint-Antoine, AP-HP, Paris, France

²³ Sorbonne Université, Paris, France

²⁴ INSERM UMRs U938, Paris, France

²⁵ Department of Hematology and Hematopoietic Cell Transplantation, City of Hope, Duarte, California

²⁶ Section of Cellular Therapy, Good Manufacturing Practices Facility, Department of Stem Cell Transplant and Cellular Therapy, University of Texas MD Anderson Cancer Center, Houston, Texas

Financial disclosure: See Acknowledgments on page 1337.

* Correspondence and reprint requests: Robert R. Jenq, MD, Departments of Genomic Medicine and Stem Cell Transplantation Cellular Therapy, Division of Cancer Medicine, University of Texas MD Anderson Cancer Center, 1515 Holcombe Boulevard, Houston, TX 77030-4000.

E-mail address: rjenq@mdanderson.org (R.R. Jenq).

** Correspondence and reprint requests: Ami S. Bhatt, MD, PhD, Division of Hematology, Department of Medicine, Department of Genetics, Stanford University, 269 Campus Drive, CCSR1 155b, Stanford, CA 94305.

E-mail address: asbhatt@stanford.edu (A.S. Bhatt).

²⁷ Department of Medicine, University of Texas MD Anderson Cancer Center, Houston, Texas

²⁸ Department of Medicine, University at Buffalo Jacobs School of Medicine and Biomedical Sciences, Buffalo, New York

²⁹ Division of Infectious Diseases, Roswell Park Comprehensive Cancer Center, Buffalo, New York

³⁰ Department of Immunology, Roswell Park Comprehensive Cancer Center, Buffalo, New York

³¹ Center for International Blood and Bone Marrow Transplant Research, Medical College of Wisconsin, Milwaukee, Wisconsin

³² Cell Therapy Laboratory and Cord Blood Bank, Department of Stem Cell Transplantation and Cellular Therapy, University of Texas, MD Anderson Cancer Center, Houston, Texas

³³ Division of Hematologic Malignancies and Cellular Therapy, Duke University School of Medicine, Duke Cancer Institute, Durham, North Carolina

³⁴ Dana-Farber Cancer Institute, Boston Children's Hospital, Boston, Massachusetts

³⁵ Department of Medicine, University of Florida Health Cancer Center, Gainesville, Florida

³⁶ Bone Marrow Transplant Program, Division of Hematology/Oncology, University of Florida College of Medicine, Florida

³⁷ Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina

³⁸ Departments of Genomic Medicine and Stem Cell Transplantation Cellular Therapy, Division of Cancer Medicine, University of Texas MD Anderson Cancer Center, Houston, Texas

³⁹ Department of Genetics and Division of Hematology, Department of Medicine, Stanford University, Stanford, California

Article history:

Received 9 January 2018

Accepted 8 February 2018

Key Words:

Microbiome

Microbiota

Hematopoietic stem cell

transplantation

Metagenomics

Prebiotic

Fecal microbiota transplant

© 2018 American Society for Blood and Marrow Transplantation.

INTRODUCTION

The human microbiome is comprised of the bacteria, archaea, viruses, fungi, and other microeukaryotes that live on and within the human host. Alterations in the microbiome are associated with adverse transplant outcomes including the expected infectious complications following allogeneic hematopoietic cell transplantation (allo-HCT) in addition to diseases that are not classically microbe associated. For example, recent data suggest an association between certain microbial community structures and mortality, disease relapse, risk of infection, and graft-versus-host disease (GVHD) [1–3]. Most studies inspecting the role of the microbiome in HCT patient outcomes, though compelling, are limited in scope: in general, data have been generated in single-center studies

or preclinical models. Here, we summarize many of the main findings of the past several decades of research on this topic, and propose areas of focus for future research that will facilitate investigating the microbiome and its role in disease (Table 1).

The Earliest Days of Microbiota Research: Germ-Free Mice and Patients

Pioneering studies conducted in the early 1970s demonstrated that mice undergoing allo-HCT in germ-free conditions experienced less GVHD and had improved survival (Table 2) [4–16]. Soon thereafter, this observation led to attempts to reproduce these conditions in patients undergoing HCT through the use of laminar-airflow isolation rooms, “sterile”

Table 1

Proposed Next Steps for Future Research Investigating the Microbiome-Host Relationship in HCT Patients

Primary Research Gaps and Strategies
1. Multi-institutional prospective studies to improve generalizability of microbiota research findings
2. Multicenter biospecimen and data collection to support collaborative and integrative “multi-omic” approaches to microbiota research
3. Expansion of microbial metagenomic studies in HCT patients beyond bacterial taxonomy
a. Investigation of the role of the virome and mycobiome in HCT outcomes
b. Elucidation of microbial genes and metabolic pathways that impact outcomes, in particular the antibiotic resistome
4. Targeting of specific microbes and/or microbial pathways to determine whether modification of the microbiota can impact microbe-disease associations
5. Investigation of the role of extraintestinal microbial populations (ie, oral, skin, vaginal, and eye) in the development of acute and chronic GVHD
6. Investigation of microbial-host dynamics associated with the development of chronic GVHD and other non-GVHD outcomes
7. Modification of antibiotic strategies that balance preservation and restoration of the microbiome with appropriate infection prevention and treatment
8. Antibiotic-sparing approaches to infection prevention and treatment; steroid-sparing approaches to GVHD prevention and treatment
9. Incorporation of standardized antibiotic and infection-related data into national transplant databases
10. Understanding the potential for using the microbiome as a biomarker for transplant outcomes and as a guide to interventions

Table 2
Early Years of Microbiota Research in Germ-Free Mice

Model system and intervention	Findings	References
Germ-free mice versus conventional mice	Significantly milder GVHD symptoms and longer survival after MHC-disparate allo-HCT in germ-free mice	Connell, 1965 [4] Jones, 1971 [5] Van Bekkum, 1961 [6] Van Bekkum, 1977 [7]
	Intact GVT and reduced GVHD after MHC-disparate allo-HCT in germ-free mice	Pollard, 1973 [8] Pollard, 1974 [9] Truitt, 1974 [10] Truitt, 1976 [11] Van Bekkum, 1974 [12]
Germ-free mice versus conventional mice versus mice with consortium of colonization resistant intestinal microflora (anaerobes)	Significantly milder GVHD symptoms and longer survival in both germ-free and colonization resistant mice whose microflora was predominantly anaerobic Conventionalization of all mice after 40 days did not induce GVHD in gnotobiotic mice that received colonization resistant microflora but did in conventional mice treated with antibiotics	
Antibiotic-treated mice versus untreated mice	Significantly milder GVHD in xenogeneic rat-to-mouse HCT; GVHD histology present but less inflamed in antibiotic-treated mice compared with conventional mice Significantly milder GVHD symptoms and longer survival after MHC-disparate allo-HCT in antibiotic-treated mice	Heit, 1973 [13] Van Bekkum, 1967 [14] Heit, 1977 [15]
Selective antibiotic decontamination of Enterobacteriaceae versus conventional mice	Mitigation of delayed-type GVHD by selective decontamination of Enterobacteriaceae was minor and dependent on mouse model	Veenendaal, 1988 [16]

MHC indicates major histocompatibility.

diets, gut decontamination with oral nonabsorbable antibiotics, and skin cleansing (Table 3) [17–29]. Between-study heterogeneity and the lack of reproducible data supporting efficacy for GVHD prevention have limited evidence-based guidance for clinical practice and prophylactic strategies. However, while many of these previous approaches have largely been abandoned, broadly-adopted modern-day recommendations to prevent infectious complications include antimicrobial prophylaxis, sterilized positive air pressure rooms, low-microbial diet, and use of barrier precautions (eg, gloves, face masks, gowns) [30]. Recently, there has been a resurgence of microbiome research across many disciplines of medicine spurred by advances in high-throughput methodologies for characterizing the microbiome, which extend beyond bacterial culture techniques and virus-specific molecular approaches for detection. Similarly, there has been rapid growth in the microbiome research as it relates to HCT.

METHODS TO INVESTIGATE THE MICROBIOME

With the advent of high-throughput molecular methods to study the microbiome, the field has grown significantly in the past decade. Commonly used methods include 16S ribosomal RNA gene sequencing for bacterial taxonomic classification, metabolomics, as well as shotgun metagenomic sequencing and subsequent taxonomic and functional classification of microbial genes; these methods have been reviewed in detail elsewhere (Table 4) [31–33]. Each of these methods provides an orthogonal approach to study the microbiome from the perspective of answering important microbiota taxonomic and functional questions, including “which microbes are there?”; “what do they make?”; “what genes do they contain?”; and “what is their relative and absolute abundance?” [34]. With the explosion of new molecular and bioinformatic approaches to study the microbiome, we anticipate an ever-growing toolkit to characterize potentially clinically relevant features of the microbiome such as antibiotic resistance, microbial virulence factors, and strain dynamics. Terms commonly used in microbiome studies and their definitions are listed in Table 5 [31,35–38].

A precedent has recently been set for the generation of multifaceted data types (ranging from shotgun metagenomic sequencing to transcriptomes and epigenomes to metabolite profiling) that facilitate multidimensional and longitudinal characterization of both the host and the microbiome. Specifically, projects of the integrative Human Microbiome Project, the U.S. National Institutes of Health Common Fund’s second phase Human Microbiome Project, have collected longitudinal samples from 3 cohorts of individuals (comprising individuals with pregnancy and preterm birth, type 2 diabetes, and inflammatory bowel disease) [39,40]. Given these advances in “multi-omic” data collection, we anticipate that the next decade of translational research in the microbiome field as it relates to HCT will extend far beyond simple characterization of community taxonomic structures within the microbiome. For example, advances in immunophenotyping and short-term, in vitro propagation of microbial mixtures will identify potential mechanistic relationships among microbes, microbial antigens, and host responses [41,42].

While the advances in phenotyping and genotyping experiments may pave the way for the identification of biomarkers that may be clinically actionable, there are challenges and limitations to their effective, wide-scale application. For example, a specific challenge is the need for rapid turnaround of next-generation sequencing results to be clinically actionable; at present, this is not routinely available due to the need to batch samples to reduce the costs of sequencing. Thus, while next-generation sequencing and metabolomic approaches are the predominant technologies used in the research setting, some intriguing and potentially more easily deployed alternatives for microbiome measurement may be more translatable. Indeed, these technologies do exist and include approaches such as species- or bacterial group-specific quantitative PCR [32] and microarray approaches [33,43]. Despite the challenges to their clinical use, it is feasible for advancements in microbiota science currently based on next-generation sequencing to be optimized for real-time use both diagnostically and prognostically in the HCT setting.

Table 3
Clinical Studies of Protective Isolation with Gut Decontamination in HCT Patients

Intervention	Control	Outcomes (intervention versus control)	References
Randomized trials			
Protective isolation (LAF isolation, skin cleansing, sterile diets) + oral antibiotic decontamination (n = 45)	Oral antibiotic decontamination (n = 44)	<ul style="list-style-type: none"> • Decreased risk of infection • Fewer patients with septicemia • Fewer days with septicemia • Longer time to first major infection • Later onset GVHD (only in aplastic anemia patients) • No difference in survival • Of note, adherence to oral antibiotic decontamination was poor 	Buckner, 1978 [17]
LAF isolation + oral antibiotic decontamination (n = 36)	Conventional rooms + hand washing and mask precautions (n = 31)	<ul style="list-style-type: none"> • Decreased incidence of infections after engraftment • Later onset acute GVHD • Trend toward decreased incidence of grade II to IV acute GVHD; not statistically significant • Increased survival 	Navari, 1984 [18]
LAF isolation + oral nonabsorbable antibiotics + prophylactic systemic antibiotics (n = 54)	LAF + oral nonabsorbable antibiotics (n = 68)	<ul style="list-style-type: none"> • Fewer episodes of septicemia • No difference in incidence or severity of GVHD • No difference in mortality 	Petersen, 1986 [19]
Conventional rooms + prophylactic systemic antibiotics (n = 45)	Conventional rooms + prophylactic granulocyte infusions (n = 67)	<ul style="list-style-type: none"> • No difference in incidence of septicemia or local major infections • No difference in GVHD incidence • No difference in mortality 	Petersen, 1986 [20]
LAF + prophylactic systemic antibiotics (n = 49)	Conventional rooms + prophylactic systemic antibiotics (n = 50)	<ul style="list-style-type: none"> • Decreased septicemia • Decreased major local infections (borderline significance) • No difference in GVHD incidence • No difference in mortality 	Petersen, 1987 [21]
Conventional rooms + ciprofloxacin + metronidazole gut decontamination (n = 68)	Conventional rooms + ciprofloxacin gut decontamination (n = 66)	<ul style="list-style-type: none"> • Decreased incidence of acute GVHD • No difference in chronic GVHD • No difference in OS 	Beelen, 1999 [22]
Observational trials			
LAF isolation + oral antibiotic decontamination (n = 39)	Conventional rooms + oral antibiotic decontamination (n = 91)	<ul style="list-style-type: none"> • Decreased incidence of acute GVHD • Increased long-term survival 	Storb, 1983 [23]
LAF isolation + oral and topical antibiotic decontamination (n = 26)	Conventional rooms + barrier nursing + oral and topical antibiotic decontamination (n = 22)	<ul style="list-style-type: none"> • No difference in GVHD incidence • No difference in mortality 	Mahmoud, 1984 [24]
Strict reverse isolation* + oral nonabsorbable antibiotics, complete gut decontamination (n = 26)	Barrier nursing + oral nonabsorbable antibiotics, selective gut decontamination (n = 15)	<ul style="list-style-type: none"> • Fewer days of fever and fewer infections • Trend toward decreased GVHD although not statistically significant 	Schmeiser, 1988 [25]
LAF + complete antibiotic oral decontamination (n = 44)	LAF + selective decontamination (n = 21)	<ul style="list-style-type: none"> • Fewer infections • Decreased acute and chronic GVHD incidence • Lower combined TRM or chronic GVHD incidence 	Vossen, 1990 [26]
Sustained growth suppression of anaerobic bacteria with oral nonabsorbable and systemic antibiotics (n = 41)	Incomplete growth suppression of anaerobic bacteria despite oral nonabsorbable and systemic antibiotics (n = 153)	<ul style="list-style-type: none"> • Decreased incidence of acute GVHD 	Beelen, 1992 [27]
Protective isolation with either LAF or HEPA filters (n = 4238)	Conventional isolation (n = 827)	<ul style="list-style-type: none"> • No difference in acute or chronic GVHD incidence • Decreased TRM and overall mortality in the first 100 days post-transplant 	Passweg, 1998 [28]
Protective isolation + successful gut decontamination with oral nonabsorbable and systemic antibiotics (n = 57)	Protective isolation -successful gut decontamination despite oral nonabsorbable and systemic antibiotic (n = 55)	<ul style="list-style-type: none"> • Decreased infectious risk • Decreased incidence of acute GVHD 	Vossen, 2014 [29]

LAF indicates laminar air flow; OS, overall survival; TRM, transplant related mortality.

* Reverse isolation was achieved in sterile plastic isolators.

Open questions

- How do we design and execute microbiome studies that permit simultaneous characterization of (1) microbes beyond taxonomy and (2) both host impact on and host response to the microbiome, thus informing a deeper understanding of microbiota function?
- What new technologies (eg, single cell sequencing, T cell receptor sequencing, long-read sequencing, and advanced imaging) will emerge as new ways to measure the

Table 4
Sequencing Technologies Used in Microbiome Research

Method	Definition	References
Metagenomics	The study of genes and noncoding genetic information in a mixed population of organisms to infer functional potential and taxonomic structure of the population and its individual organisms	Marchesi, 2015 [31]
16S ribosomal RNA/DNA sequencing	PCR amplification of bacterial RNA/DNA from the variable regions of the 16S ribosomal RNA gene for taxonomic profiling (the region selected is usually determined by the niche that is being investigated; eg, V4 for stool, V1-3 for skin)	Weisburg, 1991 [32]
Shotgun next-generation metagenomic sequencing	High-throughput DNA sequence generation and analysis from any organism or group of organisms via fragmentation, tagging, amplification, and massively parallel or deep sequencing. Allows for taxonomic identification in addition to generating information about gene presence, genetic bioregulation, and potential metabolic pathways	Loman, 2012 [33]
Metatranscriptomics	High-throughput RNA sequence generation and analysis from any organism or group of organisms via reverse transcription, tagging, amplification, and massively parallel or deep sequencing. Provides a snapshot of which genes are being transcribed	Marchesi, 2015 [31]
Metabolomics	Characterization of the collection of metabolites produced by an organism or a single tissue. The term has been used to describe characterization of the collection of metabolites produced by a collection of organisms (ie, the microbiota), although some prefer the term <i>metabonomics</i> for that definition	Marchesi, 2015 [31]
Metaproteomics	Characterization of all proteins within a clinical or environmental sample	Marchesi, 2015 [31]

microbiome, its structure, function and interactions with the host; how should we apply these approaches in the HCT setting?

- What strategies for multicenter collections of biospecimens and clinical data best support future integrative “multi-omic” approaches to illuminate host-microbe relationships as they pertain to HCT outcomes?

THE MICROBIOME AS BIOMARKER

Identifying biomarkers with high prognostic and predictive value is crucial for communicating risk to patients and selecting appropriate therapeutic strategies. Thus, it is no surprise that composition of the intestinal microbiota, which is affected by host genetic factors, immunological factors, diet, medications, lifestyle, and environmental exposures, has been analyzed as a biomarker for important clinical outcomes after HCT. During transplantation, dramatic shifts in the composition of the intestinal flora are observed [1]. These shifts in species abundance and measures of diversity have been proposed as potential biological markers associated with patient outcomes after transplantation (Table 6) [1-3,22,27,44-49]. Biomarkers such as these may prove useful in the design of clinical trials to identify patients at risk of certain outcomes, as surrogate markers of clinical outcomes, or as early predictive markers of treatment response. If a causal relationship between the microbiota and transplant outcomes can

be established, these relationships may inform the development of microbiota-based therapeutic interventions to improve transplant outcomes.

3-Indoxyl Sulfate as Biomarker of Intestinal Microbiota Health

An example of a recently proposed microbiome-derived biomarker in HCT is the small, aromatic tryptophan metabolite 3-indoxyl sulfate [46,50]. Indoxyl sulfate originates from the degradation of dietary protein-derived tryptophan to indole by the tryptophanase of commensal intestinal bacteria. After resorption of indole from the intestine, it is metabolized to indoxyl sulfate in the liver and finally excreted in the urine. Microbiota-derived indole and its derivatives are integral to the maintenance of human microbial communities through bacteriostatic effects on Gram-negative enteric bacteria, antifungal activities that provide colonization resistance to *Candida albicans*, as well as regulation of epithelial function and control of local inflammation through induction of anti-inflammatory cytokines [51,52]. Recent studies have demonstrated that indoxyl sulfate can serve as an important biomarker with lower urine levels being associated with significant and clinically relevant intestinal microbiota disruption in patients undergoing allo-HCT [3,46,53]. In the future, metabolites such as indoxyl sulfate may serve as a urine or serum marker for monitoring

Table 5
Definition of Terms

Term	Definition	References
Diversity	Measurement of the number of different types (taxa) of organisms and their abundance. Alpha-diversity and beta-diversity refer to diversity within and between samples, respectively	Lozupone, 2012 [35]
Dysbiosis	Perturbation of the taxonomic structure and function of the microbiota from the healthy state. This can be associated with the development of disease	Petersen, 2014 [36]
Germ-free mice	Mice raised in sterile conditions and free from colonization by all microorganisms. Also referred to as “axenic” mice	Giraud, 2008 [37]
Fecal microbiota transplantation	The transfer of stool from a donor to a recipient via either endoscopy, nasogastric/duodenal tube, capsules, or enema for the purpose of altering the intestinal microbiota of the recipient and restoring health. Stool is obtained from healthy related or unrelated donors, and less commonly from the intended recipient	Borody, 2013 [38]
Microbiota	The entirety of microorganisms (bacteria, archaea, viruses, fungi, and other eukaryotes) within a specific habitat	Marchesi, 2015 [31]
Microbiome	Includes the biotic (microorganisms and their genomes) and abiotic (environmental) factors present within a particular habitat. This definition is modeled after the meaning of the term “biome”. Many in the field use the term microbiome to refer to the collection of genes and genomes within a particular habitat, although this definition is redundant with “metagenome”	Marchesi, 2015 [31]

Table 6
Clinical Studies Examining the Microbiota as a Biomarker for HCT Outcomes

Microbiota Feature	Association	Sample size	References
Sustained decontamination of gut anaerobes	Lower risk of GVHD	194	Beelen, 1992 [27]
Decontamination of gut anaerobes	Lower risk of GVHD	134	Beelen, 1999 [22]
Intestinal monodomination by <i>Enterococcus</i> and Proteobacteria	Higher risk of bacteremia and intestinal GVHD	94	Taur, 2012 [44]
Intestinal monodomination, especially by <i>Enterococcus</i>	Higher risk of bacteremia and intestinal GVHD	31	Holler, 2014 [3]
Decreased duodenal Paneth cell counts at GVHD	Higher GI GVHD severity, lower GVHD treatment response, and higher NRM	142	Levine, 2013 [45]
Low intestinal microbiota diversity	Lower OS, higher TRM	80	Taur, 2014 [1]
Lower urinary 3-indoxyl sulfate	Higher intestinal microbiota dysbiosis, higher risk of GVHD	31	Holler, 2014 [3]
Lower urinary 3-indoxyl sulfate	Higher intestinal microbiota dysbiosis, higher TRM, lower OS	131	Weber, 2015 [46]
Higher fecal <i>Blautia</i> abundance	Lower GVHD-related mortality, higher OS	115	Jenq, 2015 [47]
Higher abundance or presence of a cluster of bacteria including <i>Eubacterium limosum</i> in fecal microbiota	Lower risk of relapse or progression of disease, higher OS	541	Peled, 2017 [2]
Higher gradient of positively to negatively correlated organisms at neutrophil recovery	Higher risk of severe acute GVHD	66	Golob, 2017 [48]
Picobirnavirus presence	Severe GI GVHD	44	Legoff, 2017 [49]

All studies are observational except Beelen, 1999 [22].

GI indicates gastrointestinal; NRM, nonrelapse mortality; OS, overall survival; TRM, transplant related mortality.

microbiota perturbations in patients being treated with antibiotics or in predicting the development of GVHD [46].

Open questions

- Can the composition of the intestinal microbiota serve as a biomarker for clinical outcomes after HCT and can it be used to guide interventions?
- Are there specific microbes that are causally associated with positive or negative outcomes of HCT, perhaps through their antigenic properties or through the action of generated metabolites? If so, might there be strain-specific differences in their capacity to induce inflammation or cytotoxic damage?
- How can changes in the microbiome be assayed in real time to allow for clinical decision making?

INTERACTIONS BETWEEN THE MICROBIOME AND THE IMMUNE SYSTEM

The Microbiome and Development of the Immune System

Proper immune reconstitution is central to successful HCT. To better understand immune reconstitution in patients following HCT, it is helpful to turn to the well-studied and analogous process of immune development in neonates. The adaptive immune system and microbiota undergo a process of rapid change and development over the first 3 years of human life, and these 2 processes are intimately interconnected [54]. Developmental microbiota perturbations have been associated with short-term immune consequences early in life, and there is a strong suggestion that these early perturbations may have long-term deleterious effects on immune function as well [55–60]. As successful immune reconstitution is central to HCT efficacy in the short and long term, it is critical to understand exactly how the microbiota impacts that process.

Axenic or “germ-free” animal models have been an essential tool in defining the importance of microbes to immune development [61–63]. Studies in these systems have shown

that the immune system of a germ-free neonate is underdeveloped [64]. Most notable are the changes observed in mucosal immunity, particularly in the intestine, with absence of gut-associated lymphoid tissue, including isolated lymphoid follicles, Peyer’s patches, and mesenteric lymph nodes [65]. While many similarities exist between the immunologic development process in infants and HCT recipients, some differences must also be considered. Notably, the adult intestinal microbiome is quite divergent from the infant microbiome, and individuals undergoing HCT have often received antimicrobial and other pharmacological agents that damage the microbiota composition. Thus, while the microbiota likely plays a role in immune reconstitution post-HCT, we must carefully consider both the similarities and differences to the process of infant immune development as we try to understand how the microbiota impacts both immune reconstitution and adverse immunological post-HCT outcomes such as GVHD.

The Microbiome and Its Role in Immune Reconstitution Post-HCT

Impaired immune reconstitution after HCT is a significant cause of morbidity and mortality, and has been implicated in increased risk of infections, malignancy relapse, and development of secondary malignancies [66–74]. Given recent data on the interactions between the gut microbiome and transplant outcomes discussed previously, as well as the immune system [75–78], it is reasonable to hypothesize that the microbiome plays a direct role in post-transplant immune recovery. The study of post-transplant immune reconstitution now benefits from a variety of quantitative and qualitative assays, including clinical parameters such as absolute lymphocyte counts, lymphocyte subsets (CD4⁺ and CD8⁺ T cells, natural killer cells, B cells), and antibody titers, as well as more complex functional assays and evaluations of T cell and B cell repertoire, and next-generation sequencing approaches to provide information on TCR diversity and specific clonotypes over time [79]. Currently, no integrated datasets comprising

simultaneous host and microbiome measurements are publicly available for analysis. Future prospective studies will need to integrate these areas of research to better define potential interactions between the immune system and the microbiome in HCT, as has been done in the integrative Human Microbiome Project for other diseases [39,40].

Potential Mechanisms of Immune Modulation by the Microbiota That Impact GVHD and Graft-Versus-Tumor Effect

The largest proportion of microorganisms in the body exists in the lower intestine, and thus the intestine is believed to be the major interface between the microbiome and adaptive immune system. Intestinal homeostasis is a dynamic process that includes maintenance of bowel mucosa integrity and relies heavily on the interactions between immunologic function and the community of organisms that make up the gut microbiota. HCT leads to dysbiosis and disruption of intestinal homeostasis as a result of the conditioning regimen, use of broad-spectrum antibiotics, alterations in nutrition, and donor cell-derived immune reconstitution. There is clinical evidence for the regulatory effect of gut microbiota in the maintenance of intestinal homeostasis mediated primarily through regulatory T cells [80–82]. For example, emerging data suggest that alterations in the intestinal microbiota and metabolome are associated with the incidence and severity of acute GVHD (Tables 7A and 7B) [1,3,22,26,27,29,48–50,54,76,84–96]. While the majority of the literature has focused on changes in intestinal microbiota diversity, others have focused on the role of particular organisms, such as *Blautia* spp. in protection from GVHD [47]. Reports of GVHD associated with blooms of eukaryotic viruses, such as picobirnaviruses, have also begun to emerge, suggesting a potential role for the human virome as, at the very least, a marker of this transplant complication [49]. In addition to increasing GVHD risk, disruption in intestinal homeostasis and dysbiosis is associated with increased treatment-related mortality (TRM), and decreased overall survival [2,3,48].

Commensal bacteria can also play a role in tumor immunosurveillance. Although the precise mechanisms by which intestinal microbes can promote tumor immunity are unknown, one hypothesis invokes antigen mimicry, as microbial proteins can bear close resemblance to tumor-associated antigens [96]. An alternative pathway might be through nonspecific activation of innate immune cells and pathways. Consistent with the notion of the microbiome influencing antitumor immunity, it was recently shown that specific members of the intestinal microbiota are associated with a decreased risk of relapse after allo-HCT [2]. Achieving a comprehensive understanding of the mechanisms driving both mucosal and systemic immune modulation by the gut microbiota may facilitate the simultaneous mitigation of GVHD while maintaining or improving GVT effects.

Whether particular microbiota signatures correspond to a causal or contributing factor to the development of various disease phenotypes remains to be elucidated, as most clinical studies have established only associations, with rare exception [22]. Specifically, analyses of alterations in intestinal microbiota have focused on time-course compositional descriptions and correlations with clinical and biological outcomes, in particular acute GVHD. Future research will undoubtedly bring greater focus on both intestinal and extraintestinal microbial alterations and their mechanistic impact on the development and severity of acute and chronic GVHD in addition to other transplant outcomes. The exper-

imental data so far indicate that modification of the gut ecosystem to restore intestinal homeostasis may represent a novel approach to modulate complications of HCT. While attempts at altering therapeutically the established intestinal dysbiosis could potentially improve transplant outcomes, we are in the beginning phases of comprehending the full impact and the mechanistic role of microbiota in HCT, not only for GVHD outcomes, but also for tumor relapse, infectious complications, and long-term outcomes after HCT.

Open questions

- Can we identify specific associations among the microbial taxa, antigens, or metabolites and post-HCT immune recovery?
- Is the pretreatment microbiome (before any treatment for the underlying disorder) prognostic of immunologic and other outcomes post-transplant?
- What is the role of oral and skin microbiomes as well as microbes from other organs on the incidence and severity of acute and chronic GVHD? Is there a specific set of intestinal and extraintestinal (eg, ocular, skin, vaginal) microbial taxonomic structures over time that correlate with or are causally related to chronic GVHD and late effects of HCT?
- How can we use interventions that modify the microbiome to improve post-HCT outcomes, specifically mediated by immune effects on GVHD and GVT?
- Do specific T cell responses against bacterial antigens affect donor and recipient T cell repertoires and therefore play a role in HCT outcomes?
- Do donor lymphocyte infusions, checkpoint inhibitors, chimeric antigen receptor T cell and other T cell therapies impact the microbiome? If so, does the microbiome in any way mediate clinical outcomes following these therapies?

THE MICROBIOME AND ITS ROLE IN INFECTION AND IDIOPATHIC POST-HCT DISORDERS

Infection is a major cause of nonrelapse morbidity and mortality after HCT, second only to GVHD. Unfortunately, other than administering prophylactic antibiotics or antiviral agents to susceptible patients, the therapeutic approach against these infections is largely reactive. Thus, identifying modifiable host or microbiome features that can be manipulated to prevent infection is a very attractive and promising proposition. The gut microbiota plays a critical role in maintaining colonization resistance against intestinal pathogens, and the mechanisms that underlie this regulation are becoming increasingly well understood. The composition of intestinal microbiota is actively regulated by a number of internal and external factors, ranging from diet to antibiotics, to the elements of the adaptive and innate immune system. An important element of the innate immune system that shapes the microbiota is host-derived antimicrobial peptides (AMPs). Examples of such AMPs include Paneth cell-derived α -defensins and REG3 α , which selectively eliminate noncommensals while preserving commensals, and thus serve as microbiome modulators [97,98]. Intestinal commensal bacteria can stimulate the gut epithelium to produce AMPs that kill pathogenic bacteria [99] and fungi [51]. In GVHD, for example, Paneth cell loss is associated with both reduced secretion of α -defensins and intestinal dysbiosis [84,100,101]. Additionally, disruption of gut microbial communities by antibiotics can increase susceptibility to intestinal pathogens [102,103]. Microbiota disruption that leads to gut microbial monodominance (eg, a microbiome

Table 7

Studies of Microbiota Changes Associated with GVHD in Murine Models and Microbiota Changes Associated with GVHD in Clinical Studies

A: Studies of microbiota changes associated with GVHD in murine models			
Intervention/comparison	Outcome (Intervention versus control)		References
Reduced-intensity allogeneic versus syngeneic HCT	<ul style="list-style-type: none"> GVHD accompanied by higher Enterobacteriaceae, <i>Bacteroides</i> and <i>Enterococcus spp.</i>, and lower <i>Lactobacilli</i>, <i>Clostridia</i>, <i>Bifidobacterium</i>, and <i>Bacillus spp.</i> Higher <i>Escherichia coli</i> associated with GVHD severity and reduced survival Treatment with ciprofloxacin did not affect severity of GI GVHD 		Heimesaat, 2010 [84]
MHC-disparate and MHC-matched/minor antigen-mismatched allo-HCT ± donor T cells	<ul style="list-style-type: none"> GVHD associated with: <ul style="list-style-type: none"> Loss of intestinal Paneth cells Reduction in α-defensin expression Lower levels of Firmicutes and Bacteroidetes Intestinal <i>Escherichia coli</i> expansion and dominance Lower microbiota diversity Oral administration of the antibiotic Polymyxin B, active against <i>E. coli</i>: <ul style="list-style-type: none"> Decreased GVHD severity Decreased GVHD-related mortality 		Eriguchi, 2012 [85]
MHC-disparate allo-HCT ± donor T cells	<ul style="list-style-type: none"> GHVD associated with: <ul style="list-style-type: none"> Loss of intestinal microbiota diversity Higher <i>Lactobacillus spp.</i> and Enterobacteriales Lower Clostridiales Administration of ampicillin before HCT: <ul style="list-style-type: none"> Increased GVHD severity and lethality Higher <i>Enterococcus spp.</i> and Enterobacteriaceae Lower <i>Blautia spp.</i> abundance Administration of <i>Lactobacillus johnsonii</i> prevented ampicillin-induced effects 		Jenq, 2012 [76]
MHC-disparate allo-HCT using T cells from specific pathogen-free versus germ-free donor	<ul style="list-style-type: none"> T cell donor microbiota presence or absence did not alter: <ul style="list-style-type: none"> T cell differentiation and proliferation GVHD severity GVHD-related mortality 		Tawara, 2013 [86]
MHC-disparate allo-HCT ± intestinal helminth infection	<ul style="list-style-type: none"> Infection with murine nematode <i>Heligmosomoides polygyrus</i>: <ul style="list-style-type: none"> Lowered GVHD severity Preserved GVT effect Increased Treg abundance and improved immune regulation Increased survival of GVHD mice Protective effects of helminthic infection dependent on TGF-β 		Li, 2015 [87]
MHC-matched/minor antigen-mismatched allo-HCT with antianaerobic antibiotics (imipenem-cilastin or piperacillin-tazobactam) versus antibiotics lacking in ant-anaerobic activity (aztreonam)	<ul style="list-style-type: none"> Treatment with anti-anaerobic antibiotics associated with: <ul style="list-style-type: none"> Higher mortality Higher severity of GI GVHD Increased GI inflammatory infiltration Higher levels of IL-23 (mediator of GVHD) Greater abundance of mucin-degrading <i>Akkermansia</i> 		Shono, 2016 [88]
MHC-disparate versus syngeneic allo-HCT 1) +/- intragastric lavage of 17 butyrate-producing strains of <i>Clostridia spp.</i> 2) Anti-anaerobic antibiotics ± <i>Clostridia spp.</i>	<ul style="list-style-type: none"> Gavage with <i>Clostridia spp.</i> resulted in lower GVHD severity and higher survival Anti-anaerobic antibiotics followed by gavage by <i>Clostridia spp.</i> replicated these findings 		Mathewson, 2016 [89]
MHC-disparate allo-HCT + levofloxacin 1) +/- clindamycin 2) Clindamycin ± anti-inflammatory <i>Clostridia spp.</i> (AIC)	<ul style="list-style-type: none"> Treatment with clindamycin decreased survival Clindamycin treatment + AIC increased survival 		Simms-Waldrup, 2017 [90]
B: Microbiota changes associated with GVHD in clinical studies			
Intervention or observational group	Control	Outcomes (Intervention versus control)	References
Randomized trials			
Ciprofloxacin with metronidazole prophylaxis (n = 68)	Ciprofloxacin prophylaxis (n = 66)	<ul style="list-style-type: none"> Lower incidence and severity of acute GVHD No difference in chronic GVHD or OS 	Beelen, 1999 [22]
Observational trials			
Complete GI decontamination (n = 40)	Selective GI decontamination (n = 18)	<ul style="list-style-type: none"> Lower incidence of acute and chronic GVHD Lower rate of infection Lower combined TRM and chronic GVHD 	Vossen, 1990 [26]
Sustained suppression of anaerobic intestinal flora (n = 41)	Incomplete suppression of anaerobic intestinal flora (n = 153)	<ul style="list-style-type: none"> Lower incidence of acute GVHD 	Beelen, 1992 [27]
Acute GVHD (n = 8)	No GVHD (n = 10)	<ul style="list-style-type: none"> Lower intestinal microbiota diversity Higher Lactobacillales Lower Clostridiales 	Jenq, 2012 [75]

(Continued on next page)

Table 7
(continued)

B: Microbiota changes associated with GVHD in clinical studies			
Intervention or observational group	Control	Outcomes (Intervention versus control)	References
Successful GI decontamination (n = 57)	Unsuccessful GI decontamination (n = 55)	<ul style="list-style-type: none"> • Lower incidence of acute GVHD • Lower infectious risk 	Vossen, 2014 [29]
Lowest GI microbial diversity at engraftment (n = 34)	Intermediate or high GI microbial diversity at engraftment (n = 20 intermediate, n = 26 high)	<ul style="list-style-type: none"> • Lower OS • Higher TRM, specifically mortality related to GVHD or infection 	Taur, 2014 [1]
Colonized with <i>Candida</i> in the intestine (n = 54)	Not colonized with <i>Candida</i> in the intestine (n = 99)	<ul style="list-style-type: none"> • Higher incidence of acute GVHD 	van der Velden, 2013 [91]
Acute GI GVHD (n = 8)	No GI GVHD (n = 23)	<ul style="list-style-type: none"> • On the day of transplant: <ul style="list-style-type: none"> • Higher <i>Enterococcus</i> spp. • Lower <i>Clostridia</i> spp. and <i>Eubacterium rectale</i> 	Holler, 2014 [3]
Acute GVHD (n = 5)	No GVHD (n = 5)	<ul style="list-style-type: none"> • Before transplant: <ul style="list-style-type: none"> • Lower diversity • Lower <i>Bacteroides</i> and <i>Parabacteroides</i> spp. • Lower propionate and SCFAs • Between day 0 and day+35: <ul style="list-style-type: none"> • Higher <i>Enterococcus</i> spp. • Lower <i>Faecalibacterium</i> spp • Over all time points: <ul style="list-style-type: none"> • Lower Bacteroidetes 	Biagi, 2015 [92]
Lower microbial diversity (n = 32), lower <i>Blautia</i> abundance (n = 58)	Higher microbial diversity (n = 32), higher <i>Blautia</i> abundance (n = 57)	<ul style="list-style-type: none"> • Lower microbiota diversity and lower abundance of <i>Blautia</i> spp. associated with higher GVHD-related mortality • Higher <i>Blautia</i> abundance associated with higher OS 	Jenq, 2015 [48]
Colonized with ARB pretransplant (n = 33)	Non-ARB colonized (n = 74)	<ul style="list-style-type: none"> • Higher incidence of acute GVHD and acute GI GVHD • Higher rate of bacteremia, increased infection-related mortality • Higher NRM • Lower OS 	Bilinski, 2016 [93]
Treatment of febrile neutropenia with antibiotics effective against anaerobic bacteria: <ul style="list-style-type: none"> • Imipenem-cilastin (n = 148) or • Piperacillin-tazobactam (n = 300) 	Treatment with antibiotics less effective or ineffective against anaerobic bacteria: <ul style="list-style-type: none"> • Cefepime (n = 152) • Aztreonam (n = 64) 	<ul style="list-style-type: none"> • Higher risk of 5-year GVHD-related mortality • No difference in OS 	Shono, 2016 [88]
Acute GVHD (n = 6)	No GVHD (n = 9)	<ul style="list-style-type: none"> • GVHD associated with: <ul style="list-style-type: none"> • Higher cumulative antibiotic exposure and antianaerobic antibiotic exposure, specifically to clindamycin • Higher Enterobacteriaceae, <i>Enterococcus</i> spp., and Neisseriaceae • Lower anti-inflammatory <i>Clostridia</i>, Bacteroidetes, and Actinobacteria • Depletion of <i>Ruminococcus</i> and <i>Blautia</i> spp. 	Simms-Waldrip, 2017 [90]
Pretransplant antibiotic prophylaxis or treatment (n = 239)	No pretransplant antibiotics (n = 261)	<ul style="list-style-type: none"> • Higher incidence and severity of acute GVHD and GI GVHD • Lower median and 10 year OS 	Routy, 2017 [94]
Early pretransplant antibiotics (n = 236)	Late post-transplant antibiotics (n = 297) or no antibiotics (n = 88)	<ul style="list-style-type: none"> • Lower urinary indoxyl sulfate • Lower Clostridiales • Higher acute and/or chronic GVHD-related mortality • Higher TRM with ciprofloxacin/metronidazole versus rifaximin • Lower OS 	Weber, 2017 [54]
Preconditioning low microbial diversity (n = 18)	Preconditioning intermediate (n = 48) and high diversity (n = 41)	<ul style="list-style-type: none"> • Before conditioning: <ul style="list-style-type: none"> • No difference in incidence of acute GVHD or GI GVHD • Higher Firmicutes, and a nonsignificant trend toward lower Bacteroidetes in those who later developed acute GVHD • No difference in OS, relapse, or NRM 	Doki, 2017 [95]

(Continued on next page)

Table 7
(continued)

B: Microbiota changes associated with GVHD in clinical studies			
Intervention or observational group	Control	Outcomes (Intervention versus control)	References
Severe acute GI GVHD (n = 14)	Nonsevere acute GI GVHD or no GVHD (n = 52)	<ul style="list-style-type: none"> • At the time of engraftment: <ul style="list-style-type: none"> • Lower intestinal microbiota diversity • Higher oral Actinobacteria and Firmicutes • Lower Lachnospiraceae • Higher gradient of positively to negatively correlated organisms 	Golob, 2017 [49]
Acute GI GVHD (n = 26)	No GI GVHD (n = 18)	<ul style="list-style-type: none"> • Longitudinal microbiome sampling: <ul style="list-style-type: none"> • Picobirnaviruses predictive of severe enteric GVHD occurrence • Higher picobirnaviruses before or within a week after transplant • Increased rate of detection and number of sequences of persistent DNA viruses over time • No difference in overall richness • Reduced microbial phage richness over time 	Legoff 2017 [50]
Acute GVHD (n = 34) No GVHD (n = 23)	HLA-matched sibling donors (n = 22)	<ul style="list-style-type: none"> • Before conditioning: <ul style="list-style-type: none"> • Lower recipient intestinal microbiota diversity compared with HLA-matched sibling donors • High donor intestinal microbiota diversity is associated with lower acute GVHD incidence in recipients • Low preconditioning intestinal microbiota diversity (in recipients) was not associated with higher risk of acute GVHD • Lower recipient diversity associated with lower OS 	Liu, 2017 [96]

OS indicates overall survival; TRM, transplant related mortality; SCFA, short-chain fatty acids; ARB, antibiotic-resistant bacteria; NRM, nonrelapse mortality.

dominated by Enterobacteriaceae or *Enterococcus spp.*) precedes and significantly increases the risk of bacteremia (with Enterobacteriaceae or *Enterococcus spp.*) in HCT patients [44]. In preclinical models, a 1- to 2-log fold reduction in bacterial [104] or fungal [51] gut colonization levels is sufficient to significantly decrease pathogen dissemination and mortality. Similarly, specific gut commensals can provide resistance to *Clostridium difficile* infection [102]. Thus, efforts targeted at protecting the commensal microbiome may protect against intestinal pathogens and infections.

The role of microbiome-host crosstalk and whether specific molecules or pathways in this cross-talk can be manipulated toward therapeutic benefit remains an active field of investigation. Oral administration of synthetic AMPs may restore gut ecology and shape the host immune system to decrease the risk of infection as well as reducing GVHD, while preserving the graft-versus-leukemia effect. As with nearly all antibacterial agents known to date, resistance to specific AMPs has been described [105]. An alternative strategy that leverages a larger spectrum of AMPs and thus protects against rapid acquisition of resistance might be stimulation of the intrinsic production of AMPs. Although augmenting innate cellular function or mucosal integrity is difficult, it may be possible in the future through modulation of gut microbiota or directly inducing gut mucosal immune effectors to tip the balance back toward gut homeostasis, restore colonization resistance, and reduce the risk of severe infections.

In addition to efforts focused on microbiome modulation to protect against bacterial infections, the importance of microbiota in the control of viral infections and host immune responses has been increasingly recognized. Studies specific to HCT patients or to viruses commonly encountered in HCT

(ie, cytomegalovirus, Epstein-Barr virus, adenovirus, and respiratory syncytial virus) are still very limited. The interactions among microbiota, host immune response, and viral infections are complex and multidirectional: for example, the microbiome may influence viral-specific CD8 T cell memory [106], which can modulate clinical symptoms, severity, and clearance of viral infections [107], or in reverse, a viral infection may result in a change in the microbiome through host-immune responses and changes in cytokines, including interferon [108]. Pathogenic or nonpathogenic viruses within the respiratory tract [109], skin, or gut [110] may also interact with the bacterial microbiome in what has been termed “trans-kingdom interactions” [111]. Host immune responses to prophylactic vaccines or antiviral drugs can also be influenced by the status of the microbiota, and this may impact future decision making around routine decisions such as the schedule for immunizations post-HCT [112].

A proportion of non-relapse-related mortality in HCT patients results from non-GVHD and noninfectious complications for which clinicians are unable to ascribe a clear etiology. These so-called idiopathic disorders may be related to an underlying microbiome dysbiosis or a potential infectious trigger that sparks a self-perpetuating inflammatory cascade (ie, a hit-and-run phenomenon) [113,114]. The application of next-generation sequencing methods and ultrasensitive molecular methods for both unbiased and candidate-base pathogen detection have illuminated several of these “mystery” cases [113,115]. However, recurrent and abundant candidate pathogens have not yet been identified for highly morbid diseases such as the idiopathic pneumonia syndrome. While the evidence is still preliminary in most cases, it is proposed that the microbiome or novel opportunistic pathogens may contribute to these disease phenotypes on occasion.

Table 8
Clinical Trials with Microbiota-Based Interventions in HCT Patients

Name	Comparison	Outcomes	Study type	Clinical Trials.gov ID	Status
Dietary interventions					
Comparing two diets in patients undergoing HSCT or remission induction chemo for acute leukemia and MDS	Standard hospital neutropenic diet versus diet inclusive of fresh fruits and vegetables	Incidence of major infections	Randomized open label	NCT03016130	Recruiting
Gluten-free diet in preventing GVHD in patients undergoing HCT	Gluten-free diet	GVHD	Single arm	NCT03102060	Recruiting
Randomized, prospective, multicenter study to compare enteral nutrition to parenteral nutrition as feeding support in patients with hematologic malignancies undergoing allo-HCT	Enteral versus intravenous nutrition	Mortality	Randomized open label	NCT01955772	Recruiting
Donor human milk in young children receiving bone marrow transplantation	Enteral donor breastmilk versus standard diet	Percentage of stool Lactobacillales	Randomized open label	NCT02470104	Active, not recruiting
Prebiotic and probiotic interventions					
Modification of the intestinal microbiome by diet intervention to mitigate acute GVHD	Bob's Red Mill potato starch-based prebiotic versus standard diet	GVHD	Randomized open label	NCT02763033	Not yet open
FOS in treating patients with blood cancer undergoing HCT	FOS	Prebiotic tolerability	Single arm	NCT02805075	Recruitment completed
<i>Lactobacillus rhamnosus</i> GG in reducing incidence of GVHD in patients who have undergone HCT	<i>L. rhamnosus</i> GG supplementation versus control	GVHD	Randomized open label	NCT02144701	Active, not recruiting
Fecal microbiota transplantation (FMT)					
Auto-FMT for prophylaxis of CDI in recipients of Allo-HCT	Auto-FMT versus control	CDI	Randomized open label	NCT02269150	Recruiting
Fecal transplant for steroid-resistant and steroid-dependent gut acute GVHD	Fecal microbiota transplant	Safety, GVHD	Single arm	NCT03214289	Recruiting
Antibiotic interventions					
Choosing the best antibiotic to protect friendly gut bacteria during the course of HCT	Piperacillin-tazobactam versus Cefepime during febrile neutropenia	Change in <i>Clostridial</i> abundance	Randomized open label	NCT03078010	Recruiting
Gut decontamination in pediatric allo-HCT	Oral Vancomycin-Polymixin B versus standard of care	Intestinal microbiota diversity	Randomized open label	NCT02641236	Recruiting

FOS indicates fructo-oligosaccharides; CDI, *Clostridium difficile* infection.

The role of the microbiome in modifying the incidence and clinical outcomes of infection and idiopathic disorders in HCT patients is becoming increasingly recognized. To date, most research has focused on the bacterial contribution to these disorders, but increasingly, there is an appreciation of the contribution of viruses to these disorders and to the delicate balance of the microbiome. Both host and microbial factors participate in a complicated interplay to maintain homeostasis, and we are just now starting to understand the detailed elements in this complicated interaction. Little is known about the fungal contribution to both the healthy and diseased HCT microbiome, although we anticipate this will be an area of active and productive research in the future.

Open Questions

- What is the composition of the human “virome” and “mycobiome” in HCT patients—and how do interactions among viruses, fungi, bacteria, and the host impact HCT outcomes?
- Do antimicrobial prophylaxis strategies adversely impact the microbiota and render HCT recipients susceptible to opportunistic infections beyond *C. difficile*?
- Might microbiome-targeted therapeutics, aimed at protecting against loss of diversity in the microbiome, decrease

the rate of infectious complications such as enteric Gram-negative and Gram-positive bacteremia believed to originate from the intestinal microbiome?

- Which of the idiopathic complications of HCT are related to either infections or microbial dysbiosis? In cases where the offending organism acts through a hit-and-run type of mechanism, how might we identify these etiologies using existing technologies and sampling strategies?

METHODS FOR MICROBIOTA MODIFICATION

A clear rationale exists for targeting the microbiome with the eventual intention of both fine-tuning the immune system (balancing GVHD and GVT, for example) and decreasing the risk of downstream infectious complications of HCT. Several interventional studies are ongoing that will alter microbiota by means of diet and prebiotics, antibiotics, probiotics, microbial metabolites, and fecal microbial transplantation (Table 8). Subsequently we discuss several clinical microbiome manipulation strategies and the implications of their use in future studies.

Antibiotics

Over the past 10 years, metagenomic and other culture-independent microbiota analyses have demonstrated the important role of the microbiome in health and disease [116].

Table 9
Clinical Trials Investigating the Efficacy of a Neutropenic Diet in HCT Patients

Intervention	Control	Outcomes (Intervention versus control)	References
Randomized controlled, prospective trial Unrestricted diet (n = 21)	Neutropenic diet (n = 25)	<ul style="list-style-type: none"> • No difference in infectious outcomes • No difference in nutritional status 	Lassiter, 2015 [120]
Retrospective observational study General hospital diet (n = 363)	Neutropenic diet (n = 363)	<ul style="list-style-type: none"> • Fewer microbiologically confirmed infections in those receiving a general diet • No difference in the incidence of microbiologically confirmed infections during neutropenia • Higher rate of infections after resolution of neutropenia in those receiving a neutropenic diet 	Triflilio, 2012 [121]

Patients undergoing HCT represent a natural group for this line of research for the reasons that they are (1) uniquely prone to perturbations in the normal microbiome as a result of toxicity from conditioning regimens, impaired diet, and antimicrobial exposure given for treatment and prophylaxis; (2) their propensity for infectious and immune-mediated morbidity and mortality; (3) their prolonged peritransplant hospitalization that facilitates convenient sampling; and (4) the availability of long-term outcomes that are universally gathered from transplant recipients. Early studies of prophylactic antibiotics in HCT demonstrated reductions in both infections and GVHD after suppression of the microbial flora [26]. Although the use of broad-spectrum antibiotics has led to dramatic improvement in infection-related TRM, antibiotics result in substantial microbiota disruption [53]. Importantly, the type of antibiotic therapy may determine the composition of intestinal microbiota and the extent of microbiome disruption. For example, antibiotics with anaerobic activity are associated with higher rates of GVHD-related mortality [87]. These new insights, which suggest an unfavorable impact of broad-spectrum antibiotics on intestinal microbiota and patient outcomes after HCT, raise the question of how we might preserve the protective effects of “healthy” commensal organisms without compromising treatment efficacy. In addition to the type of antibiotic, timing of treatment also appears to influence microbial diversity and may impact patient outcomes. Patients starting antibiotics before their day of transplantation showed significantly more microbiome disruption and had a higher TRM than did those who began antibiotics on or following day 0 or who did not receive antibiotics [53]. Collectively, these studies support an argument for more selective use of broad-spectrum antibiotics along with early de-escalation strategies. Such strategies would preserve the microbiome but still ensure adequate prevention and treatment of bacterial infections. Further, the benefit of gut decontamination and prophylactic antibiotics should be examined through well-designed prospective trials. Definitive support for or against these practices will only come through the conduct of multicenter prospective trials designed to assess the short-term risk of bacterial infections during neutropenia with long-term endpoints (GVHD, immune reconstitution, and microbial resistance) that may be affected by disruption of microbiome diversity.

Open question(s)

- How do we balance adequate prevention and treatment of bacterial infections with preservation of the microbiome in HCT recipients?

- How can we incorporate microbiota stewardship practices in addition to antibiotic stewardship practices in our care of HCT recipients?

Diet, Prebiotics, Probiotics, and Fecal Microbiota Transplantation

Ingested food contaminated by microbes has long been recognized as a potential source of bloodstream infection during chemotherapy-induced neutropenia with attendant gastrointestinal mucosal damage. The germ-free “sterile” diet was conceived in the 1960s as a way to reduce ingestion of potentially harmful microbes, but this was not palatable [117]. A “cooked-food” diet alternative, which eliminated raw foods with high bacterial counts, was shown in a randomized trial to have a similar effect on bacterial stool cultures as the germ-free diet, but it was also limited by patient dissatisfaction [117]. To expand and improve food palatability, Pizzo et al. [118] cultured commercially available foods and identified low-microbial foods that were deemed suitable for a neutropenic diet. The composition of neutropenic diets vary from center to center but, in general, consist of cooked and canned food products and exclude raw meat, fresh fruits, juices and vegetables, raw eggs, and unpasteurized dairy products [117,119]. Despite limited evidence to support the merits of a neutropenic diet in HCT recipients as illustrated in Table 9 [120,121], dietary restriction of fresh fruits and vegetables continues to be standard practice for neutropenic patients in some centers, which likely has an impact on the amount of fiber that is consumed by HCT patients [122]. In the early post-transplant period, nutritional oral intake often declines to the point of necessitating nutritional supplementation. Retrospective comparisons of parenteral and enteral nutrition have suggested a benefit to the enteral route [123,124]. This may be due to enteral nutrition maintaining digestive function and the mucosal barrier, thus preventing bacterial translocation [125]. An ongoing trial is currently evaluating enteral versus parenteral nutrition [126].

Specific elements of diet, called prebiotics, are particularly influential in the structure and function of the microbiota. The term “prebiotic” is traditionally applied to indigestible carbohydrates that are metabolized by gut bacteria to produce short-chain fatty acids; recently the term is being redefined to refer to any substrate that is selectively utilized by host microorganisms and that confers a health benefit [127]. Apart from a single retrospective study, little is known about how prebiotics affect transplant outcomes such as GVHD [128,129]. However, studies in patients with inflammatory bowel disease treated with prebiotics such as inulin and fructooligosaccharides (inulin-type fructans) have demonstrated that

Table 10

Probiotics and the Microbiome in Murine Models of HCT and Probiotics and the Microbiome in Clinical Studies of HCT Patients

A: Probiotics and the microbiome in murine models of HCT			
Treatment	Control	Outcomes	References
<i>Lactobacillus rhamnosus</i> GG	Ciprofloxacin	<ul style="list-style-type: none"> • Lower mortality • Lower GVHD incidence 	Gerbitz, 2004 [136]
17 butyrate-producing <i>Clostridia</i> spp. strains Anti-inflammatory <i>Clostridia</i> spp.	Phosphate-buffered saline Phosphate-buffered saline	<ul style="list-style-type: none"> • Lower GVHD severity • Higher survival 	Mathewson, 2016 [88] Simms-Waldrup, 2017 [89]
B: Probiotics and the microbiome in clinical studies of HCT patients			
Trial design	Probiotic, dose	Outcomes	References
Observational	Self-reported yogurt intake, average of 150 g/day (n = 41)	<ul style="list-style-type: none"> • Higher yogurt intake associated with more rapid neutrophil engraftment 	Tavil, 2012 [137]
Single-arm	<i>Lactobacillus plantarum</i> , 1 × 10 ¹³ CFU/kg/day (n = 30)	<ul style="list-style-type: none"> • 97% of the children received at least half of the probiotic doses • No incidence of <i>Lactobacillus</i> bacteremia reported 	Ladas, 2016 [138]
Randomized	<i>Lactobacillus rhamnosus</i> GG, 1 × 10 ⁹ /day (probiotic group, n = 20; control group, n = 11)	<ul style="list-style-type: none"> • No difference in gut microbiota diversity • No difference in GVHD incidence 	Gorshein, 2017 [139]

CFU, colony-forming units.

these agents increase microbiota diversity and are associated with a corresponding decrease in disease markers and activity [130,131]. Such studies provide a compelling rationale for studies of prebiotics in the HCT setting.

Probiotics are live microorganisms given to improve health and have long been used as part of traditional diets through the ingestion of fermented foods. Encapsulated preparations of 1 or more isolated live organisms have been used in attempts to treat a wide variety of gastrointestinal illnesses, including infectious diarrhea or gastroenteritis [132], and inflammatory bowel disease [89,133]. Some of these studies have shown evidence of efficacy, most likely mediated through direct antimicrobial effects, stimulation of immune responses that lead to upregulation of anti-inflammatory cytokines and IgA, and promotion of intestinal barrier function [134,135]. To date, probiotics in the HCT setting have been limited to preclinical models and small pilot trials (Tables 10A and 10B) [88,89,136–139]. Of course, concern exists for the potential infectious complications associated with administration of live microbial organisms in high dose [140]. Indeed, case reports of bacteremia following ingestion of probiotics suggest the importance of exercising caution and judgment in the use of live bacterial therapies [141]. Further clinical studies are needed to fully determine the safety and efficacy of probiotics in patients undergoing HCT.

Finally, fecal microbiota transplantation (FMT) is yet another intervention that could be employed to preserve or restore the gastrointestinal microbiota in patients undergoing HCT. Pioneering physicians performed FMT in non-HCT patients with recurrent or refractory *Clostridium difficile* infection (CDI) and demonstrated efficacy in up to 90% of treated patients [142,143]. Literature on FMT in HCT patients is still scant; however, the limited data to date appear encouraging, with a total of 25 reported HCT patients having undergone FMT without known complications (Tables 11A–C) [144–151]. Highlighting the need for a cautious approach in HCT populations, case reports in non-HCT patients with CDI have documented infectious complications following FMT including norovirus infection and sepsis as a result of presumed bacterial translocation [152,153]. Taken together, while limited published experience suggests that FMT can be used in immunocompromised patients with CDI [154], prospective trials evaluating safety and efficacy in HCT recipients are needed.

Undeniably, the field of FMT is rapidly growing, yet several questions remain including what guidelines the field should adopt for identification of the best FMT donors and appropriate donor stool screening before FMT in immunocompromised patients. Autologous FMT donation may have an advantage of simple traceability of the preparation and control of the inoculum during donor procedures, as well as reducing the risk of potential transmission of diseases originating from the microbiota of an external donor. The opportunity to obtain a “premorbid/baseline” stool may not always be feasible for autologous FMT, and for this reason several investigators are using third-party FMT obtained from healthy donors [155]. Case reports have explored the role of FMT for the treatment of noninfectious complications. Finally, it is interesting to speculate about the role of the stem cell donor’s microbiota [85,95], or the patient’s cohabitating family members [156] from whom the microbiota may potentially reconstitute after transplant-induced dysbiosis.

Open questions

- What is the impact of neutropenic dietary restriction and use of parenteral nutrition on long-term outcomes post-HCT?
- How might the screening protocol for FMT donors to HCT patients differ than standard screening protocols used for less immunocompromised patients?
- Can targeting of specific microbes or microbial pathways result in modification of microbe-disease associations?
- Is there a role for genetically modified bacteria in the post-HCT setting? How might this tool be safely and effectively leveraged?
- Are there novel bacterial natural products (small molecules, proteins, glycolipids, sugars) that can be investigated for salutary drug-like effects in the HCT patient population?
- Might we create a defined microbial consortium as the next-generation microbial therapeutic to supplant FMT in the HCT recipient population?
- How stable are microbial populations and microbial genomes over time? How does horizontal gene transfer affect the medium- and long-term safety and efficacy of potential novel bacterial therapies?

Table 11

A. Fecal Microbiota Transplantation in Auto- and Allo-HCT Patients with Recurrent CDI, Fecal Microbiota transplantation in Allo-HCT Patients with GVHD, and Fecal Microbiota Transplantation in Allo-HCT Patients with antibiotic-resistant bacteria

A: Fecal microbiota transplantation in auto- and allo-HCT patients with recurrent CDI						
Patient population	Total patients	Route	Donor type	Outcomes	Adverse events	References
21-year-old woman; allo-HCT	1	NG tube	Spouse	Resolution of CDI	None	Neemann, 2012 [144]
60-year-old woman; allo-HCT	1	Push enteroscopy	Anonymous, 2 donors	Resolution of CDI	None	de Castro, 2015 [145]
64-year-old man; auto-HCT	1	Enema	Anonymous, 2 sequential FMTs from different donors	First FMT resulted in resolution of CDI. Recurrence in 6 months. Second FMT performed at 6 months resulted in resolution of CDI durable for at least 7 months	None	Mittal, 2015 [146]
Allo-HCT patients	7	NJ tube or colonoscopy	Anonymous	Resolution of CDI in 6 patients; 1 patient recurred at day +156 post-FMT after receiving antibiotics. Repeat FMT with resolution of symptoms lasting >4 months	None	Webb, 2016 [147]
Auto- and allo-HCT patients	8	Oral capsules	Anonymous	No recurrence of disease in 7 patients; 1 had recurrence of CDI day +179 after FMT	None	Moss, 2017 [148]
B: Fecal microbiota transplantation in allo-HCT patients with GVHD						
Steroid-resistant GI GVHD	4	ND tube	Related	Resolution of diarrhea in 3 patients, partial resolution in 1 patient	None	Kakihana, 2016 [149]
Steroid-refractory GI GVHD	3	Colonoscopy	Related or anonymous	Resolution of diarrhea in 2 patients, improvement in GI GVHD in third patient	None	Spindelboeck, 2017 [150]
C: Fecal microbiota transplantation in allo-HCT patients with antibiotic-resistant bacteria						
Hematologic disorders + ARB colonized	20 (8 HCT)	ND tube	Anonymous	<ul style="list-style-type: none"> 75% of patients (60% of FMTs) achieved complete ARB decolonization at 1 month 93% of FMTs achieved complete ARB decolonization at 6 months 	<ul style="list-style-type: none"> Vomiting (n = 1) Transient, grade 1 diarrhea within 3 days (n = 25) Abdominal pain (n = 2) Ileus (n = 2) 	Bilinski, 2017 [151]

CDI indicates *Clostridium difficile* infection; NG, nasogastric; NJ, nasojejunal; ND, nasoduodenal; ARB, antibiotic-resistant bacteria.

MAXIMIZING THE OPPORTUNITY FOR IMPACT

Identifying associations between the microbiome and clinical phenotypes is critically important. With the advent of technologies such as metagenomic sequencing, metabolomics, and improved tools for studying human immunology, it is becoming increasingly affordable and feasible to perform longitudinal molecular characterization of patients following HCT. As we transition toward an increased reliance on human samples for the generation and querying of biological hypotheses, it is important that the same rigor used in carefully controlled *in vitro* or animal experiments be applied in the clinical setting. This is particularly important in light of the sometimes conflicting results seen between preclinical and clinical studies, possibly as a result of microbiota variability between animal strains or differences in practice between centers, but also potentially as a result of variation in sample management. Samples must be collected, stored, and processed in a reproducible manner to avoid the unintended introduction of bias in the results [157]. This is of utmost importance when studying low-biomass or low-microbial-burden samples, where the chance introduction of ambient microorganisms through handling or processing may confound the ability to draw robust, reproducible, and generalizable conclusions. Similar to standard sample collection and data generation practices, strict procedures for management of data generated through high-throughput techniques

will ensure data quality and accuracy critical to reliable interpretation and analysis. For example, current efforts to understand the microbiome and its impact on transplant outcomes are often limited by incomplete information on antibiotic exposure and diet. The bone and marrow transplant community has a strong track record of collecting detailed clinical data regarding GVHD and immunosuppressive medication administration. This same rigor needs to be applied to infection reporting and antimicrobial exposure to draw meaningful conclusions regarding microbiome findings. To this end, the Center for International Blood and Marrow Transplant Research currently has a working group tasked with the development of reporting standards for infectious disease endpoints.

As clinical infectious data collection strategies improve, inclusion of patient-reported outcomes (PROs) should also be prioritized in microbiome-oriented studies. PROs provide greater accuracy of treatment-related symptoms than clinician report [158], and given the role that the microbiome likely plays in mediating symptoms such as diarrhea, constipation, flatus, and abdominal discomfort, inclusion of PROs will be critical in studying the impact of microbiome-modifying therapies. Specifically, validated instruments of physical function and health-related quality of life are available and include patient-reported measures of gastrointestinal symptoms, with domains of severity, frequency, and interference related to




Figure 1. Structuring the design of future interventions aimed at establishing microbe-host disease causality.

these symptoms contained within the NCI PRO–common terminology criteria for adverse events (<http://healthcare.delivery.cancer.gov/pro-ctcae>). The successful incorporation of PROs will require (1) the development of a scalable infrastructure for participating sites; (2) consensus choice of relevant measures [159]; (3) collection time points so that PROs can be appropriately linked to clinical data consistently across research studies and clinical practice; and (4) application of statistically sound approaches to handle missing data is of paramount importance.

It is expected that the collection of high-quality high-resolution molecular, clinical, and patient-reported symptom data will hasten the identification of potentially relevant and useful molecular biomarkers as well as associations between microbiome alterations or specific microbes and clinically relevant outcomes. We anticipate that “big-data” approaches, including clinical-informatics efforts that extract information from the electronic medical record, will be a key part of this effort. As more and more data are collected, it will be increasingly important to leverage machine-learning approaches to data interpretation and analysis, as is already being done in the field of cancer genomics and beyond.

Perhaps most importantly, we must not lose sight of the importance of moving “beyond association” in study design. Strategies to model the hypothesized interactions between the microbiome and host, both in vitro and in vivo in models ranging from cell lines to small and large animal models, will allow for carefully controlled experiments to be carried out—something that is of course difficult or challenged by ethical considerations in patient studies. Last, whenever possible, it is critical that well designed and thoughtfully targeted interventional studies of microbiome modification be performed in multicenter settings. This will provide the highest level of prospective data to help guide clinical practice, and the multicenter nature of these studies will ensure the highest level of generalizability. When these studies are done, adhering to rigorous standards of biospecimen collection, clinical data collection, and PRO data collection will allow for the ac-

curate measurement of the consequences of the tested interventions. This will shed light on potential mechanisms of action of these interventions, and will inform the next iteration of interventions (Figure 1).

Open Questions

- How can we facilitate more universal sample collection and support multi-institutional studies of the microbiome to improve the generalizability of findings?
- How do we identify and implement standardized methods for sample and data collection that can be used for multicenter prospective clinical trials that study the microbiome in HCT patients?
- How can we best collect information about infections and antimicrobial medication use in both the clinical trial setting and for registry purposes?
- What are the critical clinical data that need to be collected for meaningful analysis of microbiome studies in HCT?
- How can we hasten the design and execution of microbiome-targeted interventional clinical trials? What funding sources exist to support these critically important efforts, given the relative paucity of classical industry partners in this space?

FUTURE DIRECTIONS

While the importance of the microbiome in immunological development, protection against infections, and patient symptoms has been investigated in the past decades, much remains to be understood on the importance and relevance of the contribution of the stability, the resilience and the redundancy of the microbial composition after transplantation. The field has come a long way from the earliest days of microbiome research, now nearly a half a century ago. There has been an explosion in the number of high-throughput tools for microbiome measurement and an increasing number of single-institution biospecimen collections. These tools and resources have facilitated the testing of only a limited number of translationally important hypotheses. The taxonomic

diversity of the microbiome has been shown to be a potential biomarker of HCT outcomes, ranging from overall survival to relapse. These early, single-institution findings are certainly compelling, and warrant further investigation. Interactions between the microbiome and immune system have been described for decades; deep immunophenotyping tools such as TCR sequencing and high-dimensional mass cytometry are now facilitating investigating the temporal relationships between the microbiome and immune system during immune reconstitution. Early data suggest a potential role for the microbiome in improving post-transplant immune reconstitution and helping to achieve the elusive goal of effective GVT without GVHD. A growing set of tools for microbiome manipulation using diet, prebiotics, probiotics, and even FMT are being tested rigorously, both in preclinical models and in humans. It is anticipated that larger randomized multi-institutional studies of these approaches and their efficacy will be initiated. Beyond GVHD, the clear role of the microbiome in mediating risk of infection and perhaps idiopathic disorders also poses an exciting opportunity for investigation, and the potential to improve non-relapse-, non-GVHD-related morbidity and mortality. To maximize the impact of microbiome-focused investigation, there are many targets that represent low-hanging fruit: improving infection and antibiotic-related data collection, incorporating PROs, and the application of newer methods such as shotgun sequencing, metabolomics, metaproteomics, advanced microscopy, and beyond, that will allow us to extend our investigational reach beyond the taxonomic realm. A more functional characterization of how these communities are structured, how they interact and what they do will undoubtedly inform progress in developing precision microbiome diagnostic and therapeutic strategies. Advances in technology have revealed many opportunities to better understand the mechanisms that underlie microbiome-host interactions. For example, recent studies have brought to the fore the role of metabolomes and host genes that are critical [88], yet, there have been so far no corresponding studies on the RNA transcripts (metatranscriptomics) and proteins produced by the microbes in these processes in HCT patients. The abundance and transcript levels of genes encoding microbial resistance to antibiotics, drug metabolism, and resistance to host mediated immune responses, for example, could shed light on better exploitation of the microbiome in HCT. Advances in bioengineering have resulted in the ability to generate microbes with specific, salutary effects. For example, oral administration of commensal bacteria genetically engineered to regulate endogenous or recombinant gene expression to alter their metabolic ability could hold great promise for restoring intestinal homeostasis and modulating host immune systems [160]. New technologies are rapidly being developed and applied; thus, in the coming years research will better define the role of microbiome on GVHD, GVT, infectious complications, and transplant outcomes. As this happens, we hope that carefully considered and planned investigations ranging from basic microbiology to immunology to large-scale, randomized-controlled interventional clinical trials will together result in improved outcomes for HCT and related patient populations. Simultaneously, we anticipate that such efforts will result in an improved understanding of the basic biological underpinnings of microbial bioregulation, microbiome community interactions, and human immunology. Vast opportunities exist for both scientific and translational advances in the realm of microbiome sciences. To capitalize

quickly on these prospects for maximum impact, we propose 10 areas of focus (Table 1) that may have the greatest promise for breakthrough discoveries regarding the dynamic and complex microbiota-host relationship in patients undergoing HCT.

ACKNOWLEDGMENTS

Funding: TMA is supported by the American Society of Blood and Marrow Transplantation New Investigator Award and by the NIH/NCATS Clinical and Translational Science Award KL2 TR000083 and UL1 TR001085. ASB is supported by the Damon Runyon Cancer Research Foundation. RRJ is supported by NIH/NHLBI RO1 HL124112 and CPRIT RR160089. JUP, MAP, and MvdB are supported in part by an NIH P01 CA23766 and NIH/NCI P30 CA 008748. The Blood and Marrow Transplant Clinical Trials Network is supported in part by grants #U10HL069294 and #U24HL138660 from the National Heart, Lung, and Blood Institute and the National Cancer Institute. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Financial disclosure: The authors have nothing to disclose.

Conflict of interest statement: J.U.P receives licensing fees and research support from Seres Therapeutics. M.v.d.B receives research support from Seres Therapeutics and has served as a consultant for Jazz Pharmaceuticals, Novartis, Regeneron Pharmaceuticals, Flagship Ventures, Boehringer Ingelheim, Merck, and Evolo. R.R.J is on the advisory committee of and holds patents with or receives royalties from Seres Therapeutics; he has consulted for Ziopharm Oncology. L.S.K consults with and receives funding from Bristol Myers Squibbs and Kymab, Ltd; she receives research funding from Regeneron Pharmaceuticals. M.M.M has received honoraria and research support from MaaT Pharma.

REFERENCES

1. Taur Y, Jenq RR, Perales M-A, et al. The effects of intestinal tract bacterial diversity on mortality following allogeneic hematopoietic stem cell transplantation. *Blood*. 2014;124(7):1174–1182.
2. Peled JU, Devlin SM, Staffas A, et al. Intestinal microbiota and relapse after hematopoietic-cell transplantation. *J Clin Oncol*. 2017;35(15):1650–1659.
3. Holler E, Butzhammer P, Schmid K, et al. Metagenomic analysis of the stool microbiome in patients receiving allogeneic stem cell transplantation: loss of diversity is associated with use of systemic antibiotics and more pronounced in gastrointestinal graft-versus-host disease. *Biol Blood Marrow Transplant*. 2014;20(5):640–645.
4. Connell MS, Wilson R. The treatment of x-irradiated germfree Cfw and C3h mice with isologous and homologous bone marrow. *Life Sci*. 1965;4:721–729.
5. Jones JM, Wilson R, Bealmear PM. Mortality and gross pathology of secondary disease in germfree mouse radiation chimeras. *Radiat Res*. 1971;45(3):577–588.
6. van Bekkum D, Vos O. Treatment of secondary disease in radiation chimaeras. *Int J Radiat Biol*. 1961;3:173–181.
7. van Bekkum DW, Knaan S. Role of bacterial microflora in development of intestinal lesions from graft-versus-host reaction. *J Natl Cancer Inst*. 1977;58(3):787–790.
8. Pollard M, Truitt RL. Allogeneic bone marrow chimerism in germfree mice. I. Prevention of spontaneous leukemia in AKR mice. *Proc Soc Exp Biol Med*. 1973;144(2):659–665.
9. Pollard M, Truitt RL. Allogeneic bone marrow chimerism in germ-free mice. II. Prevention of reticulum cell sarcomas in SJL-J mice. *Proc Soc Exp Biol Med*. 1974;145(2):488–492.
10. Truitt RL, Pollard M, Srivastava KK. Allogeneic bone marrow chimerism in germfree mice. 3. Therapy of leukemic AKR mice. *Proc Soc Exp Biol Med*. 1974;146(1):153–158.
11. Truitt RL, Pollard M. Allogeneic bone marrow chimerism in germ-free mice. IV. Therapy of “Hodgkin’s-like” reticulum cell sarcoma in SJL mice. *Transplantation*. 1976;21(1):12–16.
12. van Bekkum DW, Roodenburg J, Heidt PJ, van der Waaij D. Mitigation of secondary disease of allogeneic mouse radiation chimeras by

- modification of the intestinal microflora. *J Natl Cancer Inst.* 1974; 52(2):401-404.
13. Heit H, Wilson R, Flidner TM, Kohne E. Mortality of secondary disease in antibiotic-treated mouse radiation chimeras. In: Heneghan JB, ed. *Germ Free Research: Biological Effects and Gnotobiotic Environment*. New York: Academic Press; 1973:477-485.
 14. van Bekkum DW, de Vries MJ, van der Waay D. Lesions characteristic of secondary disease in germfree heterologous radiation chimeras. *J Natl Cancer Inst.* 1967;38(2):223-231.
 15. Heit H, Heit W, Kohne E, Flidner TM, Hughes P. Allogeneic bone marrow transplantation in conventional mice: I. Effect of antibiotic therapy on long term survival of allogeneic chimeras. *Blut.* 1977;35(2):143-153.
 16. Veenendaal D, de Boer F, Van der Waaij D. Effect of selective decontamination of the digestive tract of donor and recipient on the occurrence of murine delayed-type graft-versus-host disease. *Med Microbiol Immunol (Berl)*. 1988;177(3):133-144.
 17. Buckner CD, Clift RA, Sanders JE, et al. Protective environment for marrow transplant recipients: a prospective study. *Ann Intern Med.* 1978;89(6):893-901.
 18. Navari RM, Buckner CD, Clift RA, et al. Prophylaxis of infection in patients with aplastic anemia receiving allogeneic marrow transplants. *Am J Med.* 1984;76(4):564-572.
 19. Petersen FB, Buckner CD, Clift RA, et al. Laminar air flow isolation and decontamination: a prospective randomized study of the effects of prophylactic systemic antibiotics in bone marrow transplant patients. *Infection.* 1986;14(3):115-121.
 20. Petersen FB, Buckner CD, Clift RA, et al. Prevention of nosocomial infections in marrow transplant patients: a prospective randomized comparison of systemic antibiotics versus granulocyte transfusions. *Infect Control.* 1986;7(12):586-592.
 21. Petersen FB, Buckner CD, Clift RA, et al. Infectious complications in patients undergoing marrow transplantation: a prospective randomized study of the additional effect of decontamination and laminar air flow isolation among patients receiving prophylactic systemic antibiotics. *Scand J Infect Dis.* 1987;19(5):559-567.
 22. Beelen DW, Elmaagacli A, Müller K-D, Hirche H, Schaefer UW. Influence of intestinal bacterial decontamination using metronidazole and ciprofloxacin or ciprofloxacin alone on the development of acute graft-versus-host disease after marrow transplantation in patients with hematologic malignancies: final results and long-term follow-up of an open-label prospective randomized trial. *Blood.* 1999;93(10):3267-3275.
 23. Storb R, Prentice RL, Buckner CD, et al. Graft-versus-host disease and survival in patients with aplastic anemia treated by marrow grafts from HLA-identical siblings. Beneficial effect of a protective environment. *N Engl J Med.* 1983;308(6):302-307.
 24. Mahmoud HK, Schaefer UW, Schünig F, et al. Laminar air flow versus barrier nursing in marrow transplant recipients. *Blut.* 1984;49(5):375-381.
 25. Schmeiser T, Kurrle E, Arnold R, Krieger D, Heit W, Heimpel H. Antimicrobial prophylaxis in neutropenic patients after bone marrow transplantation. *Infection.* 1988;16(1):19-24.
 26. Vossen JM, Heidt PJ, van den Berg H, Gerritsen EJ, Hermans J, Dooren LJ. Prevention of infection and graft-versus-host disease by suppression of intestinal microflora in children treated with allogeneic bone marrow transplantation. *Eur J Clin Microbiol Infect Dis.* 1990;9(1):14-23.
 27. Beelen DW, Haralambie E, Brandt H, et al. Evidence that sustained growth suppression of intestinal anaerobic bacteria reduces the risk of acute graft-versus-host disease after sibling marrow transplantation. *Blood.* 1992;80(10):2668-2676.
 28. Passweg JR, Rowlings PA, Atkinson KA, et al. Influence of protective isolation on outcome of allogeneic bone marrow transplantation for leukemia. *Bone Marrow Transplant.* 1998;21(12):1231-1238.
 29. Vossen JM, Guiot HFL, Lankester AC, et al. Complete suppression of the gut microbiome prevents acute graft-versus-host disease following allogeneic bone marrow transplantation. *PLoS ONE.* 2014;9(9):doi:10.1371/journal.pone.0105706. e105706.
 30. Tomblyn M, Chiller T, Einsele H, et al. Guidelines for preventing infectious complications among hematopoietic cell transplantation recipients: a global perspective. *Biol Blood Marrow Transplant.* 2009;15(10):1143-1238.
 31. Marchesi JR, Ravel J. The vocabulary of microbiome research: a proposal. *Microbiome.* 2015;3:31.
 32. Weisburg WG, Barns SM, Pelletier DA, Lane DJ. 16S ribosomal DNA amplification for phylogenetic study. *J Bacteriol.* 1991;173(2):697-703.
 33. Loman NJ, Constantinidou C, Chan JZM, et al. High-throughput bacterial genome sequencing: an embarrassment of choice, a world of opportunity. *Nat Rev Microbiol.* 2012;10(9):599-606.
 34. Manzo VE, Bhatt AS. The human microbiome in hematopoiesis and hematologic disorders. *Blood.* 2015;126(3):311-318.
 35. Lozupone CA, Stombaugh JI, Gordon JL, Jansson JK, Knight R. Diversity, stability and resilience of the human gut microbiota. *Nature.* 2012;489(7415):220-230.
 36. Petersen C, Round JL. Defining dysbiosis and its influence on host immunity and disease. *Cell Microbiol.* 2014;16(7):1024-1033.
 37. Giraud A. Axenic mice model. *Methods Mol Biol.* 2008;415:321-336.
 38. Borody TJ, Paramsothy S, Agrawal G. Fecal microbiota transplantation: indications, methods, evidence, and future directions. *Curr Gastroenterol Rep.* 2013;15(8):337.
 39. Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R, Gordon JL. The human microbiome project. *Nature.* 2007;449(7164):804-810.
 40. Integrative HMP (iHMP) Research Network Consortium. The integrative human microbiome project: dynamic analysis of microbiome-host omics profiles during periods of human health and disease. *Cell Host Microbe.* 2014;16(3):276-289.
 41. Auchtung JM, Robinson CD, Britton RA. Cultivation of stable, reproducible microbial communities from different fecal donors using minibioreactor arrays (MBRAs). *Microbiome.* 2015;3:42.
 42. Kim HJ, Li H, Collins JJ, Ingber DE. Contributions of microbiome and mechanical deformation to intestinal bacterial overgrowth and inflammation in a human gut-on-a-chip. *Proc Natl Acad Sci USA.* 2016;113(1):E7-E15.
 43. Rajilić-Stojanović M, Heilig GHJ, Molenaar D, et al. Development and application of the human intestinal tract chip, a phylogenetic microarray: analysis of universally conserved phylotypes in the abundant microbiota of young and elderly adults. *Environ Microbiol.* 2009;11(7):1736-1751.
 44. Taur Y, Xavier JB, Lipuma L, et al. Intestinal domination and the risk of bacteremia in patients undergoing allogeneic hematopoietic stem cell transplantation. *Clin Infect Dis.* 2012;55(7):905-914.
 45. Levine JE, Huber E, Hammer ST, et al. Low Paneth cell numbers at onset of gastrointestinal graft-versus-host disease identify patients at high risk for nonrelapse mortality. *Blood.* 2013;122(8):1505-1509.
 46. Weber D, Oefner PJ, Hiergeist A, et al. Low urinary indoxyl sulfate levels early after transplantation reflect a disrupted microbiome and are associated with poor outcome. *Blood.* 2015;126(14):1723-1728.
 47. Jenq RR, Taur Y, Devlin SM, et al. Intestinal blautia is associated with reduced death from graft-versus-host disease. *Biol Blood Marrow Transplant.* 2015;21(8):1373-1383.
 48. Golob JL, Pergam SA, Srinivasan S, et al. Stool microbiota at neutrophil recovery is predictive for severe acute graft vs host disease after hematopoietic cell transplantation. *Clin Infect Dis.* 2017;65(12):1984-1991.
 49. Legoff J, Resche-Rigon M, Bouquet J, et al. The eukaryotic gut virome in hematopoietic stem cell transplantation: new clues in enteric graft-versus-host disease. *Nat Med.* 2017;23(9):1080-1085.
 50. Niwa T. Uremic toxicity of indoxyl sulfate. *Nagoya J Med Sci.* 2010;72(1-2):1-11.
 51. Fan D, Coughlin LA, Neubauer MM, et al. Activation of HIF-1 α and LL-37 by commensal bacteria inhibits *Candida albicans* colonization. *Nat Med.* 2015;21(7):808-814.
 52. Lee J-H, Lee J. Indole as an intercellular signal in microbial communities. *FEMS Microbiol Rev.* 2010;34(4):426-444.
 53. Weber D, Jenq RR, Peled JU, et al. Microbiota disruption induced by early use of broad-spectrum antibiotics is an independent risk factor of outcome after allogeneic stem cell transplantation. *Biol Blood Marrow Transplant.* 2017;23(5):845-852.
 54. Gomez de Agüero M, Ganai-Vonarburg SC, Fuhrer T, et al. The maternal microbiota drives early postnatal innate immune development. *Science.* 2016;351(6279):1296-1302.
 55. Arrieta M-C, Stiemsma LT, Dimitriu PA, et al. Early infancy microbial and metabolic alterations affect risk of childhood asthma. *Sci Transl Med.* 2015;7(307):307ra152.
 56. Shaw SY, Blanchard JF, Bernstein CN. Association between the use of antibiotics in the first year of life and pediatric inflammatory bowel disease. *Am J Gastroenterol.* 2010;105(12):2687-2692.
 57. Dominguez-Bello MG, Costello EK, Contreras M, et al. Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc Natl Acad Sci USA.* 2010;107(26):11971-11975.
 58. Sevelsted A, Stokholm J, Bønnelykke K, Bisgaard H. Cesarean section and chronic immune disorders. *Pediatrics.* 2015;135(1):e92-e98.
 59. Cox LM, Yamanishi S, Sohn J, et al. Altering the intestinal microbiota during a critical developmental window has lasting metabolic consequences. *Cell.* 2014;158(4):705-721.
 60. Kuo C-H, Kuo H-F, Huang C-H, Yang S-N, Lee M-S, Hung C-H. Early life exposure to antibiotics and the risk of childhood allergic diseases: an update from the perspective of the hygiene hypothesis. *J Microbiol Immunol Infect.* 2013;46(5):320-329.
 61. Ma D, Leulier F, Storelli G, Mitchell M. Studying host-microbiota mutualism in *Drosophila*: harnessing the power of gnotobiotic flies. *Biomed J.* 2015;38(4):285.
 62. Milligan-Myhre K, Charette JR, Phennic RT, et al. Study of Host-Microbe Interactions in Zebrafish. In: *Methods in Cell Biology*, Vol. 105. Elsevier; 2011:87-116. Available at: <http://linkinghub.elsevier.com/retrieve/pii/B9780123813206000047>. Accessed September 12, 2017.
 63. Hörmansperger G, Schaubeck M, Haller D. Intestinal microbiota in animal models of inflammatory diseases. *ILAR J.* 2015;56(2):179-191.

64. Bauer H, Horowitz RE, Levenson SM, Popper H. The response of the lymphatic tissue to the microbial flora. Studies on germfree mice. *Am J Pathol.* 1963;42:471–483.
65. Stěpánková R, Kovářů F, Kruml J. Lymphatic tissue of the intestinal tract of germfree and conventional rabbits. *Folia Microbiol (Praha).* 1980;25(6):491–495.
66. Baron F. Kinetics of engraftment in patients with hematologic malignancies given allogeneic hematopoietic cell transplantation after nonmyeloablative conditioning. *Blood.* 2004;104(8):2254–2262.
67. Bartelink IH, Belitser SV, Knibbe CAJ, et al. Immune reconstitution kinetics as an early predictor for mortality using various hematopoietic stem cell sources in children. *Biol Blood Marrow Transplant.* 2013;19(2):305–313.
68. Geddes M, Storek J. Immune reconstitution following hematopoietic stem-cell transplantation. *Best Pract Res Clin Haematol.* 2007;20(2):329–348.
69. Gratama JW, Boeckh M, Nakamura R, et al. Immune monitoring with iTAg MHC Tetramers for prediction of recurrent or persistent cytomegalovirus infection or disease in allogeneic hematopoietic stem cell transplant recipients: a prospective multicenter study. *Blood.* 2010;116(10):1655–1662.
70. Kim DH, Sohn SK, Won DI, Lee NY, Suh JS, Lee KB. Rapid helper T cell recovery above 200 × 106/l at 3 months correlates to successful transplant outcomes after allogeneic stem cell transplantation. *Bone Marrow Transplant.* 2006;37(12):1119–1128.
71. Novitzky N, Davison GM, Hale G, Waldmann H. Immune reconstitution at 6 months following T-cell depleted hematopoietic stem cell transplantation is predictive for treatment outcome. *Transplantation.* 2002;74(11):1551–1559.
72. Porrata LF, Litzow MR, Inwards DJ, et al. Infused peripheral blood autograft absolute lymphocyte count correlates with day 15 absolute lymphocyte count and clinical outcome after autologous peripheral hematopoietic stem cell transplantation in non-Hodgkin's lymphoma. *Bone Marrow Transplant.* 2004;33(3):291–298.
73. Savani BN. Factors associated with early molecular remission after T cell-depleted allogeneic stem cell transplantation for chronic myelogenous leukemia. *Blood.* 2006;107(4):1688–1695.
74. Storek J, Espino G, Dawson MA, Storer B, Flowers ME, Maloney DG. Low B-cell and monocyte counts on day 80 are associated with high infection rates between days 100 and 365 after allogeneic marrow transplantation. *Blood.* 2000;96(9):3290–3293.
75. Jenq RR, Ubeda C, Taur Y, et al. Regulation of intestinal inflammation by microbiota following allogeneic bone marrow transplantation. *J Exp Med.* 2012;209(5):903–911.
76. Levy M, Kolodziejczyk AA, Thaiss CA, Elinav E. Dysbiosis and the immune system. *Nat Rev Immunol.* 2017;17(4):219–232.
77. Hooper LV, Littman DR, Macpherson AJ. Interactions between the microbiota and the immune system. *Science.* 2012;336(6086):1268–1273.
78. Josefsson KS, Baldrige MT, Kadmon CS, King KY. Antibiotics impair murine hematopoiesis by depleting the intestinal microbiota. *Blood.* 2017;129(6):729–739.
79. van den Brink MRM, Velardi E, Perales M-A. Immune reconstitution following stem cell transplantation. *Hematology.* 2015;2015(1):215–219.
80. Atarashi K, Tanoue T, Oshima K, et al. Treg induction by a rationally selected mixture of Clostridia strains from the human microbiota. *Nature.* 2013;500(7461):232–236.
81. Atarashi K, Tanoue T, Shima T, et al. Induction of colonic regulatory T cells by indigenous clostridium species. *Science.* 2011;331(6015):337–341.
82. Smith PM, Howitt MR, Panikov N, et al. The microbial metabolites, short-chain fatty acids, regulate colonic Treg cell homeostasis. *Science.* 2013;341(6145):569–573.
83. Heimesaat MM, Nogai A, Bereswill S, et al. MyD88/TLR9 mediated immunopathology and gut microbiota dynamics in a novel murine model of intestinal graft-versus-host disease. *Gut.* 2010;59(8):1079–1087.
84. Eriguchi Y, Takashima S, Oka H, et al. Graft-versus-host disease disrupts intestinal microbial ecology by inhibiting Paneth cell production of α -defensins. *Blood.* 2012;120(1):223–231.
85. Tawara I, Liu C, Tamaki H, et al. Influence of donor microbiota on the severity of experimental graft-versus-host-disease. *Biol Blood Marrow Transplant.* 2013;19(1):164–168.
86. Li Y, Chen H-L, Bannick N, et al. Intestinal helminths regulate lethal acute graft-versus-host disease and preserve the graft-versus-tumor effect in mice. *J Immunol.* 2015;194(3):1011–1020.
87. Shono Y, Docampo MD, Peled JU, et al. Increased GVHD-related mortality with broad-spectrum antibiotic use after allogeneic hematopoietic stem cell transplantation in human patients and mice. *Sci Transl Med.* 2016;8(339):339ra71.
88. Mathewson ND, Jenq R, Mathew AV, et al. Gut microbiome-derived metabolites modulate intestinal epithelial cell damage and mitigate graft-versus-host disease. *Nat Immunol.* 2016;17(5):505–513.
89. Simms-Waldrip TR, Sunkersett G, Coughlin LA, et al. Antibiotic-induced depletion of anti-inflammatory clostridia is associated with the development of graft-versus-host disease in pediatric stem cell transplantation patients. *Biol Blood Marrow Transplant.* 2017;23(5):820–829.
90. van der Velden WJFM, Netea MG, de Haan AFJ, Huls GA, Donnelly JP, Blijlevens NMA. Role of the mycobiome in human acute graft-versus-host disease. *Biol Blood Marrow Transplant.* 2013;19(2):329–332.
91. Biagi E, Zama D, Nastasi C, et al. Gut microbiota trajectory in pediatric patients undergoing hematopoietic SCT. *Bone Marrow Transplant.* 2015;50(7):992–998.
92. Bilinski J, Robak K, Peric Z, et al. Impact of gut colonization by antibiotic-resistant bacteria on the outcomes of allogeneic hematopoietic stem cell transplantation: a retrospective, single-center study. *Biol Blood Marrow Transplant.* 2016;22(6):1087–1093.
93. Routy B, Letendre C, Enot D, et al. The influence of gut-decontamination prophylactic antibiotics on acute graft-versus-host disease and survival following allogeneic hematopoietic stem cell transplantation. *Oncoimmunology.* 2017;6(1):e1258506.
94. Doki N, Suyama M, Sasajima S, et al. Clinical impact of pre-transplant gut microbial diversity on outcomes of allogeneic hematopoietic stem cell transplantation. *Ann Hematol.* 2017;96(9):1517–1523.
95. Liu C, Frank DN, Horch M, et al. Associations between acute gastrointestinal GvHD and the baseline gut microbiota of allogeneic hematopoietic stem cell transplant recipients and donors. *Bone Marrow Transplant.* 2017;52(12):1643–1650.
96. Zitvogel L, Ayyoub M, Routy B, Kroemer G. Microbiome and anticancer immunosurveillance. *Cell.* 2016;165(2):276–287.
97. Ayabe T, Satchell DP, Wilson CL, Parks WC, Selsted ME, Ouellette AJ. Secretion of microbicidal α -defensins by intestinal Paneth cells in response to bacteria. *Nat Immunol.* 2000;1(2):113–118.
98. Masuda K, Sakai N, Nakamura K, Yoshioka S, Ayabe T. Bactericidal activity of mouse α -defensin cryptdin-4 predominantly affects noncommensal bacteria. *J Innate Immun.* 2011;3(3):315–326.
99. Cash HL, Whitham CV, Behrendt CL, Hooper LV. Symbiotic bacteria direct expression of an intestinal bactericidal lectin. *Science.* 2006;313(5790):1126–1130.
100. Eriguchi Y, Nakamura K, Hashimoto D, et al. Decreased secretion of Paneth cell α -defensins in graft-versus-host disease. *Transpl Infect Dis.* 2015;17(5):702–706.
101. Weber D, Frauenschläger K, Ghimire S, et al. The association between acute graft-versus-host disease and antimicrobial peptide expression in the gastrointestinal tract after allogeneic stem cell transplantation. *PLoS ONE.* 2017;12(9):e0185265.
102. Buffie CG, Jarchum I, Equinda M, et al. Profound alterations of intestinal microbiota following a single dose of clindamycin results in sustained susceptibility to Clostridium difficile-induced colitis. *Infect Immun.* 2012;80(1):62–73.
103. Koh AY, Köhler JR, Coggshall KT, Van Rooijen N, Pier GB. Mucosal damage and neutropenia are required for Candida albicans dissemination. *PLoS Pathog.* 2008;4(2):e35.
104. Brandl K, Plitas G, Mihu CN, et al. Vancomycin-resistant enterococci exploit antibiotic-induced innate immune deficits. *Nature.* 2008;455(7214):804–807.
105. Bell G, Gouyon P-H. Arming the enemy: the evolution of resistance to self-proteins. *Microbiology (Reading, Engl.).* 2003;149(Pt 6):1367–1375.
106. Tanaka K, Sawamura S, Satoh T, Kobayashi K, Noda S. Role of the indigenous microbiota in maintaining the virus-specific CD8 memory T cells in the lung of mice infected with murine cytomegalovirus. *J Immunol.* 2007;178(8):5209–5216.
107. de Steenhuijsen Pijters WA, Heinonen S, Hasrat R, et al. Nasopharyngeal microbiota, host transcriptome and disease severity in children with respiratory syncytial virus infection. *Am J Respir Crit Care Med.* 2016;194(9):1104–1115.
108. Nakamura S, Davis KM, Weiser JN. Synergistic stimulation of type I interferons during influenza virus coinfection promotes Streptococcus pneumoniae colonization in mice. *J Clin Invest.* 2011;121(9):3657–3665.
109. Allander T, Tammi MT, Eriksson M, Bjerkner A, Tiveljung-Lindell A, Andersson B. Cloning of a human parvovirus by molecular screening of respiratory tract samples. *Proc Natl Acad Sci USA.* 2005;102(36):12891–12896.
110. Lim ES, Zhou Y, Zhao G, et al. Early life dynamics of the human gut virome and bacterial microbiome in infants. *Nat Med.* 2015;21(10):1228–1234.
111. Pfeiffer JK, Virgin HW. Viral immunity. Transkingdom control of viral infection and immunity in the mammalian intestine. *Science.* 2016;351(6270):doi:10.1126/science.aad5872.
112. Tarabichi Y, Li K, Hu S, et al. The administration of intranasal live attenuated influenza vaccine induces changes in the nasal microbiota and nasal epithelium gene expression profiles. *Microbiome.* 2015;3:74.
113. Seo S, Renaud C, Kuypers JM, et al. Idiopathic pneumonia syndrome after hematopoietic cell transplantation: evidence of occult infectious etiologies. *Blood.* 2015;125(24):3789–3797.
114. Herrera AF, Soriano G, Bellizzi AM, et al. Cord colitis syndrome in cord-blood stem-cell transplantation. *N Engl J Med.* 2011;365(9):815–824.

115. Bhatt AS, Freeman SS, Herrera AF, et al. Sequence-based discovery of *Bradyrhizobium enterica* in cord colitis syndrome. *N Engl J Med*. 2013;369(6):517–528.
116. Blaser MJ. The microbiome revolution. *J Clin Invest*. 2014;124(10):4162–4165.
117. Preisler HD, Goldstein IM, Henderson ES. Gastrointestinal “sterilization” in the treatment of patients with acute leukemia. *Cancer*. 1970;26(5):1076–1081.
118. Pizzo PA, Purvis DS, Waters C. Microbiological evaluation of food items. For patients undergoing gastrointestinal decontamination and protected isolation. *J Am Diet Assoc*. 1982;81(3):272–279.
119. Smith LH, Besser SG. Dietary restrictions for patients with neutropenia: a survey of institutional practices. *Oncol Nurs Forum*. 2000;27(3):515–520.
120. Lassiter M, Schneider SM. A pilot study comparing the neutropenic diet to a non-neutropenic diet in the allogeneic hematopoietic stem cell transplantation population. *Clin J Oncol Nurs*. 2015;19(3):273–278.
121. Trifilio S, Helenowski I, Giel M, et al. Questioning the role of a neutropenic diet following hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2012;18(9):1385–1390.
122. Moody K, Charlson ME, Finlay J. The neutropenic diet: what's the evidence? *J Pediatr Hematol Oncol*. 2002;24(9):717–721.
123. Guièze R, Lemal R, Cabrespine A, et al. Enteral versus parenteral nutritional support in allogeneic haematopoietic stem-cell transplantation. *Clin Nutr*. 2014;33(3):533–538.
124. Seguy D, Duhamel A, Rejeb MB, et al. Better outcome of patients undergoing enteral tube feeding after myeloablative conditioning for allogeneic stem cell transplantation. *Transplantation*. 2012;94(3):287–294.
125. Szeluga DJ, Stuart RK, Brookmeyer R, Utermohlen V, Santos GW. Nutritional support of bone marrow transplant recipients: a prospective, randomized clinical trial comparing total parenteral nutrition to an enteral feeding program. *Cancer Res*. 1987;47(12):3309–3316.
126. Lemal R, Cabrespine A, Pereira B, et al. Could enteral nutrition improve the outcome of patients with haematological malignancies undergoing allogeneic haematopoietic stem cell transplantation? A study protocol for a randomized controlled trial (the NEPHA study). *Trials*. 2015;16:136.
127. Gibson GR, Hutkins R, Sanders ME, et al. Expert consensus document: the International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nat Rev Gastroenterol Hepatol*. 2017;14(8):491–502.
128. Iyama S, Sato T, Tatsumi H, et al. Efficacy of enteral supplementation enriched with glutamine, fiber, and oligosaccharide on mucosal injury following hematopoietic stem cell transplantation. *Case Rep Oncol*. 2014;7(3):692–699.
129. Andermann TM, Rezvani A, Bhatt AS. Microbiota manipulation with prebiotics and probiotics in patients undergoing stem cell transplantation. *Curr Hematol Malig Rep*. 2016;11(1):19–28.
130. Lindsay JO, Whelan K, Stagg AJ, et al. Clinical, microbiological, and immunological effects of fructo-oligosaccharide in patients with Crohn's disease. *Gut*. 2006;55(3):348–355.
131. Casellas F, Borrueal N, Torrejón A, et al. Oral oligofructose-enriched inulin supplementation in acute ulcerative colitis is well tolerated and associated with lowered faecal calprotectin. *Aliment Pharmacol Ther*. 2007;25(9):1061–1067.
132. Margreiter M, Ludl K, Phleps W, Kaehler ST. Therapeutic value of a *Lactobacillus gasseri* and *Bifidobacterium longum* fixed bacterium combination in acute diarrhea: a randomized, double-blind, controlled clinical trial. *Int J Clin Pharmacol Ther*. 2006;44(5):207–215.
133. Kruis W, Fric P, Pokrotnieks J, et al. Maintaining remission of ulcerative colitis with the probiotic *Escherichia coli* Nissle 1917 is as effective as with standard mesalazine. *Gut*. 2004;53(11):1617–1623.
134. Vitetta L, Briskey D, Alford H, Hall S, Coulson S. Probiotics, prebiotics and the gastrointestinal tract in health and disease. *Inflammopharmacology*. 2014;22(3):135–154.
135. Preidis GA, Versalovic J. Targeting the human microbiome with antibiotics, probiotics, and prebiotics: gastroenterology enters the metagenomics era. *Gastroenterology*. 2009;136(6):2015–2031.
136. Gerbitz A, Schultz M, Wilke A, et al. Probiotic effects on experimental graft-versus-host disease: let them eat yogurt. *Blood*. 2004;103(11):4365–4367.
137. Tavil B, Koksali E, Yalcin SS, Uckan D. Pretransplant nutritional habits and clinical outcome in children undergoing hematopoietic stem cell transplant. *Exp Clin Transplant*. 2012;10(1):55–61.
138. Ladas EJ, Bhatia M, Chen L, et al. The safety and feasibility of probiotics in children and adolescents undergoing hematopoietic cell transplantation. *Bone Marrow Transplant*. 2016;51(2):262–266.
139. Gorshein E, Wei C, Ambrosy S, et al. *Lactobacillus rhamnosus* GG probiotic enteric regimen does not appreciably alter the gut microbiome or provide protection against GVHD after allogeneic hematopoietic stem cell transplantation. *Clin Transplant*. 2017;31(5):doi:10.1111/ctr.12947.
140. Salminen MK, Rautelin H, Tynkkynen S, et al. *Lactobacillus bacteremia*, clinical significance, and patient outcome, with special focus on probiotic *L. rhamnosus* GG. *Clin Infect Dis*. 2004;38(1):62–69.
141. Mehta A, Rangarajan S, Borate U. A cautionary tale for probiotic use in hematopoietic SCT patients—*Lactobacillus acidophilus* sepsis in a patient with mantle cell lymphoma undergoing hematopoietic SCT. *Bone Marrow Transplant*. 2013;48(3):461–462.
142. Gough E, Shaikh H, Manges AR. Systematic review of intestinal microbiota transplantation (fecal bacteriotherapy) for recurrent *Clostridium difficile* infection. *Clin Infect Dis*. 2011;53(10):994–1002.
143. Kassam Z, Lee CH, Yuan Y, Hunt RH. Fecal microbiota transplantation for *Clostridium difficile* infection: systematic review and meta-analysis. *Am J Gastroenterol*. 2013;108(4):500–508.
144. Neemann K, Eichele DD, Smith PW, Bociak R, Akhtari M, Freifeld A. Fecal microbiota transplantation for fulminant *Clostridium difficile* infection in an allogeneic stem cell transplant patient. *Transpl Infect Dis*. 2012;14(6):E161–E165.
145. de Castro CG, Ganc AJ, Ganc RL, Petrolli MS, Hamerschlack N. Fecal microbiota transplant after hematopoietic SCT: report of a successful case. *Bone Marrow Transplant*. 2015;50(1):145.
146. Mittal C, Miller N, Meighani A, Hart BR, John A, Ramesh M. Fecal microbiota transplant for recurrent *Clostridium difficile* infection after peripheral autologous stem cell transplant for diffuse large B-cell lymphoma. *Bone Marrow Transplant*. 2015;50(7):1010.
147. Webb BJ, Brunner A, Ford CD, Gazdik MA, Petersen FB, Hoda D. Fecal microbiota transplantation for recurrent *Clostridium difficile* infection in hematopoietic stem cell transplant recipients. *Transpl Infect Dis*. 2016;18(4):628–633.
148. Moss EL, Falconer SB, Tkachenko E, et al. Long-term taxonomic and functional divergence from donor bacterial strains following fecal microbiota transplantation in immunocompromised patients. *PLoS ONE*. 2017;12(8):doi:10.1371/journal.pone.0182585. e0182585.
149. Kakihana K, Fujioka Y, Suda W, et al. Fecal microbiota transplantation for patients with steroid-resistant acute graft-versus-host disease of the gut. *Blood*. 2016;128(16):2083–2088.
150. Spindelboeck W, Schulz E, Uhl B, et al. Repeated fecal microbiota transplantations attenuate diarrhea and lead to sustained changes in the fecal microbiota in acute, refractory gastrointestinal graft-versus-host-disease. *Haematologica*. 2017;102(5):e210–e213.
151. Bilinski J, Grzesiowski P, Sorensen N, et al. Fecal microbiota transplantation in patients with blood disorders inhibits gut colonization with antibiotic-resistant bacteria: results of a prospective, single-center study. *Clin Infect Dis*. 2017;65(3):364–370.
152. Quera R, Espinoza R, Estay C, Rivera D. Bacteremia as an adverse event of fecal microbiota transplantation in a patient with Crohn's disease and recurrent *Clostridium difficile* infection. *J Crohns Colitis*. 2014;8(3):252–253.
153. Schwartz M, Gluck M, Koon S. Norovirus gastroenteritis after fecal microbiota transplantation for treatment of *Clostridium difficile* infection despite asymptomatic donors and lack of sick contacts. *Am J Gastroenterol*. 2013;108(8):1367.
154. Di Bella S, Gouliouris T, Petrosillo N. Fecal microbiota transplantation (FMT) for *Clostridium difficile* infection: focus on immunocompromised patients. *J Infect Chemother*. 2015;21(4):230–237.
155. Smith M, Kassam Z, Edelstein C, Burgess J, Alm E. OpenBiome remains open to serve the medical community. *Nat Biotechnol*. 2014;32(9):867.
156. Song SJ, Lauber C, Costello EK, et al. Cohabiting family members share microbiota with one another and with their dogs. *eLife*. 2013;2:e00458.
157. Sinha R, Abu-Ali G, Vogtmann E, et al. Assessment of variation in microbial community amplicon sequencing by the Microbiome Quality Control (MBQC) project consortium. *Nat Biotechnol*. 2017;35(11):1077–1086.
158. Atkinson TM, Ryan SJ, Bennett AV, et al. The association between clinician-based common terminology criteria for adverse events (CTCAE) and patient-reported outcomes (PRO): a systematic review. *Support Care Cancer*. 2016;24(8):3669–3676.
159. Shaw BE, Lee SJ, Horowitz MM, Wood WA, Rizzo JD, Flynn KE. Can we agree on patient-reported outcome measures for assessing hematopoietic cell transplantation patients? A study from the CIBMTR and BMT CTN. *Bone Marrow Transplant*. 2016;51(9):1173–1179.
160. Mimeo M, Tucker AC, Voigt CA, Lu TK. Programming a human commensal bacterium, bacteroides thetaiotaomicron, to sense and respond to stimuli in the murine gut microbiota. *Cell Syst*. 2016;2(3):214.