

Review

Remission in psoriatic arthritis—where are we now?

Laura C. Coates^{1,a}, Philip G. Conaghan¹, Maria Antonietta D'Agostino^{2,3}, Maarten De Wit⁴, Oliver FitzGerald⁵, Tore K. Kvien⁶, Rik Lories⁷, Philip Mease^{8,9}, Peter Nash¹⁰, Georg Schett¹¹, Enrique R. Soriano¹² and Paul Emery¹

Abstract

Advances in treatments and treatment strategies for PsA have led to many patients responding well to management of their disease, and targeting remission as a treatment goal is now a possibility. Treat to target is a strategy aimed at maximizing benefit, irrespective of the type of medication used, by monitoring disease activity and using it to guide therapy. The measurement of response to treatment has been the subject of wide discussions among experts for some time, and many instruments exist. Comparisons of the different measures and their different strengths and weaknesses is ongoing. The impact of modern imaging techniques on monitoring disease progression is also evolving, and advanced techniques using both MRI and US have the potential to improve management of PsA through identification of risk factors for poor prognosis as well as accurate assessment of inflammation and damage, including subclinical disease. Increased understanding of the pathways that drive the pathogenesis of PsA will be key to identifying specific biomarkers for the disease and developing effective treatment strategies. Targets for response, considerations for use of a treat to target strategy in PsA, different imaging techniques and serological aspects of remission are all discussed in this review, and areas for further research are identified.

Key words: PsA, remission, treat to target, tight control, disease activity, imaging, biomarkers, Doppler ultrasound, MRI

Rheumatology key messages

- Recent advances in pharmacological treatment and treatment strategies for PsA have improved responses in many patients.
- Targets for response in PsA have been widely discussed, but little evidence is available to guide consensus.
- Advances in imaging techniques have potential to improve PsA management and provide outcomes for remission.

Introduction

The treatment goals for patients with PsA are control of disease activity, improvement of physical function and

quality of life and prevention of structural damage to joints [1–3]. In the last few years, advances in pharmacological treatment of PsA, particularly the introduction of

¹Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, and National Institute for Health Research (NIHR) Leeds Musculoskeletal Biomedical Research Unit, Leeds, UK, ²APHP, Hôpital Ambroise Paré, Rheumatology Department, Boulogne-Billancourt, 92100, ³INSERM U1173, Laboratoire d'Excellence INFLAMEX, UFR Simone Veil, Versailles-Saint-Quentin University, Saint-Quentin en Yvelines, 78180, France, ⁴Department of Medical Humanities, VU University Medical Centre, Amsterdam, The Netherlands, ⁵Department of Rheumatology, St Vincent's University Hospital, and Conway Institute for Biomolecular Research, University College Dublin, Ireland, ⁶Department of Rheumatology, Diakonhjemmet Hospital, Oslo, Norway, ⁷Skeletal Biology and Engineering Research Center, KU Leuven, Leuven, Belgium; Division of Rheumatology, UZ Leuven, Leuven, Belgium, ⁸University of Washington School of Medicine, ⁹Clinical Research Division, Swedish Medical Center, Seattle, WA, USA, ¹⁰Department of Medicine,

University of Queensland, Brisbane, Queensland, Australia, ¹¹Department of Internal Medicine 3 and Institute for Clinical Immunology, Friedrich Alexander University of Erlangen-Nuremberg, Erlangen, Germany and ¹²Rheumatology Unit, Internal Medical Services, Hospital Italiano de Buenos Aires, Buenos Aires, Argentina

Submitted 17 February 2017; revised version accepted 14 August 2017

^aPresent address: Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Oxford, UK.

Correspondence to: Laura Coates, Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, Botnar Research Centre, University of Oxford, Windmill Road, Oxford, OX3 7LD, UK.
E-mail: lauraccoates@gmail.com

biologic therapies, have enabled excellent responses to be achieved in many patients [4]. However, PsA is a heterogeneous disease and measuring its response to treatment, both in the clinic and in clinical trials, has been the subject of wide debate.

Advances in treatment strategies for rheumatic diseases have also occurred. Treat to target (T2T) is aimed at maximizing benefit, irrespective of the type of medication used, by monitoring disease activity using the best current measures and remission criteria [5]. The Tight Control of disease activity in RA [6] study showed that escalating therapy in a T2T strategy could improve outcomes in RA. The study investigated an intensive treatment strategy consisting of frequent, objective assessment of patients, intensive use of intra-articular steroid injections if needed, and a structured protocol for the escalation of treatment in patients with active disease despite treatment. The targets in RA have become more stringent over time, related to a greater ability to achieve remission as new, better treatments are developed [7].

Evidence for T2T in PsA only began to emerge in 2013, and many treatments and outcome measures have been borrowed from RA. There has been little agreement on what target(s) for response should be used in PsA [8], and a literature review by the EULAR showed that there were few relevant studies on T2T in PsA [8].

The use of MRI, US and CT in the study of PsA has permitted a better understanding of the various pathologies of the PsA phenotypes. These sensitive imaging techniques have highlighted the high frequency of subclinical inflammation and added insights into the persistence of inflammation and structural damage after therapy [9–12].

This review provides an overview of the current status of targeting remission in PsA, including a focus on areas that need more research. It resulted from a consensus meeting with an expert panel of clinicians involved in PsA routine management and research in February 2016.

Considerations in applying T2T in PsA clinical practice

Specific aspects and challenges of remission in PsA

In order to assess remission, it must first be defined. Remission implies that at a minimum, the inflammatory disease process will be controlled such that the patient has no symptoms and no long-term functional or structural joint consequences [13]. Even in clinical manifestations, PsA is a multi-faceted disease with varied rheumatological and dermatological presentations. Beyond this, PsA not only has clinical manifestations, but is also characterized by structural and immunological changes. Therefore PsA remission may encompass more than remission of the clinical signs and symptoms of musculoskeletal and skin disease.

Core domains for assessment of PsA were defined by OMERACT in 2006 [14] and updated in 2016 [15, 16], and a core set of domain criteria for minimal disease activity (MDA) in PsA have also been defined [7, 17]. Ideally, the

target for remission should be feasible for clinical use and, as PsA is a heterogeneous condition, should include assessment of all key different domains. As yet, there are no reliable serum markers of PsA disease activity.

Another major factor affecting quality of life for PsA patients is comorbidity, and this aspect needs to be considered when setting realistic expectations of disease remission. A large proportion of PsA patients have comorbidities, which are often under-recognized and under-treated, which may influence treatment, prognosis and outcomes; they include cardiovascular disease, obesity, metabolic syndrome, depression, uveitis and cancer [18, 19]. One study has found that 42% of PsA patients have three or more comorbidities; however, the incremental effects of comorbidities on quality of life relate more to the type rather than the number of comorbidities [20]. Targeted treatment is therefore an important concept in achieving patient-defined remission.

Patient perspectives on disease activity, treatment and remission in PsA

Patients with PsA and their physicians may view the disease differently, and there is a discrepancy between patient and physician assessment of joint activity [21]. An analysis of 565 patients found that patients scored their disease worse than physicians, with the discordance greater for joints than for skin parameters. Similar discrepancy is well documented in RA [22], but has been less well studied in PsA.

Patient education in PsA is often not optimal and PsA patients are less empowered than those with RA [23]. However, a recent study showed that the difference between patients' and physicians' global assessment of disease activity as well as the difference between tender and swollen joint count were associated with a reduced risk of achieving remission, both in PsA and RA [24].

Results of the Multinational Assessment of Psoriasis and Psoriatic Arthritis Survey [25] showed that 59% of surveyed PsA patients were receiving no treatment or only topical treatment. This is partly due to low expectations on the part of patients that dermatologists or rheumatologists will be able to offer effective treatments.

For patients, the impact of disease on quality of life and function is important. Although individuals may have very different expectations of how their disease is managed, aspects of disease and treatment that are important to patients are not adequately covered by the self-report measures (both patient reported outcomes and existing composite scores) most often used in PsA patients [26]. These include the impact of environmental factors, societal attitudes towards individuals with psoriasis (PsO) or PsA, the increased feeling of isolation from social activities, and treatment burden, resulting in, for instance, lack of leisure time. Expectations are an important factor in disease management. For example, there is evidence that RA patients consider remission more as a feeling of returning to normality, rather than an absence or reduction of symptoms [27]. Treatment clearly impacts quality of life for PsA patients, and there is evidence that treatment

early in the course of the disease (<2 years' disease duration) led to greater improvements in arthritis scores and quality of life measures compared with those treated after having the disease for >2 years [28].

Clinical remission in PsA: how to measure it

Comparison of different instruments used to assess disease activity and measure outcomes

The composite measures used to assess disease activity are compared in Table 1, built by the authors' consensus while writing the paper. These composite measures combine individual measures of disease activity into a single score and, while this may be a more efficient approach than comparing across individual scores, the ability to distinguish between changes in disease activity in individual clinical features may be lost [29]. Different outcome measures may be used in clinical practice from those used in clinical trials although, if being used to guide treatment decisions in a trial, these measures must be feasible. A joint count assessing 68 joints for tenderness and 66 joints for swelling is employed in virtually all randomized controlled trials to constitute the primary outcome measure and is endorsed by OMERACT [14]. However, the Disease Activity Score 28 (DAS28) developed for RA [30] is an often-used measure of disease activity and remission in PsA in clinical practice around the world and is a secondary measure used in randomized controlled trials. However, it was not developed for PsA patients, it is purely a measure of joint inflammation and confined to 28 joints and it does not assess disease in common domains of PsA involvement, that is, DIP joints, feet or ankles, skin and nails.

Work is ongoing to compare different measures. The GRAppa Composite Exercise project aimed to develop new composite measures in PsA and compare them with existing indices [31]. The new indices included the psoriatic arthritis DAS and the Group for Research in Psoriasis and Psoriatic Arthritis (GRAPPA) composite index, which uses the arithmetic mean of desirability functions. These have been compared with existing indices such the Composite Psoriatic arthritis Disease Activity Index, Disease Activity index for Psoriatic Arthritis and DAS28.

A recent study in patients with active PsA demonstrated that different remission criteria provide different results [32], while the performance of six composite activity indices was compared in a real-world study [33]; all six showed good discriminant capacity, but the proportions of patients classified in the disease activity levels differed and, in particular, the rate of patients in remission was clearly different among the indices. Of note, none of the existing composite measures, including MDA, capture the original (2006) [14] nor the updated (2016) [15, 16] PsA core set.

Is T2T applicable in PsA?

The preferred target (state) of a T2T approach is remission or inactive disease as the primary goal and low disease activity or MDA as the secondary goal. The Tight Control of Psoriatic Arthritis (TICOPA) study [34] has recently shown that treating to target by escalating therapy with

a greater use of combination DMARDs and biologics in the tight control arm of the study significantly improves joint outcomes for newly diagnosed patients (Fig. 1) [34]. In the standard care arm patients were reviewed every 12 weeks in a general rheumatology outpatient clinic supervised by a consultant rheumatologist. No formal measures of disease activity were used to guide treatment decisions and there was no restriction on prescribing. In contrast, in the tight control arm patients were seen every 4 weeks by the study physician and treated according to a predefined treatment protocol. At each visit, patients were assessed for MDA criteria. Those not achieving MDA had their treatment escalated to the maximum dose according to the protocol. Patients achieving the MDA criteria continued on their current therapy.

Patients who received tight control treatment did experience more treatment-related adverse and serious adverse events than those receiving standard care, reporting more colds, nausea, fatigue and gastrointestinal upsets than those in the control arm (only partly explained by more frequent visits and recording of adverse events). However, despite larger doses of methotrexate in the tight control arm, liver enzyme abnormalities were similar in both arms. Patients in the tight control arm also required 27% more TNF inhibitor (TNFi) usage compared with those on standard care.

Patients in the TICOPA study were selected for early disease, and current T2T concepts may be more appropriate for newly diagnosed patients, and may be more difficult to apply in patients with longer disease duration with relatively more damage. This damage may affect the optimal primary target as patients with longstanding disease may be unable to meet these stringent criteria.

TICOPA is the first trial of strategy in PsA and further strategy trials are needed to weigh effectiveness against safety, since adverse events were also higher in the tight control arm of the TICOPA study compared with the standard care arm.

In two additional studies, a delay in diagnosis and intervention by 6 months demonstrated an impact on structural damage and long-term functional outcomes [35–37]. Data from the Swedish Early Psoriatic Arthritis Register [38] suggest that a shorter time between onset of symptoms and diagnosis is associated with better clinical outcomes at 5 years. It therefore appears that, as is the case with RA, early intervention combined with a tight control strategy is important to prevent irreversible damage.

Insights from modern imaging


Much less information is available on the use of US and MRI in PsA compared with RA, and imaging outcomes for remission in PsA are still evolving. What is clear is that MRI and US have the potential to improve PsA management [39]. Both techniques offer capability for assessing both inflammation and damage, with MRI enabling visualization of the spine in axial disease. Both may evaluate peripheral joints, with US being more patient friendly while providing multiple joint examinations in real time, though it is unable to visualize intra-bone pathology (osteitis). MRI can

TABLE 1 Comparison of features of clinical remission instruments

Features of instrument	MDA [7, 29, 55]	PASDAS [31]	CPDAI [78, 79]	DAPSA [80, 81]	RAPID3 [82]	P-sAID ^a [83]	DAS28 [30]
Developed for PSA?	Yes	Yes	Yes	Yes	No, generic	Yes	No, RA
Approx. time for patient to perform, min	2-5	2-5	5-10	1-2	2-5	2-5	1-2
Approx. time for assessor to perform, min	5-10	5-10	5-10	5-10	1	1	3-5
Complex calculation required	No	Yes	No	No	No	No	Yes
Continuous measure of disease activity	No	Yes	Yes	Yes	Yes	Impact not activity ^a	Yes
Measures peripheral arthritis	Yes	Yes	Yes	Yes	No	Impact ^a	Yes
Measures enthesitis	Yes	Yes	Yes	No	No	Impact ^a	No
Measures skin disease	Yes	Within global only	Yes	No	No	Impact ^a	No
Sensitive to change in PSA	Yes	Yes	Yes	Yes	Yes	Yes	Polyarticular only
Additional comments		Requires SF-36 and CRP	Development not evidence based	Requires CRP cutoffs based on physician opinion only, peripheral arthritis only	No physician exam	Impact measure rather than activity	Peripheral arthritis only and measures only 28 joints

^aP-sAID measures the impact of the disease on the patient rather than disease activity but identifies impact of the disease in many domains including MSK and skin. CPDAI, Composite Psoriatic arthritis Disease Activity Index; DAPSA, Disease Activity Index for Psoriatic Arthritis; DAS28, Disease Activity Score 28; MDA, minimal disease activity; PASDAS, Psoriatic Arthritis Disease Activity Score; P-sAID, Psoriatic Arthritis Impact of Disease; RAPID3, Routine Assessment of Patient Index Data.

Fig. 1 Proportion of patients achieving an ACR response at 48 weeks in tight control of psoriatic arthritis



Tight control vs standard of care. American College of Rheumatology Response criteria ACR20, 50 and 70. Data taken from [34].

evaluate only one joint or a joint area during one session, and may be less acceptable to patients due to the enclosed nature of the technique.

Although there are no typical US patterns characterizing PsA synovitis, with the exception of possibly more intense intra-articular vascularization seen in inflamed tissue, US has demonstrated good accuracy in assessing synovitis in PsA [10, 39–42]. In addition, the presence of US-detected synovitis has been shown to be associated with long-term radiographic erosion progression and poor outcomes [23]. Recently, Ficjan *et al.* [11] in a prospective and longitudinal study, developed an US composite score for the assessment of inflammatory and structural lesions in PsA, which demonstrates good metric properties including good sensitivity to change. US has also been shown to be of added value in assessing enthesitis and dactylitis. US can also be used for visualizing structural changes and inflammatory activity at the psoriatic skin and nail level; thickening of both the epidermis and the dermis is the most constant US pathological finding in psoriatic plaques, whereas the hypoechoic band in the upper dermis is associated with power Doppler (PD) activity (an expression of neoangiogenesis) and is particularly detectable in the active stages of the disease [43, 44].

Recommendations on imaging in spondyloarthritis (SpA) have been proposed by EULAR, including use of X-rays, US or MRI [45]. In axial SpA, the recommendation is for disease activity to be monitored with MRI of the sacroiliac joints and/or the spine, whereas conventional radiography should be used for long-term monitoring of structural damage. Similarly, for peripheral SpA, the recommendation is for US and MRI to be considered when monitoring disease activity (particularly synovitis and enthesitis), and conventional radiography is recommended to monitor structural damage.

The EULAR recommendations reflect the benefits of advanced imaging in assessing inflammation rather than assessing damage on X-rays, which has previously been an issue for trials conducted over short periods of time and trials that are not placebo controlled, where there is little

radiographic structural progression. The recommendations and recent evidence from clinical trials suggest that the field could be moving towards a time when X-rays are of limited value for imaging in SpA clinical studies.

Using imaging to monitor disease activity


Multiple studies have shown that MRI and US can detect inflammatory and structural lesions [46] and identify risk factors for poor prognosis in PsA [39, 47]. In terms of quantifying change, most US composite scores have been developed for the assessment of inflammatory and structural lesions in PsA (in terms of quantifying change), and they have demonstrated construct validity, sensitivity to change, reliability and feasibility [11]. The OMERACT PsA MRI score has similarly demonstrated good performance metrics [11, 48]. Several studies have now demonstrated the use of imaging to monitor disease activity and therapeutic response. A study of more than 300 SpA patients being treated with TNFi showed that PD US is a reliable method to monitor therapeutic response by measuring enthesitis [49], while US had a pivotal role in differential diagnosis and treatment monitoring in a patient with early PsA undergoing an aggressive tight control treatment programme and being monitored by US [50]. Similarly MRI has demonstrated responsiveness in PsA clinical studies [51].

Imaging of subclinical disease and remission

In line with the concept of subclinical disease first described in RA (inflammation detected by modern imaging but not examination), studies have found discrepancies between modern imaging and clinical findings, uncovering issues with accurate detection and clinical assessment of inflammation [9] and enthesitis, tenosynovitis or perisynovitis (i.e. extracapsular inflammation) in PsA patients in clinical remission [52]. In a study of newly diagnosed PsA comparing clinical examination with US in 49 patients, three-quarters were found to have sub-clinical synovitis, most frequently in the wrist and knee (Fig. 2) [53]. In patients on treatment, subclinical synovitis has been detected using US in patients classified as being in remission (as defined by MDA or DAS28) [9, 52]. There is some evidence that US detected synovitis might predict short-term flares in PsA patients in remission. However, it is not clear how important these US-detected manifestations really are and whether a T2T approach based on imaging would be superior to one based on clinical assessments. Some studies have shown that US can detect inflammatory and structural lesions and identify risk factors for poor prognosis in PsA [47]. Most of the studies have found discrepancies between US and clinical findings, uncovering issues with accurate detection and assessment of inflammation [9] and enthesitis, tenosynovitis or perisynovitis in PsA patients in clinical remission [52].

Enthesitis is another key, but often underestimated, feature of PsA, and therefore assessment of enthesitis with imaging is important, particularly as clinical measurements are often unreliable. Enthesitis may be predictive of flares, can predict clinical outcome, and can be

Fig. 2 Subclinical synovitis in 49 patients with early PsA



(A) US positive, clinical exam negative; (B) US negative, clinical exam positive. Reproduced from Freeston *et al.* Is there subclinical synovitis in early psoriatic arthritis? A clinical comparison with gray-scale and power Doppler ultrasound. *Arthritis Care Res* 2014;**66**:432–9 [53]. Copyright 2014. With permission from John Wiley & Sons.

present, although at a lower level, in remission or low disease activity states [54, 55]. A number of studies have been published supporting the validity of US in the assessment of entheses [56–60]. A recent study in newly diagnosed PsA found that three-quarters had sub-clinical synovitis, most frequently in the wrist and knee (Fig. 2).

Using contrast-enhanced US to detect persistent joint inflammation among patients in clinical remission showed that this technique is sufficiently sensitive to identify the presence of synovitis and thereby monitor remission [61]. Although there are limited data, there is some evidence that US detected synovitis might predict short-term flares in PsA patients in remission [62]. However, while PD US assessment may have an important role in monitoring treatment, its use at every clinic visit may not be feasible due to the expertise required, time and financial constraints [63, 64]. Significant training is required to obtain and interpret US images and such expertise may not be available at every centre. Further developments such as whole-body MRI could provide additional tools for use in clinical decision making, allowing the assessment of disease activity in axial and peripheral sites, and improving the detection of inflammatory changes in PsA in locations that are difficult to assess clinically [65]. Again, feasibility is an important consideration given the equipment required and the costs associated with scanning.

Serological and immunological aspects of remission in PsA

Two hypotheses have been formulated for the pathogenesis of PsA: firstly, that PsA is a classic autoimmune disease, or alternatively, that it begins with microtrauma at the enthesis, which then initiates innate immune events [66]. A better understanding of the key pathological pathways that drive progression from skin to bone involvement is needed in order to develop more effective treatment strategies.

Several studies on the origins of PsA have revealed signs of subclinical synovitis and enthesitis by MRI and US examination in the joints of patients who have PsO but not PsA [67–70], although the significance of these findings is not clear. Enthesitis has also been documented in healthy controls [59], and in patients with PsO without arthritis [71]. PsO patients also have a greater risk of developing enthesophytes than healthy controls [72]. There is also evidence to suggest that skin–bone interactions are triggered by IL-17, and IL-17 overexpression in mice with chronic skin inflammation induces bone loss through inhibition of osteoblast-mediated bone formation [73]. Finally, recent data show that BMI may also have an effect on the development of enthesitis, with overweight patients having less chance of fulfilling MDA criteria for tender enthesal points [74].

There may be differences in the pathologies of the various phenotypes of PsA, in terms of presence of certain cytokines/immune cells in synovitis and enthesitis; for example, T cell concentration changes or abnormalities in early disease may be predictive of progression and/or response to therapy [75]. Genetic factors, such as IL-23R

polymorphisms, may also predispose to exaggerated cytokine production and a hyperproliferative response, which can combine with mechanical stress factors into clinically apparent skin disease and clinically unapparent enthesal proliferation [76].

Biochemical markers of inflammation

The concept of immunological remission in PsA is only beginning to be understood. Standard biomarkers of inflammation are not particularly helpful in judging inflammatory disease activity in PsA. Unlike the situation in RA, there are few established biomarkers for immunological pathology in PsA. As the IL-17 pathway is integral in PsO and psoriatic disease [77], the IL-17–IL-23 pathway may provide more reliable markers for PsA in future, and recently changes in CD3⁺ T cell expression in PsA synovium have been shown to correlate with clinical response to treatment [75]. Biomarkers are under review as part of the OMERACT/GRAPPA initiative and several new biomarkers for PsA have been proposed, including calprotectin, serum amyloid A and myeloid-related protein, although none has been extensively validated to date. In the future, newer approaches such as proteomics may reveal better biomarkers of disease activity for PsA.

Conclusions

Advances in the treatment of PsA, particularly the introduction of biologic therapies, have allowed the disease to be controlled in many patients; however, measuring response to treatment in PsA patients is widely debated, partly caused by the heterogeneity of the disease. Changes have also occurred in treatment strategies for rheumatic diseases and the development of treat to target approaches have led to a change in the established treatment paradigm.

Both MRI and US techniques have the potential to improve PsA management, and imaging outcomes for remission in PsA are still evolving. The concept of immunological remission in PsA is only just beginning to be discussed and biomarkers for the disease are yet to be fully identified.

While remission is the ultimate goal for PsA patients and their physicians, questions on what exactly we should aim to achieve still remain; this review has examined the current status of targeting remission in PsA, with a focus on areas that need more research.

Acknowledgements

The authors thank Tina Patrick, PhD, of Novartis Ireland Ltd for providing medical writing support with the drafting of this manuscript, which was funded by Novartis in accordance with Good Publication Practice (GPP3) guidelines (<http://www.ismpp.org/gpp3>).

Funding: P.G.C. and P.E. are supported in part by the National Institute for Health Research Leeds Musculoskeletal Biomedical Research Unit. The views expressed are those of the author(s) and not necessarily

those of the National Health Service, the National Institute for Health Research or the Department of Health.

Disclosure statement: L.C.C. has received research funding and/or honoraria from AbbVie, Amgen, Bristol Myers Squibb (BMS), Celgene, Janssen, Lilly, Merck Sharp Dohme (MSD), Novartis, Pfizer, Sun Pharma and union chimique belge (UCB). P.G.C. has been on consultancies or speakers bureau for Abbvie, BMS, Eli Lilly, Novartis, Pfizer, Roche. M.d.W. has received consulting fees through Stitching Tools from AbbVie, BMS, Celgene, Eli Lilly, Novartis and Roche. P.N. has consulted for AbbVie, Amgen, BMS, Celgene, Lilly, Pfizer, Janssen, UCB, Novartis, Roche and Sanofi, been on the speaker bureau of AbbVie, Amgen, BMS, Celgene, Lilly, Pfizer, Janssen, UCB, Novartis, Roche and Sanofi and received research funding from AbbVie, Amgen, BMS, Celgene, Lilly, Pfizer, Janssen, UCB, Novartis, Roche and Sanofi. E.R.S. has received fees for speaking and/or consulting from AbbVie, BMS, Janssen, Novartis, Pfizer, Roche and UCB and received research funding to Hospital Italiano de Buenos Aires from AbbVie, BMS, Janssen, Novartis, Pfizer, Roche and UCB. T.K.K. has received fees for speaking and/or consulting from AbbVie, Biogen, BMS, Boehringer Ingelheim, Celgene, Celltrion, Eli Lilly, Epirus, Hospira, Merck-Serono, MSD, Mundipharma, Novartis, Oktal, Orion Pharma, Hospira/Pfizer, Roche, Sandoz and UCB and received research funding to Diakonhjemmet Hospital from AbbVie, BMS, MSD, Pfizer, Roche and UCB. P.E. has undertaken clinica trials and provided expert advice to Pfizer, MSD, UCB, Roche, Novartis, Samsung, Sandoz and Lilly. P.M. has received fees for research, consulting and speaking from AbbVie, Amgen, BMS, Celgene, Janssen, Novartis, Pfizer and UCB; received speaking fees from Genentech; and received research fees and consulting fees from Lilly and Sun. All other authors have declared no conflicts of interest.

References

- Coates LC, Kavanaugh A, Mease PJ *et al.* Group for Research and Assessment of Psoriasis and Psoriatic Arthritis 2015 treatment recommendations for psoriatic arthritis. *Arthritis Rheumatol* 2016;68:1060–71.
- Gossec L, Smolen JS, Ramiro S *et al.* European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2015 update. *Ann Rheum Dis* 2016;75:499–510.
- Smolen JS, Braun J, Dougados M *et al.* Treating spondyloarthritis, including ankylosing spondylitis and psoriatic arthritis, to target: recommendations of an international task force. *Ann Rheum Dis* 2014;73:6–16.
- Gottlieb A, Korman NJ, Gordon KB *et al.* Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 2. Psoriatic arthritis: overview and guidelines of care for treatment with an emphasis on the biologics. *J Am Acad Dermatol* 2008;58:851–64.
- Smolen JS. Treat-to-target as an approach in inflammatory arthritis. *Curr Opin Rheumatol* 2016;28:297–302.
- Grigor C, Capell H, Stirling A *et al.* Effect of a treatment strategy of tight control for rheumatoid arthritis (the TICORA study): a single-blind randomised controlled trial. *Lancet* 2004;364:263–9.
- Coates LC, Fransen J, Helliwell PS. Defining minimal disease activity in psoriatic arthritis: a proposed objective target for treatment. *Ann Rheum Dis* 2010;69:48–53.
- Schoels MM, Braun J, Dougados M *et al.* Treating axial and peripheral spondyloarthritis, including psoriatic arthritis, to target: results of a systematic literature search to support an international treat-to-target recommendation in spondyloarthritis. *Ann Rheum Dis* 2014;73:238–42.
- Michelsen B, Diamantopoulos AP, Hammer HB *et al.* Ultrasonographic evaluation in psoriatic arthritis is of major importance in evaluating disease activity. *Ann Rheum Dis* 2016;75:2108–13.
- Wiell C, Szkudlarek M, Hasselquist M *et al.* Ultrasonography, magnetic resonance imaging, radiography, and clinical assessment of inflammatory and destructive changes in fingers and toes of patients with psoriatic arthritis. *Arthritis Res Therapy* 2007;9:R119.
- Ficjan A, Husic R, Gretler J *et al.* Ultrasound composite scores for the assessment of inflammatory and structural pathologies in Psoriatic Arthritis (PsASon-Score). *Arthritis Res Ther* 2014;16:476.
- Curran SA, FitzGerald OM, Costello PJ *et al.* Nucleotide sequencing of psoriatic arthritis tissue before and during methotrexate administration reveals a complex inflammatory T cell infiltrate with very few clones exhibiting features that suggest they drive the inflammatory process by recognizing autoantigens. *J Immunol* 2004;172:1935–44.
- Kavanaugh A, Fransen J. Defining remission in psoriatic arthritis. *Clin Exp Rheumatol* 2006;24:S-83-7.
- Gladman DD, Mease PJ, Strand V *et al.* Consensus on a core set of domains for psoriatic arthritis. *J Rheumatol* 2007;34:1167–70.
- Orbai AM, Mease PJ, de Wit M *et al.* Report of the GRAPPA-OMERACT Psoriatic Arthritis Working Group from the GRAPPA 2015 Annual Meeting. *J Rheumatol* 2016;43:965–9.
- Orbai AM, de Wit M, Mease P *et al.* International patient and physician consensus on a psoriatic arthritis core outcome set for clinical trials. *Ann Rheum Dis* 2017;76:673–80.
- Coates LC, Helliwell PS. Defining low disease activity states in psoriatic arthritis using novel composite disease instruments. *J Rheumatol* 2016;43:371–5.
- Haroon M, FitzGerald O. Psoriatic arthritis: complexities, comorbidities and implications for the clinic. *Expert Rev Clin Immunol* 2016;12:405–16.
- Haroon M, Gallagher P, Heffernan E, FitzGerald O. High prevalence of metabolic syndrome and of insulin resistance in psoriatic arthritis is associated with the severity of underlying disease. *J Rheumatol* 2014;41:1357–65.
- Husted JA, Thavaneswaran A, Chandran V, Gladman DD. Incremental effects of comorbidity on quality of life in patients with psoriatic arthritis. *J Rheumatol* 2013;40:1349–56.

- 21 Eder L, Thavaneswaran A, Chandran V, Cook R, Gladman DD. Factors explaining the discrepancy between physician and patient global assessment of joint and skin disease activity in psoriatic arthritis patients. *Arthritis Care Res* 2015;67:264–72.
- 22 Barton JL, Imboden J, Graf J *et al*. Patient-physician discordance in assessments of global disease severity in rheumatoid arthritis. *Arthritis Care Res* 2010;62:857–64.
- 23 Helliwell P, Coates L, Chandran V *et al*. Qualifying unmet needs and improving standards of care in psoriatic arthritis. *Arthritis Care Res* 2014;66:1759–66.
- 24 Michelsen B, Kristianslund EK, Hammer HB *et al*. Discordance between tender and swollen joint count as well as patient's and evaluator's global assessment may reduce likelihood of remission in patients with rheumatoid arthritis and psoriatic arthritis: data from the prospective multicentre NOR-DMARD study. *Ann Rheum Dis* 2017;76:708–11.
- 25 Lebwohl MG, Bachelez H, Barker J *et al*. Patient perspectives in the management of psoriasis: results from the population-based Multinational Assessment of Psoriasis and Psoriatic Arthritis Survey. *J Am Acad Dermatol* 2014;70:871–81.e1–30.
- 26 Stamm TA, Nell V, Mathis M *et al*. Concepts important to patients with psoriatic arthritis are not adequately covered by standard measures of functioning. *Arthritis Rheum* 2007;57:487–94.
- 27 van Tuyl LH, Hewlett S, Sadlonova M *et al*. The patient perspective on remission in rheumatoid arthritis: 'You've got limits, but you're back to being you again'. *Ann Rheum Dis* 2015;74:1004–10.
- 28 Kirkham B, de Vlam K, Li W *et al*. Early treatment of psoriatic arthritis is associated with improved patient-reported outcomes: findings from the etanercept PRESTA trial. *Clin Exp Rheumatol* 2015;33:11–9.
- 29 Soriano ER. Defining remission in psoriatic arthritis: are we getting closer? *J Rheumatol* 2015;42:907–8.
- 30 van Gestel AM, Haagsma CJ, van Riel PL. Validation of rheumatoid arthritis improvement criteria that include simplified joint counts. *Arthritis Rheum* 1998;41:1845–50.
- 31 Helliwell PS, FitzGerald O, Fransen J *et al*. The development of candidate composite disease activity and responder indices for psoriatic arthritis (GRACE project). *Ann Rheum Dis* 2013;72:986–91.
- 32 Van den Bosch F, Kavanaugh A, Kron M, Kupper H, Mease PJ. Clinical remission in patients with active psoriatic arthritis treated with adalimumab and correlations in joint and skin manifestations. *J Rheumatol* 2015;42:952–9.
- 33 Salaffi F, Ciapetti A, Carotti M, Gasparini S, Gutierrez M. Disease activity in psoriatic arthritis: comparison of the discriminative capacity and construct validity of six composite indices in a real world. *Biomed Res Int* 2014;2014:528105.
- 34 Coates LC, Moverley AR, McParland L *et al*. Effect of tight control of inflammation in early psoriatic arthritis (TICOPA): a UK multicentre, open-label, randomised controlled trial. *Lancet* 2015;386:2489–98.
- 35 Haroon M, Gallagher P, FitzGerald O. Diagnostic delay of more than 6 months contributes to poor radiographic and functional outcome in psoriatic arthritis. *Ann Rheum Dis* 2015;74:1045–50.
- 36 Tillett W, Jadon D, Shaddick G *et al*. Smoking and delay to diagnosis are associated with poorer functional outcome in psoriatic arthritis. *Ann Rheum Dis* 2013;72:1358–61.
- 37 Gladman DD, Thavaneswaran A, Chandran V, Cook RJ. Do patients with psoriatic arthritis who present early fare better than those presenting later in the disease?. *Ann Rheum Dis* 2011;70(12):2152–4.
- 38 Theander E, Husmark T, Alenius GM *et al*. Early psoriatic arthritis: short symptom duration, male gender and preserved physical functioning at presentation predict favourable outcome at 5-year follow-up. Results from the Swedish Early Psoriatic Arthritis Register (SwePsA). *Ann Rheum Dis* 2014;73:407–13.
- 39 Coates LC, Hodgson R, Conaghan PG, Freeston JE. MRI and ultrasonography for diagnosis and monitoring of psoriatic arthritis. *Best Pract Res Clin Rheumatol* 2012;26:805–22.
- 40 Gutierrez M, Filippucci E, De Angelis R *et al*. A sonographic spectrum of psoriatic arthritis: "the five targets". *Clin Rheumatol* 2010;29:133–42.
- 41 De Simone C, Caldarola G, D'Agostino M *et al*. Usefulness of ultrasound imaging in detecting psoriatic arthritis of fingers and toes in patients with psoriasis. *Clin Dev Immunol* 2011;2011:390726.
- 42 Sankowski AJ, Lebkowska UM, Cwikla J, Walecka I, Walecki J. The comparison of efficacy of different imaging techniques (conventional radiography, ultrasonography, magnetic resonance) in assessment of wrist joints and metacarpophalangeal joints in patients with psoriatic arthritis. *Pol J Radiol* 2013;78:18–29.
- 43 Gutierrez M, De Angelis R, Bernardini ML *et al*. Clinical, power Doppler sonography and histological assessment of the psoriatic plaque: short-term monitoring in patients treated with etanercept. *Br J Dermatol* 2011;164:33–7.
- 44 Gutierrez M, Wortsman X, Filippucci E *et al*. High-frequency sonography in the evaluation of psoriasis: nail and skin involvement. *J Ultrasound Med* 2009;28:1569–74.
- 45 Mandl P, Navarro-Compan V, Terslev L *et al*. EULAR recommendations for the use of imaging in the diagnosis and management of spondyloarthritis in clinical practice. *Ann Rheum Dis* 2015;74:1327–39.
- 46 Kaeley GS. Review of the use of ultrasound for the diagnosis and monitoring of enthesitis in psoriatic arthritis. *Curr Rheumatol Reports* 2011;13:338–45.
- 47 El Miedany Y, El Gaafary M, Youssef S, Ahmed I, Nasr A. Tailored approach to early psoriatic arthritis patients: clinical and ultrasonographic predictors for structural joint damage. *Clin Rheumatol* 2015;34:307–13.
- 48 Boyesen P, McQueen FM, Gandjbakhch F *et al*. The OMERACT Psoriatic Arthritis Magnetic Resonance Imaging Score (PsAMRIS) is reliable and sensitive to change: results from an OMERACT workshop. *J Rheumatol* 2011;38:2034–8.
- 49 Naredo E, Battle-Gualda E, Garcia-Vivar ML *et al*. Power Doppler ultrasonography assessment of entheses in

- spondyloarthropathies: response to therapy of enthesal abnormalities. *J Rheumatol* 2010;37:2110-7.
- 50 Bandinelli F, Bonacci E, Matucci-Cerinic M. Ultrasound-integrated tight control in early psoriatic arthritis during adalimumab treatment. *Clin Exp Rheumatol* 2013;31:440-2.
 - 51 Glinatsi D, Bird P, Gandjbakhch F *et al.* Validation of the OMERACT Psoriatic Arthritis Magnetic Resonance Imaging Score (PsAMRIS) for the Hand and Foot in a Randomized Placebo-controlled Trial. *J Rheumatol* 2015;42:2473-9.
 - 52 Husic R, Gretler J, Felber A *et al.* Disparity between ultrasound and clinical findings in psoriatic arthritis. *Ann Rheum Dis* 2014;73:1529-36.
 - 53 Freeston JE, Coates LC, Nam JL *et al.* Is there subclinical synovitis in early psoriatic arthritis? A clinical comparison with gray-scale and power Doppler ultrasound. *Arthritis Care Res* 2014;66:432-9.
 - 54 Perrotta FM, Astorri D, Zappia M *et al.* An ultrasonographic study of enthesis in early psoriatic arthritis patients naive to traditional and biologic DMARDs treatment. *Rheumatol Int* 2016;36:1579-83.
 - 55 Marin J, Acosta-Felquer ML, Ferreyra-Garrot L *et al.* Patients with psoriatic arthritis fulfilling the minimal disease activity criteria do not have swollen and tender joints, but have active skin. *J Rheumatol* 2016;43:907-10.
 - 56 Filippucci E, Aydin SZ, Karadag O *et al.* Reliability of high-resolution ultrasonography in the assessment of Achilles tendon enthesopathy in seronegative spondyloarthropathies. *Ann Rheum Dis* 2009;68:1850-5.
 - 57 de Miguel E, Munoz-Fernandez S, Castillo C, Cobo-Ibanez T, Martin-Mola E. Diagnostic accuracy of enthesis ultrasound in the diagnosis of early spondyloarthritis. *Ann Rheum Dis* 2011;70:434-9.
 - 58 D'Agostino MA, Aegerter P, Bechara K *et al.* How to diagnose spondyloarthritis early? Accuracy of peripheral enthesitis detection by power Doppler ultrasonography. *Ann Rheum Dis* 2011;70:1433-40.
 - 59 D'Agostino MA, Said-Nahal R, Hacquard-Bouder C *et al.* Assessment of peripheral enthesitis in the spondylarthropathies by ultrasonography combined with power Doppler: a cross-sectional study. *Arthritis Rheum* 2003;48:523-33.
 - 60 Wang CH, Feng Y, Ren Z *et al.* Performance of ultrasound to monitor Achilles enthesitis in patients with ankylosing spondylitis during TNF- α antagonist therapy. *Clin Rheumatol* 2015;34:1073-8.
 - 61 Bonifati C, Elia F, Graceffa D *et al.* Clinical and contrast-enhanced ultrasound echography outcomes in psoriatic arthritis patients after one year of continuous therapy with anti-TNF drugs. *ISRN Dermatol* 2014;2014:932721.
 - 62 Marin J, Acosta-Felquer ML, Ferreyra-Garrot L *et al.* Utility of power Doppler ultrasound-detected synovitis for the prediction of short term flare in psoriatic arthritis patients in clinical remission [abstract]. *Arthritis Rheumatol* 2015;67(Suppl 10):161347.
 - 63 Diaz-Torne C, Moragues C, Toniolo E *et al.* Impact of ultrasonography on treatment decision in rheumatoid arthritis: the IMPULSAR study. *Rheumatol Int* 2017;37:891-6.
 - 64 Naredo E, Rodriguez M, Campos C *et al.* Validity, reproducibility, and responsiveness of a twelve-joint simplified power doppler ultrasonographic assessment of joint inflammation in rheumatoid arthritis. *Arthritis Rheum* 2008;59:515-22.
 - 65 Weckbach S, Schewe S, Michaely HJ *et al.* Whole-body MR imaging in psoriatic arthritis: additional value for therapeutic decision making. *Eur J Radiol* 2011;77:149-55.
 - 66 FitzGerald O, Haroon M, Giles JT, Winchester R. Concepts of pathogenesis in psoriatic arthritis: genotype determines clinical phenotype. *Arthritis Res Therapy* 2015;17:115.
 - 67 Naredo E, Moller I, de Miguel E *et al.* High prevalence of ultrasonographic synovitis and enthesopathy in patients with psoriasis without psoriatic arthritis: a prospective case-control study. *Rheumatology* 2011;50:1838-48.
 - 68 Gisondi P, Tinazzi I, El-Dalati G *et al.* Lower limb enthesopathy in patients with psoriasis without clinical signs of arthropathy: a hospital-based case-control study. *Ann Rheum Dis* 2008;67:26-30.
 - 69 Offidani A, Cellini A, Valeri G, Giovagnoni A. Subclinical joint involvement in psoriasis: magnetic resonance imaging and X-ray findings. *Acta Derm Venereol* 1998;78:463-5.
 - 70 Faustini F, Simon D, Oliveira I *et al.* Subclinical joint inflammation in patients with psoriasis without concomitant psoriatic arthritis: a cross-sectional and longitudinal analysis. *Ann Rheum Dis* 2016;75:2068-74.
 - 71 Moshrif A, Mosallam A, Mohamed EE, Gouda W, Doma M. Subclinical enthesopathy in patients with psoriasis and its association with other disease parameters: a power Doppler ultrasonographic study. *Eur J Rheumatol* 2017;4:24-8.
 - 72 Simon D, Faustini F, Kleyer A *et al.* Analysis of periarticular bone changes in patients with cutaneous psoriasis without associated psoriatic arthritis. *Ann Rheum Dis* 2016;75:660-6.
 - 73 Uluckan O, Wagner EF. Role of IL-17A signalling in psoriasis and associated bone loss. *Clin Exp Rheumatol* 2016;34(Suppl 98):17-20.
 - 74 Eder L, Thavaneswaran A, Chandran V, Cook RJ, Gladman DD. Obesity is associated with a lower probability of achieving sustained minimal disease activity state among patients with psoriatic arthritis. *Ann Rheum Dis* 2015;74:813-7.
 - 75 Pontifex EK, Gerlag DM, Gogarty M *et al.* Change in CD3 positive T-cell expression in psoriatic arthritis synovium correlates with change in DAS28 and magnetic resonance imaging synovitis scores following initiation of biologic therapy - a single centre, open-label study. *Arthritis Res Ther* 2011;13:R7.
 - 76 Cargill M, Schrodi SJ, Chang M *et al.* A large-scale genetic association study confirms IL12B and leads to the identification of IL23R as psoriasis-risk genes. *Am J Hum Genet* 2007;80:273-90.
 - 77 Durham LE, Kirkham BW, Taams LS. Contribution of the IL-17 pathway to psoriasis and psoriatic arthritis. *Curr Rheumatol Reports* 2015;17:55.
 - 78 Ritchlin CT, Kavanaugh A, Gladman DD *et al.* Treatment recommendations for psoriatic arthritis. *Ann Rheum Dis* 2009;68(9):1387-94.

- 79 Mumtaz A, Gallagher P, Kirby B *et al.* Development of a preliminary composite disease activity index in psoriatic arthritis. *Ann Rheum Dis* 2011;70:272–7.
- 80 Schoels MM, Aletaha D, Alasti F, Smolen JS. Disease activity in psoriatic arthritis (PsA): defining remission and treatment success using the DAPSA score. *Ann Rheum Dis* 2016;75:811–8.
- 81 Aletaha D, Alasti F, Smolen JS. Disease activity states of the DAPSA, a psoriatic arthritis specific instrument, are valid against functional status and structural progression. *Ann Rheum Dis* 2017;76:418–21.
- 82 Pincus T, Bergman MJ, Yazici Y. RAPID3—an index of physical function, pain, and global status as “vital signs” to improve care for people with chronic rheumatic diseases. *Bull NYU Hospital Joint Dis* 2009;67:211–25.
- 83 Gossec L, de Wit M, Kiltz U *et al.* A patient-derived and patient-reported outcome measure for assessing psoriatic arthritis: elaboration and preliminary validation of the Psoriatic Arthritis Impact of Disease (PsAID) questionnaire, a 13-country EULAR initiative. *Ann Rheum Dis* 2014;73:1012–9.