

## Superspace formulation of the local RG equation

---

Roberto Auzzi<sup>a,b</sup> and Boaz Keren-Zur<sup>c</sup>

<sup>a</sup>*Dipartimento di Matematica e Fisica, Università Cattolica,  
Via Musei 41, 25121 Brescia, Italy*

<sup>b</sup>*INFN Sezione di Perugia,  
Via A. Pascoli, 06123 Perugia, Italy*

<sup>c</sup>*Institut de Théorie des Phénomènes Physiques, EPFL,  
CH-1015 Lausanne, Switzerland*

*E-mail:* [roberto.auzzi@unicatt.it](mailto:roberto.auzzi@unicatt.it), [boaz.kerenzur@epfl.ch](mailto:boaz.kerenzur@epfl.ch)

**ABSTRACT:** We present the superspace formulation of the local RG equation, a framework for the study of supersymmetric RG flows in which the constraints of holomorphy and  $R$ -symmetry are manifest. We derive the consistency conditions associated with super-Weyl symmetry off-criticality and initiate the study of their implications. As examples, we derive an expression for the  $a$ -function, and present an analog of the  $a$ -maximization equation, which is valid off-criticality. We also apply this machinery to the study of conformal manifolds and give a simple proof that the metric on such manifolds is Kähler.

**KEYWORDS:** Superspaces, Anomalies in Field and String Theories, Renormalization Group

**ARXIV EPRINT:** [1502.05962](https://arxiv.org/abs/1502.05962)

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The generalized super-Weyl symmetry</b>	<b>3</b>
2.1	Background sources and the generating functional	3
2.2	Flavor symmetries	5
2.3	The super-Weyl symmetry	7
2.4	The superconformal symmetry	9
2.4.1	$R$ -symmetry and non-renormalization theorems	9
2.4.2	Anomalous super-Weyl weights	10
2.4.3	Primary operators	11
2.5	Ambiguity	11
2.6	Comparison with the general formulation	12
2.7	Consistency conditions	14
2.8	Super-Weyl variation of functions	14
2.8.1	SUSY covariant derivative	14
2.8.2	Lie derivative	15
2.8.3	Functions transforming covariantly under super-Weyl symmetry	15
<b>3</b>	<b>Consistency conditions for the generalized super-Weyl anomaly</b>	<b>16</b>
3.1	The main idea	16
3.2	The super-Weyl anomaly	17
3.3	The super-Weyl anomaly in the presence of chiral $U(1)$ anomalies	19
3.3.1	Chiral anomalies, ambiguity and Weyl anomaly coefficients	21
3.4	Scheme dependence of the anomaly	22
3.5	Consistency conditions	23
3.5.1	Constraints setting anomaly coefficients to zero	23
3.5.2	Constraints which can be used to algebraically eliminate anomaly coefficients	23
3.5.3	Reality constraints	24
3.5.4	Other constraints	25
3.5.5	Comments on the consistency conditions	26
<b>4</b>	<b>General implications</b>	<b>26</b>
4.1	$a$ -function and $a$ -maximization	26
4.1.1	$a$ -maximization	27
4.1.2	$\tilde{a}$ off-criticality	29
4.2	The conformal manifold	29
<b>5</b>	<b>Conclusions</b>	<b>31</b>

<b>A</b>	<b>Formulas and conventions</b>	<b>32</b>
A.1	Classical superWeyl variations	32
A.2	Covariant derivatives	33
A.3	Useful identities	33
<b>B</b>	<b>Useful definitions and results</b>	<b>33</b>
B.1	Notations	33
B.2	Lie derivatives in parameter space	34
B.3	Super-Weyl variations of functions	34
B.4	Integrations by parts	35
B.5	Computation of the consistency conditions	35
<b>C</b>	<b>Basic examples</b>	<b>36</b>
C.1	The Wess-Zumino model	36
C.2	Gauge theories	37
<b>D</b>	<b>The super-Weyl anomaly and the Zamolodchikov metric</b>	<b>38</b>

---

## 1 Introduction

Renormalization group (RG) flows describe a trajectory in the space of theories, induced by a change of scale. It is a major challenge of quantum field theory to characterize these trajectories, and understand their structure. One of the theoretical tools used for this purpose is the local RG equation, formulated in [1–3] (for recent reviews, see also [4–6]). The idea is to consider the RG evolution as a dilatation symmetry transformation, which is explicitly broken by interaction terms in the Lagrangian. Promoting the coupling constants  $\lambda$  to background fields  $\lambda(x)$ , the symmetry can be formally restored by assigning them with compensating transformation properties, determined by their  $\beta$ -function and anomalous dimensions. Moreover, introducing a background metric  $g_{\mu\nu}$ , the global dilatation symmetry can be promoted to a local Weyl symmetry. The local RG (LRG) equation is nothing but the anomalous Ward identity for the generalized Weyl symmetry. Roughly speaking, it takes the following form

$$\Delta_{\sigma}^W \mathcal{W}[g, \lambda] \equiv \int d^4x \sigma \left( 2g_{\mu\nu} \frac{\delta}{\delta g_{\mu\nu}(x)} + \beta \cdot \frac{\delta}{\delta \lambda(x)} + \dots \right) \mathcal{W}[g, \lambda] = \mathcal{A}_{\sigma}^W[g, \lambda], \quad (1.1)$$

where  $\mathcal{W}$  is the generating functional for correlation functions of composite operators,  $\Delta_{\sigma}^W$  is a generator for the Weyl symmetry transformation, and  $\mathcal{A}_{\sigma}^W$  is the anomaly function, which is a local function of the background fields. These objects will be defined in more detail below. The major results of this formalism are based on the Wess-Zumino consistency conditions for this anomaly

$$[\Delta_{\sigma_2}^W, \Delta_{\sigma_1}^W] \mathcal{W} = \Delta_{\sigma_2}^W \mathcal{A}_{\sigma_1} - \Delta_{\sigma_1}^W \mathcal{A}_{\sigma_2} = 0. \quad (1.2)$$

This equation gives non-trivial relations between the various anomaly coefficients, which are functions of the coupling constants. These constraints on functions of the couplings can be translated into constraints on the RG flow. A prominent example is the proof for the irreversibility of perturbative unitary RG flows [2, 3].

The goal of this work is to apply the formalism of the local RG equation to the study of supersymmetric RG flows. Supersymmetric RG flows are known to have a rich structure and non-trivial properties, such as the non-renormalization theorems for the superpotential [7] and the exact formula for the  $\beta$ -function in gauge theories [8, 9]. The derivation of these results is based on two properties of supersymmetric theories —  $R$ -symmetry and holomorphy. Supersymmetry can be introduced to the formalism of the LRG equation by directly specializing the field content and the couplings to the supersymmetric case, leading to some interesting constraints [10] (and more recently [6] and [11]). However, the formulation of the equation in components does not exploit the power of holomorphy. For this purpose we formulate the LRG equation using superspace notation,<sup>1</sup> with a supergravity background. The new equation we find, the superspace local RG (SLRG) equation, corresponds to the Ward-identity of the super-Weyl symmetry off-criticality, and it makes holomorphy and the relation with  $R$ -symmetry manifest.

In the framework described here, the non-renormalization theorem and the NSVZ formula [8, 9] appear as a natural part of the construction. In addition, we find a generalization of the consistency conditions (1.2), and begin the exploration of their implications. As in the framework of the LRG equation, we define a function  $\tilde{a}$ , which is a continuation off-criticality of the  $a$  coefficient in the Weyl anomaly, and find a relation between its derivative with respect to the coupling  $\lambda^I$  and the  $\beta$  function:

$$\frac{\partial}{\partial \lambda^I} \tilde{a} = \bar{\beta}^{\bar{J}} \chi_{I\bar{J}} - \beta^J (\partial_J \bar{w}_I - \partial_I \bar{w}_J) \tag{1.3}$$

where  $\chi$  and  $\bar{w}$  are functions of the couplings appearing as anomaly coefficients. A similar result is derived from the LRG consistency conditions. A remarkable result which appears only in the SLRG framework is that the tensor  $\chi_{I\bar{J}}$  can be written as a gradient of a function we denote by  $\Omega_I$ , plus a function of other anomaly coefficients ( $w_{\bar{J}}, \xi_{[IK]\bar{J}}^1, \zeta_{AI}^1$ ):

$$\chi_{I\bar{J}} - \partial_I w_{\bar{J}} - 2i \xi_{[IK]\bar{J}}^1 \beta^K + \zeta_{AI}^1 \partial_{\bar{J}} \Gamma^A = -\partial_{\bar{J}} \Omega_I. \tag{1.4}$$

In addition, we find a consistency condition which coincides with the  $a$ -maximization formula [13] at the fixed point. Plugging these results back into (1.3), we find the following expression for  $\tilde{a}$  in terms of the anomalous dimension matrices  $\gamma$

$$\tilde{a} = -\frac{1}{128\pi^2} \text{Tr}[\gamma^2] + \frac{1}{192\pi^2} \text{Tr}[\gamma^3] - \beta^I \tilde{\Omega}_I + \text{const}, \tag{1.5}$$

where  $\tilde{\Omega}_I = \Omega_I - \bar{w}_I$ . A similar ansatz for  $\tilde{a}$ , where  $\tilde{\Omega}_I$  was an undetermined function, was conjectured in [10] and checked to be consistent up to 4-loop order (see e.g. [6, 11, 14, 15]), under some assumptions regarding the anomalous dimensions, on which we comment below.

---

<sup>1</sup>Superfield formalism simplifies also the derivation of renormalization group equation for SUSY breaking parameters, using analytic continuation into superspace [12].

In [14–16]  $\tilde{\Omega}_I$  is interpreted as a Lagrange multiplier imposing the vanishing of the  $\beta$  function at the fixed point.

Another potential application of the SLRG framework is the study of manifolds of fixed points (conformal manifolds) [17]. To demonstrate the usefulness of this approach we give a simple proof for the fact that the Zamolodchikov metric on this manifold is Kähler (for an earlier proof see also [18]). We believe that the same machinery is suitable for the investigation of additional properties of the conformal manifold (as in [19] and [20]).

The paper is organized as follows: In section 2 we define the basic ingredients used in our formalism, namely the background fields and the generating functional  $\mathcal{W}$ , and present the generalized super-Weyl (SW) symmetry. We highlight the differences between the general formulation of the LRG equation and the superspace formulation (SLRG), which are the consequences of the choice of the holomorphic scheme. Using the generator of the SW symmetry we define a generator of infinitesimal super-conformal (SC) symmetry transformations acting on the sources. Using these expression we easily extract some constraints on the RG flow related to  $R$ -symmetry. We conclude with a discussion of the equation in components, and some technical aspects regarding the SW variation of functions.

In section 3 we define the super-Weyl anomaly, and its consistency conditions. There are several aspects in which this discussion differs from the presentation of the consistency conditions in [3]. First, following the methodology of [21], we introduce non-gauge-invariant SW anomaly terms which are necessary in the presence of chiral anomalies. Next, we discuss a peculiar feature of the Wess-Zumino consistency conditions of the SW anomaly, which implies that some of the consistency conditions are given only up to some unknown functions. Finally, and most importantly, we find many more equation than in the LRG framework. This is to be expected, as the SLRG equation describes a larger symmetry. An initial exploration of their implications, as described above, is given in section 4. Some of the relevant definitions and formulas appear in the appendix. Throughout this paper we use the notations of [22].

## 2 The generalized super-Weyl symmetry

### 2.1 Background sources and the generating functional

Our goal is to study supersymmetric RG flows in the vicinity of superconformal fixed points. The main tools at our disposal are a set of background source fields  $\mathcal{J}$  and a functional of these fields  $\mathcal{W}[\mathcal{J}]$ , which generates the renormalized correlation functions of the composite operators in the superconformal theory

$$\frac{\delta}{\delta \mathcal{J}_1(z_1)} \cdots \frac{\delta}{\delta \mathcal{J}_n(z_n)} \mathcal{W}[\mathcal{J}] \Big|_{\mathcal{J}=0} = i^{1-n} \langle \mathbf{T} \{ \mathcal{O}_1(z_1) \dots \mathcal{O}_n(z_n) \} \rangle . \quad (2.1)$$

Such a functional is schematically given (if the fixed point has a Lagrangian description) by

$$e^{-i\mathcal{W}[\mathcal{J}]} = \int D\Phi e^{-i \int (\mathcal{L}_0[\Phi] + \mathcal{J}\mathcal{O} + \mathcal{L}_{c.t.}(\mathcal{J}, \Phi))} \quad (2.2)$$

where  $\Phi$  are the dynamical fields and  $\mathcal{L}_0[\Phi]$  is the Lagrangian at the fixed point. A non-dynamical supergravity background, which is a useful way to realize the super-Weyl

symmetry of the theory, is implicit in this notation. In order to ensure the finiteness of the correlation functions,  $\mathcal{W}$  must contain a set of local counterterms  $\mathcal{L}_{c.t.}(\mathcal{J}, \Phi)$ , which consists of functions of the sources and their derivatives. The existence of these counterterms is responsible for the appearance of the anomaly. It should be emphasized, however, that eq. (2.1) is sufficient as a definition for  $\mathcal{W}$ , and that the formalism is valid also for conformal fixed points with no Lagrangian description.

The functional derivatives in eq. (2.1) evaluated with a small, non-zero, background value for  $\mathcal{J}$  gives the correlation functions in a theory where a deformation  $\int \mathcal{J}\mathcal{O}$  is turned on. In that sense, the background source fields  $\mathcal{J}$  can be understood as the coupling constants promoted to a coordinate dependent field. There is yet another way in which we can take advantage of these background fields: in a given theory, some of the global symmetries of the fixed point may be broken explicitly by the deformations. However, one can formally restore these symmetries by assigning transformation properties to the background fields, which compensate for the non-invariance of the theory. The background fields  $\mathcal{J}$  thus play several different roles in our framework: coupling constants, sources for composite operators, and compensators for broken symmetries.

The idea of compensating background fields is well known: for example, it has been exploited in the past for the study of broken flavor symmetries [23]. The application of this idea to Weyl symmetry, which was formulated in [3], is the relevant realization of this idea for our discussion. In that paper, the background fields were used to define a generalized form of the Weyl symmetry, which is valid off-criticality, and the consistency conditions associated with this symmetry [24] lead to non-trivial constraints on RG flows. Here we specialize to supersymmetric theories and we take this idea one step further: we introduce superspace notations, and consider the sources as compensators of the super-Weyl (SW) symmetry of the super-conformal (SC) fixed point.

We will restrict our discussion to RG flows induced by marginal deformation. As discussed in [20], in a flat background there are only two possible options for such a deformation: chiral operators integrated over  $d^2\theta$  (superpotential deformations) which we denote by  $\mathcal{O}_I$ , or generic operators integrated in  $d^4\theta$  (Kähler deformations) which we denote by  $J_{\bar{i}j}$  (the structure of the indices will be explained in the next section). In order for the Kähler deformation  $J$  to be marginal, it should correspond to a conserved current ( $\bar{D}^2 J = 0$ ), and then  $\int d^4\theta J$  does not deform the Lagrangian. If the current  $J$  is not conserved, then the deformation is irrelevant by unitarity and will be neglected in the following analysis.

A general discussion of RG flows requires introduction of an infinite set of sources, for an infinite set of operators. However, for the discussion of RG flows induced by marginal deformations, with no mass parameters, it is sufficient to introduce sources for the marginal operators only. We will consider a chiral source  $\lambda^I$  and a real source  $Z^{j\bar{i}}$  defined such that<sup>2</sup>

$$\frac{\delta}{\delta\lambda^I(z)}\mathcal{W} \equiv [\mathcal{O}_I(z)] \quad \frac{\delta}{\delta Z^{j\bar{i}}(z)}\mathcal{W} \equiv [J_{\bar{i}j}(z)]. \quad (2.3)$$

In our notations, the brackets mean that the operator is renormalized. The source  $Z$  can be understood as the normalization of the kinetic term. For example, in the case of a

---

<sup>2</sup>We use the conventions  $\frac{\delta\lambda(z_+)}{\delta\lambda(z_+)} = \varphi^{-3}\delta^{(4)}(x-x')\delta^{(2)}(\theta-\theta')$ ,  $\frac{\delta Z(z)}{\delta Z(z')} = E\delta^{(4)}(x-x')\delta^{(2)}(\theta-\theta')\delta^{(2)}(\bar{\theta}-\bar{\theta}')$ .

Wess-Zumino model the coupling of the sources takes the form

$$\int d^8z E^{-1} Z^{j\bar{i}} J_{\bar{i}j} + \int d^6z \varphi^3 \lambda^I \mathcal{O}_I + \text{c.c.} \tag{2.4}$$

where

$$J_{\bar{i}j} \equiv \bar{\Phi}_{\bar{i}} \Phi_j, \quad \lambda^I \mathcal{O}_I \equiv \lambda^{ijk} \Phi_i \Phi_j \Phi_k. \tag{2.5}$$

Alternatively, one can work in components, set the normalization of the kinetic terms to unity and absorb the wavefunction renormalization into the bare fields. However, by doing that one loses information about the constraints imposed by holomorphy. Our choice to work in the holomorphic scheme forces us to introduce  $Z$  as an independent source, and to work with the non-physical holomorphic coupling constants  $\lambda$ .

As mentioned above, the generating functional is defined in a curved supergravity background. We will use the old-minimal formulation of supergravity, and the notations of [22]. For the purpose of the discussion here, it is sufficient to mention that the supergravity fields include a superfield containing the background metric  $H^{\alpha\dot{\alpha}}$  and a chiral field  $\varphi$  known as the chiral compensator.  $\varphi$  will play an important role in the realization of the super-Weyl symmetry on  $\mathcal{W}$ . These fields can also be understood as the sources for the supercurrent (Ferrara-Zumino multiplet [25])  $\mathcal{T}_{\alpha\dot{\alpha}}$  and a chiral operator  $\mathcal{T}$

$$\frac{\delta}{\delta H^{\alpha\dot{\alpha}}(z)} \mathcal{W} = -\frac{1}{2} [\mathcal{T}_{\alpha\dot{\alpha}}(z)], \quad \frac{\delta}{\delta \varphi^3(z)} \mathcal{W} \equiv \frac{1}{3} [\mathcal{T}(z)], \tag{2.6}$$

with which we can write the supersymmetric generalization of the conservation of energy momentum tensor:

$$0 = \bar{\mathcal{D}}^{\dot{\alpha}} [\mathcal{T}_{\alpha\dot{\alpha}}] + \frac{2}{3} \mathcal{D}_{\alpha} [\mathcal{T}]. \tag{2.7}$$

Using this Ward identity we can write the  $\theta^2$  component of the chiral superfield  $\mathcal{T}$  as

$$\mathcal{T}|_{\theta^2} = \frac{1}{2} T_{\mu\nu}^{\mu} + i \frac{3}{4} \partial_{\mu} j_5^{\mu} \tag{2.8}$$

where  $T_{\mu\nu}$  is the energy momentum tensor and  $j_5^{\mu}$  is the lowest component of the supercurrent, which at the fixed point coincides with the  $R$  current.

## 2.2 Flavor symmetries

As mentioned in the previous section, the coupling constants which induce the RG flow break explicitly some of the symmetries of the fixed point. In this section we discuss global, internal, perhaps anomalous, symmetries, which we will denote by  $G$  and refer to as “flavor” symmetries. We will assume that the marginal operators  $\mathcal{O}_I$  reside in some representation of the symmetry group, and their transformation rule is given by

$$\delta_{\omega}^G \mathcal{O}_I = -\omega^A \mathcal{O}_J (T_A)^J_I \tag{2.9}$$

where  $(T_A)_J^I$  are the generators of the symmetry, and  $\omega^A$  is some constant transformation parameter. Obviously, the global symmetry can be restored (up to anomalies) by assigning the following transformation properties to the fields

$$\delta_\omega^G \lambda^I = \omega^A (T_A)_J^I \lambda^J . \tag{2.10}$$

The framework of the background sources and generating functional  $\mathcal{W}$  can be used to generate the Ward identities for the flavor symmetries. This is done by promoting the global symmetry to a local one and using background gauge fields. Conveniently, the necessary background fields were already introduced in the above discussion. Indeed, the sources  $Z$  can be interpreted either as wavefunction renormalization, or equivalently, as gauge fields which act as sources for the Noether currents of the theory. In the example of a Wess-Zumino model, we can use the more familiar notation  $Z^{i\bar{j}} \equiv (e^{-V})^{i\bar{j}}$ , where  $V$  is a real vector superfield,<sup>3</sup> and we can write

$$Z^{i\bar{j}} \Phi_i \bar{\Phi}_{\bar{j}} \equiv \Phi_i (e^{-V})^{i\bar{j}} \bar{\Phi}_{\bar{j}} \tag{2.11}$$

and Noether currents can be defined as

$$(e^V)_{\bar{j}i} (T_A)_k^i \frac{\delta}{\delta (e^V)_{\bar{j}k}} \mathcal{W} = -[\Phi_i (T_A)_k^i (e^{-V})^{k\bar{j}} \bar{\Phi}_{\bar{j}}] \equiv [J_A] . \tag{2.12}$$

The indices  $i$  ( $\bar{i}$ ) run in the fundamental (antifundamental) of the global symmetry group (which may be broken by the chiral couplings  $\lambda^I$ ). It is always possible to decompose the representation of the chiral couplings  $\lambda^I$  in product of fundamentals; from this decomposition we can infer how  $e^V$  acts on the couplings  $\lambda^I$ .

A few words about our index convention are in order. A super-gauge transformation can be parameterized by the exponential of a chiral gauge parameter:  $(e^{i\Lambda})_j^i$ , where  $\Lambda = \Lambda^A T_A$  and  $T_A$  is a basis of hermitian matrices. As a mnemonic device, the indices for the sources are chosen as:

$$\lambda^I, \quad \bar{\lambda}^{\bar{I}}, \quad (e^V)_{\bar{j}k}, \quad (e^{-V})^{i\bar{j}} . \tag{2.13}$$

The gauge field  $e^V$  and its inverse transform as:

$$(e^V)_{\bar{j}k} \rightarrow (e^{i\bar{\Lambda}})_{\bar{j}}^{\bar{l}} (e^V)_{\bar{l}i} (e^{-i\Lambda})_k^i, \quad (e^{-V})^{i\bar{j}} \rightarrow (e^{i\Lambda})_k^i (e^{-V})^{k\bar{l}} (e^{-i\bar{\Lambda}})_{\bar{l}}^{\bar{j}} . \tag{2.14}$$

The chiral coupling transform as:

$$\lambda^I \rightarrow (e^{i\Lambda})_J^I \lambda^J, \quad \bar{\lambda}^{\bar{I}} \rightarrow \bar{\lambda}^{\bar{J}} (e^{-i\bar{\Lambda}})_{\bar{J}}^{\bar{I}} . \tag{2.15}$$

The representation according to which  $\lambda^I$  transforms can be obtained from some opportune tensor product of the fundamental representation, which has indices  $i$ . It is sometimes convenient to write objects as  $Y_j^i$  and  $\bar{Y}_{\bar{j}}^{\bar{i}}$  in term of adjoint indices:  $Y_j^i = Y^A (T_A)_j^i$ , where matrices  $(T_A)_j^i$  are the hermitian generators of the fundamental representation. We can then write the same object in an arbitrary representation; for example, in the representation

---

<sup>3</sup>In our normalization the  $\theta\sigma^\mu\bar{\theta}$  component of  $V$  is twice the ‘‘canonical’’ gauge field  $A_\mu$ , which is defined as the one appearing in the covariant derivated of fermions as  $D_\mu\psi_\Phi = \partial_\mu\psi_\Phi - iA_\mu^A T_A\psi_\Phi$ .



of the chiral coupling:  $Y_J^I = Y^A(T_A)_J^I$ . We denote with  $\bar{T}_A$  the complex conjugate of the generator  $T_A$ :  $(\bar{T}_A)_{\bar{j}}^{\bar{i}} = ((T_A)_j^i)^* = (T_A)_i^j$ ; with this notation  $\bar{Y}_{\bar{j}}^{\bar{i}} = \bar{Y}^A(\bar{T}_A)_{\bar{j}}^{\bar{i}}$ . The generators of the anti-fundamental representation are:  $(\tilde{T}_A)_{\bar{j}}^{\bar{i}} = -(\bar{T}_A)_{\bar{j}}^{\bar{i}}$ .

In general, one may consider chiral operators, and associated chiral sources, which are invariant under all flavor symmetries of the fixed point. However, the occurrence of such singlet chiral sources is rare (in fact the only example we are aware of are the holomorphic gauge couplings in pure super Yang-Mills theory). Moreover, in most cases it is impossible to write a holomorphic function of the  $\lambda$ 's which is a singlet of the symmetries. We refer to this property as the ‘‘absence of chiral singlets’’. In theories where it applies, we will show below that it can be used to derive strong constraints on the flow.

We are now ready to define a generator for the flavor symmetry transformations

$$\begin{aligned} \Delta_\Lambda^G \equiv & \int d^6z \varphi^3 i(\Lambda\lambda)^I \frac{\delta}{\delta\lambda^I} + c.c \\ & - \int d^8z E^{-1} (i(e^V \Lambda)_{\bar{i}j} - i(\bar{\Lambda} e^V)_{\bar{i}j}) \frac{\delta}{\delta(e^V)_{\bar{i}j}}, \end{aligned} \tag{2.16}$$

The anomalous invariance equation for  $\mathcal{W}$  is written as

$$\Delta_\Lambda^G \mathcal{W} = \mathcal{A}_\Lambda^G, \tag{2.17}$$

where  $\mathcal{A}_\Lambda^G$  is the anomaly involving background gauge fields, which we will further discuss in section 3.3. Anomalies involving dynamical gauge fields are realized in this formula by assigning transformation properties to the holomorphic gauge couplings (see section C.2). Taking the transformation parameter  $\Lambda(z)$  to be the chiral delta function  $\delta_+(z - y)$  and evaluating in the flat and constant background, we find the operator equation

$$(T_A \lambda)^I [\mathcal{O}_I(y)] = -\frac{1}{4} \bar{D}^2 [J_A(y)]. \tag{2.18}$$

In this equation the anomaly vanishes because we take the background sources to be constant. Finally, the Ward identity for a correlation function is given by

$$\begin{aligned} \Delta_\Lambda^G \frac{\delta}{\delta\lambda^{I_1}(z_1)} \cdots \frac{\delta}{\delta\lambda^{I_n}(z_n)} \mathcal{W} = & \left[ \Delta_\Lambda^G, \frac{\delta}{\delta\lambda^{I_1}(z_1)} \cdots \frac{\delta}{\delta\lambda^{I_n}(z_n)} \right] \mathcal{W} \\ & + \frac{\delta}{\delta\lambda^{I_1}(z_1)} \cdots \frac{\delta}{\delta\lambda^{I_n}(z_n)} \mathcal{A}_\Lambda^G. \end{aligned} \tag{2.19}$$

### 2.3 The super-Weyl symmetry

In a supergravity background,  $\mathcal{W}$  possess another important symmetry, which is broken by the marginal deformations — the super-Weyl (SW) symmetry. This symmetry is realized as a variation of the chiral compensator  $\delta_\sigma^{\text{SW}} \varphi = \sigma \varphi$ . At the fixed point this symmetry is generated by

$$\Delta_\sigma^{\text{SW}} \equiv \int d^6z \sigma \varphi \frac{\delta}{\delta\varphi} + c.c \tag{2.20}$$

As discussed in [3], in order to formally restore the Weyl symmetry broken by quantum effects, the background sources  $\lambda$  and  $Z$  (or equivalently  $V$ ) must be given Weyl transformation properties, which compensate for the running of the couplings. The Weyl symmetry can thus be generated by an operator similar to the one defined in eq. (2.16):

$$\begin{aligned} \Delta_\sigma^{\text{SW}} \equiv & \int d^6z \varphi^3 \sigma \left( 3 \frac{\delta}{\delta \varphi^3} + b^I \frac{\delta}{\delta \lambda^I} \right) + c.c \\ & + \int d^8z E^{-1} (\sigma \Gamma^A (e^V T_A)_{\bar{i}j} + \bar{\sigma} \bar{\Gamma}^A (\bar{T}_A e^V)_{\bar{i}j}) \frac{\delta}{\delta (e^V)_{\bar{i}j}} . \end{aligned} \quad (2.21)$$

We rewrote the variation of  $\varphi$  to allow usage of eq. (2.6).  $b^I = b^I(\lambda)$  is a holomorphic function of chiral couplings we will refer to as the holomorphic  $\beta$ -function, and  $\Gamma^A = \Gamma^A(\lambda, \bar{\lambda}, e^V)$  is a superfield which contains the anomalous dimensions of the dynamical fields. In the following sections we will discuss these functions and the model independent constraints imposed on them by the different symmetries of the theory.

As mentioned above, a full analysis of the Weyl symmetry off-criticality requires introduction of sources for each of the composite operators in the spectrum, and not only the marginal ones. However, for the discussion of a flow in which none of the dimensionful parameters obtains a non-zero VEV, it suffices to introduce only dimensionless sources. There is one exception we are aware of — there could be contributions to the local RG equations from D-terms involving chiral dimension 2 operators (e.g.  $\int d^8z E^{-1} \eta \Phi^2$ , where  $\eta$  is some function of the couplings). However, such terms can be eliminated by a choice of improvement (see, e.g. discussion in [26, 27]), and we leave the study of such terms for the future.

The supersymmetric local RG (SLRG) equation, which is the subject of our discussion, is nothing but the (anomalous) Ward identity for this symmetry

$$\Delta_\sigma^{\text{SW}} \mathcal{W} = \mathcal{A}_\sigma^{\text{SW}} . \quad (2.22)$$

As an operator equation (going to the flat and constant background) this can be written as

$$[\mathcal{T}] = - \left( b^I [\mathcal{O}_I] - \frac{1}{4} \Gamma^A \bar{\nabla}^2 [J_A] \right) \quad (2.23)$$

Using the operator equation (2.18) one can rewrite this in a more familiar form

$$[\mathcal{T}] = -\beta^I [\mathcal{O}_I] \quad (2.24)$$

where we identify the physical  $\beta$ -function (which corresponds to the numerator of NSVZ [8]) as

$$\beta^I \equiv b^I + \Gamma^A (T_A \lambda)^I . \quad (2.25)$$

Plugging this into eq. (2.8), and taking the real part, we find the expected trace anomaly

$$T_\mu^\mu = \beta^I [O_I] + c.c \quad (2.26)$$

where  $O_I$  is minus the  $\theta^2$  component of  $\mathcal{O}_I$ .

## 2.4 The superconformal symmetry

The superconformal symmetry can be defined as a combination of super-diffeomorphisms and super-Weyl transformations which keep the superspace interval, and equivalently, the chiral compensator  $\varphi$ , fixed. Following [28, 29], the infinitesimal diff transformations, corresponding to superconformal symmetry are parameterized by a vector  $h^\mu$  and a spinor  $\eta^\alpha$  (which is a function of  $h$ ) and generated by the operator

$$\mathcal{L}_h^{diff} \equiv h^\mu \partial_\mu + \eta^\alpha \partial_\alpha + \bar{\eta}_{\dot{\alpha}} \bar{\partial}^{\dot{\alpha}} . \quad (2.27)$$

Under such diff-transformation of the coordinates the compensator  $\varphi$  transforms as

$$\delta_h \varphi = \sigma_h \varphi \quad (2.28)$$

where  $\sigma_h$  is some chiral superfield which is a function of the transformation parameters. This variation can be canceled by a further Weyl rescaling, generated using the operator  $\Delta_\sigma^{SW}$ . Combining the two operations, we can define a generator of superconformal transformations which is schematically given by:

$$\Delta_h^{SC} = \int d^8 z E^{-1} \mathcal{L}_h^{diff} [\mathcal{J}] \frac{\delta}{\delta \mathcal{J}} - \Delta_{\sigma_h}^{SW} . \quad (2.29)$$

The transformation properties of the composite operators under the SC symmetry are obtained by computing the commutator

$$\delta^{SC} \left( \frac{\delta}{\delta \mathcal{J}} \mathcal{W} \right) \equiv \left[ \Delta_h^{SC}, \frac{\delta}{\delta \mathcal{J}} \right] \mathcal{W} . \quad (2.30)$$

The super-diff parameters  $h$  and  $\eta$  corresponding to infinitesimal dilatations (with a scale factor  $d$ ) and  $R$  symmetry rotations (with a phase  $r$ ) are

$$\begin{aligned} h^\mu &= x^\mu d + 2(\theta \sigma^\mu \bar{\theta}) r \\ \eta^\alpha &= \theta^\alpha \left( \frac{1}{2} d + i r \right) . \end{aligned} \quad (2.31)$$

The corresponding infinitesimal diff generator and Weyl transformation parameters are

$$\begin{aligned} \mathcal{L}_{h(d,r)}^{diff} &= d \left( x^\mu \partial_\mu + \frac{1}{2} \theta^\alpha \frac{\partial}{\partial \theta^\alpha} + \frac{1}{2} \bar{\theta}^{\dot{\alpha}} \frac{\partial}{\partial \bar{\theta}^{\dot{\alpha}}} \right) + i r \left( \theta^\alpha \frac{\partial}{\partial \theta^\alpha} - \bar{\theta}^{\dot{\alpha}} \frac{\partial}{\partial \bar{\theta}^{\dot{\alpha}}} \right) \\ \sigma_{h(d,r)} &= d - i \frac{2}{3} r . \end{aligned} \quad (2.32)$$

### 2.4.1 $R$ -symmetry and non-renormalization theorems

Let us write the generator of  $R$  symmetry off-criticality using eqs. (2.29)

$$\begin{aligned} \Delta_{h(r)}^{SC} &= \int d^6 z \varphi^3 \left( (\mathcal{L}_{h(r)}^{diff} \lambda)^I + i \frac{2}{3} r b^I \frac{\delta}{\delta \lambda^I} \right) + c.c \\ &+ \int d^8 z E^{-1} \left( (\mathcal{L}_{h(r)}^{diff} e^V)_{i\bar{j}} + i \frac{2}{3} r ((e^V \Gamma)_{i\bar{j}} - (\bar{\Gamma} e^V)_{i\bar{j}}) \right) \frac{\delta}{\delta (e^V)_{i\bar{j}}} + c.c . \end{aligned} \quad (2.33)$$

The first implication of this expression is that if we want to avoid assigning a non-vanishing, coupling dependent,  $R$ -charge to the background gauge fields, we must consider only hermitean  $\Gamma$  matrices

$$\bar{\Gamma}_{\bar{i}}^{\bar{k}}(e^V)_{\bar{k}j} = (e^V)_{\bar{i}k}\Gamma_j^k. \quad (2.34)$$

This constraint will be applied in all the results described in this paper.

Another constraint, with far reaching implications, is derived from the requirement that the  $R$ -symmetry acts linearly on the sources. By inspection of (2.33), this can be satisfied only if the holomorphic  $\beta$ -function can be written as

$$b^I = (q\lambda)^I \quad (2.35)$$

where  $q_I^J$  is some matrix of numbers which is at the moment unconstrained. We conclude that the holomorphic  $\beta$  function, as defined here, is either vanishing or linear in the couplings. This result is consistent with the non-renormalization theorem for the superpotential [7]. Indeed, superpotentials terms have vanishing holomorphic  $\beta$ -function,<sup>4</sup> and as explained in the next section, this corresponds to the fact that they have  $R$ -charge 2.

Our conclusion seems to be in contradiction with the well known fact that the holomorphic  $\beta$ -function of the holomorphic gauge couplings  $S \equiv \frac{4\pi}{g_h^2} - i\frac{\Theta}{2\pi}$ , is a constant. However, as a source for the marginal deformation associated with a gauge theory, we use the background field  $\lambda_G = e^{-S}$  (see appendix C.2 for a more detailed descriptions of our conventions for gauge couplings). This basis for the background sources has the virtues of transforming linearly under the Konishi symmetry, and vanishing in the limit of the free theory, and therefore it is more suitable to our formalism. In this basis, we find that holomorphic  $\beta$ -function is linear, in agreement with our discussion above.

#### 2.4.2 Anomalous super-Weyl weights

Using the commutator with  $\Delta_{h(d,r)}^{SC}$  we can compute the variation of the renormalized operators. We find the following transformation properties, evaluated with flat background:

$$\begin{aligned} \delta_d^{SC}[\mathcal{O}_I] &= -\mathcal{L}_{h(d)}^{diff}[\mathcal{O}_I] - d(3\delta_I^J + \gamma_I^J + \tilde{\gamma}_I^J - \Gamma^A(T_A)_I^J)[\mathcal{O}_J] \\ \delta_r^{SC}[\mathcal{O}_I] &= -\mathcal{L}_{h(r)}^{diff}[\mathcal{O}_I] + ir\left(2\delta_I^J - \frac{2}{3}q_I^J\right)[\mathcal{O}_J] \\ \delta_d^{SC}[J_A] &= -\mathcal{L}_{h(d)}^{diff}[J_A] - d(2\delta_A^B + 2\gamma_A^B)[J_B] \\ \delta_r^{SC}[J_A] &= -\mathcal{L}_{h(r)}^{diff}[J_A] \end{aligned} \quad (2.36)$$

where the generator of superdiffs is defined in (2.32) and the anomalous dimension matrices are given by

$$\gamma_I^J = \partial_I\beta^J \quad \tilde{\gamma}_I^J = \partial_I\Gamma^A(T_A\lambda)^I \quad \gamma_A^B = \partial_I\Gamma^B(T_A\lambda)^I. \quad (2.37)$$

---

<sup>4</sup>Due to an inherent ambiguity in our definition of  $b$ , see section 2.5, there could be parameterization of the local RG equation in which the holomorphic beta function  $b$  is linear in the coupling. This has no physical significance, because the physical beta function  $\beta$  is independent of this ambiguity.

Basing on the second line of (2.36) we can also define the  $R$  charge matrix  $T_R$  for the composite operators

$$(T_R)_I^J \equiv 2\delta_I^J - \frac{2}{3}q_I^J \tag{2.38}$$

where  $q$  is the matrix defined in (2.35). At the fixed point, where  $\Gamma^A(T_A\lambda) = -(q\lambda)^I$ , we find the expected relation between the anomalous dimension and the  $R$  charge

$$\Gamma^A T_A = \frac{3}{2}(T_R - 2) . \tag{2.39}$$

### 2.4.3 Primary operators

It is also useful to compute the action of the special superconformal symmetry on the operators (with the corresponding fermionic transformation parameters  $s^\alpha$ ). At the fixed point, one would expect the primary operators  $\mathcal{O}_I$  to be annihilated by the generator of this transformations. The corresponding SW transformation parameter is given by

$$\sigma_{h(s)} \propto s^\alpha \theta_\alpha . \tag{2.40}$$

Using the commutator (2.30) we find the following result

$$\delta_s^{SC}[\mathcal{O}_I] \propto \bar{s}_{\dot{\alpha}} \partial_I \bar{\Gamma}^A \bar{\nabla}^{\dot{\alpha}} [J_B] . \tag{2.41}$$

We conclude that at the fixed point

$$\partial_I \bar{\Gamma}^A = 0 \quad \forall \mathcal{O}_I \text{ primary} . \tag{2.42}$$

Due to the Ward identities of the flavor symmetries, there are linear combinations of the marginal operators,  $(T_A\lambda)^I \mathcal{O}_I$ , which can be written as derivatives of currents, and therefore are not primary. Such operators do not necessarily satisfy condition (2.42).

## 2.5 Ambiguity

In the presence of a non-anomalous  $U(1)$  symmetries, there is an equivalence class of local RG equations, related by addition of  $U(1)$  Ward identities

$$\Delta_\sigma^{\text{SW}} \rightarrow (\Delta_\sigma^{\text{SW}})' = \Delta_\sigma^{\text{SW}} + \Delta_{\Lambda=ia\sigma}^G , \tag{2.43}$$

where  $a^A$  is a vector of numbers<sup>5</sup> defining the linear combination of  $U(1)$  symmetries added to the local RG equation (in fact, as will be explained in section 3.3, this ambiguity exists also in the case of anomalous  $U(1)$ s). This ambiguity can be translated into an ambiguity in the holomorphic function  $b$  and the anomalous dimension  $\Gamma$

$$\Delta_\sigma^{\text{SW}}(b^I, \Gamma^A) \rightarrow \Delta_\sigma^{\text{SW}}(b^{I'}, \Gamma^{A'}) = \Delta_\sigma^{\text{SW}}(b^I, \Gamma^A) + \Delta_{\Lambda=ia\sigma}^G , \tag{2.44}$$

---

<sup>5</sup>In principle we might contemplate the possibility that  $a^A$  as is a non-constant holomorphic function of the chiral couplings, but this would lead to a contradiction with the argument from the previous section regarding the linear transformation of the sources under the  $R$ -symmetry.

where

$$\begin{aligned} b^{I'} &= b^I - a^A (T_A \lambda)^I \\ \Gamma^{A'} &= \Gamma^A + a^A . \end{aligned} \tag{2.45}$$

Under this transformation the  $R$  charge defined in (2.38) changes as

$$T'_R = T_R + \frac{2}{3} a^A T_A . \tag{2.46}$$

Recall that in the local RG equation we are keeping only the information about composite operators, while the elementary fields, which are not assigned with independent sources, can be left unspecified. The freedom to redefine  $(\Gamma^A, b^I)$  is due to the fact that we can change the dimensions of the elementary component fields, as long as we keep the dimensions of the composite operators unchanged. This ambiguity is, therefore, not physical. The physical  $\beta$  function and the anomalous dimension matrices of the composite operators, defined in eq. (2.37), are insensitive to this ambiguity.

Due to this ambiguity, even in theories where the holomorphic  $\beta$ -functions vanishes, there could be, in principle, parameterization of the equation in which the  $b$  is linear in the couplings. This is still in agreement with our non-renormalization theorem of section 2.4.1. Finally, let us comment that in the general formulation of the local RG equation there is an ambiguity which is similar in form but of different origin. As was demonstrated in [30], in certain cases there is a freedom to define a phase difference between the bare and renormalized sources. This ambiguity in the definition of the bare Lagrangian leads to an ambiguity in the  $\beta$  function. However, this artifact does not appear in the holomorphic scheme we are using, in which the normalization of the wavefunction is kept as an independent field.

## 2.6 Comparison with the general formulation

The original general expression for the LRG equation in the presence of scalar marginal operators, and vectors of dimension 3, is given by

$$\Delta_\sigma^W = \int d^4x \left( 2\sigma g_{\mu\nu} \frac{\delta}{\delta g_{\mu\nu}(x)} + \sigma \beta^I \frac{\delta}{\delta \lambda^I(x)} + (\sigma \rho_I^A \nabla_\mu \lambda^I - \partial_\mu \sigma S^A) \frac{\delta}{\delta A_\mu^A(x)} \right) \tag{2.47}$$

where  $g_{\mu\nu}$  is the metric,  $A_\mu^A$  are the background gauge fields, while  $\rho$  and  $S$  are some covariant functions of the sources  $\lambda$  [3].

An important observation is that in this formalism there is no source for the kinetic term, which corresponds to choosing canonical wave function renormalization. In the superspace formulation such a term exists (related to the lowest component of  $\Gamma$ ), and therefore the matching of the Ward identities in the two formulations is non-trivial. However, limiting ourselves to the operator equation, i.e. (2.23), we can eliminate the contribution of the kinetic term, and compare the resulting equations. For this purpose we focus on a specific point in parameter space, which is defined by the specific value  $\tilde{\lambda}$  for the coupling constants. We then use the ambiguity discussed in section 2.5 to subtract from  $\Gamma^A$  the

numerical value of its lowest component at that point, which we denote by  $\tilde{\Gamma} \equiv \Gamma(\tilde{\lambda}, \bar{\tilde{\lambda}})$  (it is not possible to subtract  $\Gamma^A$  as a function, because it is not holomorphic):

$$\Delta_{\sigma}^{\text{SW}}(b^I, \Gamma^A) \mathcal{W} \Big|_{\lambda=\tilde{\lambda}} = \Delta_{\sigma}^{\text{SW}}(b^I + \tilde{\Gamma}^A (T_A \lambda)^I, (\Gamma^A - \tilde{\Gamma}^A)) \mathcal{W} \Big|_{\lambda=\tilde{\lambda}} + \Delta_{\Lambda=i\sigma\tilde{\Gamma}}^G \mathcal{W} \Big|_{\lambda=\tilde{\lambda}}. \tag{2.48}$$

Not surprisingly, the elimination of the wave function renormalization leads to the appearance to the physical  $\beta$ -function  $\beta^I = b^I + \tilde{\Gamma}^A (T_A \lambda)^I$ .

Another step necessary to make the comparison is to focus on the case where just the real part of the lowest component of the super-Weyl transformation parameter  $\sigma$ , which we denote by  $\sigma_c$ , is non zero. To keep things simple, just the lowest component of the chiral coupling  $\lambda$  is taken non-zero, and  $e^V$  is written in the WZ gauge:

$$\begin{aligned} \lambda^I &= \lambda_c^I + i\theta\sigma^\mu\bar{\theta}\partial_\mu\lambda_c^I + \frac{\theta^2\bar{\theta}^2}{4}\square\lambda_c^I \\ (e^V)_{\bar{i}j} &= \delta_{\bar{i}j} + 2\theta\sigma^\mu\bar{\theta}(A_\mu)_{\bar{i}j} - \theta^2\bar{\theta}^2\left((A_\mu A^\mu)_{\bar{i}j} + m_{\bar{i}j}^2\right) \end{aligned} \tag{2.49}$$

where the  $D$ -term of  $V$  (the soft mass) is denoted by  $m^2$ . Expanding the generator of the SW symmetry in components, and comparing with (2.47) we find

$$\beta^I = b^I + \Gamma^A (T_A \lambda)^I, \quad \rho_I^A = -\partial_I \Gamma^A, \quad \rho_{\bar{I}}^A = \partial_{\bar{I}} \Gamma^A, \quad S^A = 0, \tag{2.50}$$

consistent with the results of [30]. A non-vanishing  $S^A$  function could, in principle, appear if  $\Gamma$  had an anti-hermitean components. The vanishing of  $S^A$  is, therefore, related to the  $R$ -symmetry constraints discussed in section 2.4.1. This relation between  $R$ -symmetry and the vanishing of  $S$  was discussed in [30].<sup>6</sup>

As explained in [3], in the presence of dimension two operators there could be contributions to the local RG equation which are non-vanishing in the limit where all the mass parameters are set to zero. In the SUSY case, one type of such contributions is related to holomorphic dimension 2 operators. As mentioned above, we chose to leave the discussion of such terms for future study. Another kind of dimension 2 operators are the soft SUSY-breaking mass term for the scalars in a chiral multiplet (sourced by the  $D$ -term of  $e^V$ ):

$$\Delta_{\sigma}^W \supset - \int d^4x \sigma (D_{\bar{I}}^a \nabla^2 \lambda^I + E_{IJ}^a \nabla_{\mu} \lambda^I \nabla^{\mu} \lambda^J) \frac{\delta}{\delta(m^2)^a}, \tag{2.51}$$

where in the case of the Wess-Zumino model the index  $a$  can be replaced by the product of a fundamental and anti-fundamental indices  $i\bar{j}$ :

$$D_{\bar{I}}^{i\bar{j}} = -\partial_{\bar{I}} \Gamma^A (T_A)^i_j \delta^{j\bar{j}}, \quad E_{IJ}^{i\bar{j}} = -\partial_{IJ} \Gamma^A (T_A)^i_j \delta^{j\bar{j}}. \tag{2.52}$$

---

<sup>6</sup>Notice, however, that this constraint is an artifact of the holomorphic scheme. It is not a direct consequence of supersymmetry. When working in components one can choose a scheme in which  $S^A$  is non-zero.

## 2.7 Consistency conditions

More constrains on  $b$  and  $\Gamma$  can be derived from the commutation relation of the generator of SW symmetry with the generators of flavor symmetries

$$[\Delta_{\sigma}^{\text{SW}}, \Delta_{\Lambda}^G] = 0 . \quad (2.53)$$

This condition implies that  $b^I$  and  $\Gamma^A$  are covariant functions of the sources, namely

$$\begin{aligned} \Delta_{\Lambda}^G b^I &= i\Lambda^A (T_A b)^I \\ \Delta_{\Lambda}^G \Gamma^A &= i([\Lambda, \Gamma])^A . \end{aligned} \quad (2.54)$$

Another constraint can be derived from the commutation relation of the super-Weyl symmetry generator with itself

$$[\Delta_{\sigma_1}^{\text{SW}}, \Delta_{\sigma_2}^{\text{SW}}] = 0 . \quad (2.55)$$

Imposing the vanishing of terms proportional to  $\sigma_1 \bar{\sigma}_2 - \sigma_2 \bar{\sigma}_1$  in this commutator, we find the equation

$$\bar{\beta}^{\bar{I}} \partial_{\bar{I}} (e^V \Gamma) = \beta^I \partial_I (\bar{\Gamma} e^V) . \quad (2.56)$$

This constraint is used extensively in the computations described below. When written in components, this constraint agrees with the LRG result

$$\beta^I \rho_I^A + \bar{\beta}^{\bar{I}} \rho_{\bar{I}}^A = 0 . \quad (2.57)$$

## 2.8 Super-Weyl variation of functions

In this section we introduce some notations and terminology which facilitate the computation of Weyl variations of functions, and the analysis of the Weyl consistency conditions.

### 2.8.1 SUSY covariant derivative

To ensure the covariance of the super-Weyl anomaly under the global symmetries it is necessary to define a covariant supersymmetric derivative which involves the background gauge fields

$$\nabla^{\alpha} \lambda^I = \mathcal{D}^{\alpha} \lambda^I + (A^{\alpha})^I_J \lambda^J , \quad \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}} = \bar{\mathcal{D}}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}} + (B^{\dot{\alpha}})^{\bar{I}}_{\bar{J}} \bar{\lambda}^{\bar{J}} , \quad (2.58)$$

with  $\mathcal{D}^{\alpha}$  the standard chiral covariant derivative in the curved background, and we introduce the notation

$$(A^{\alpha})^I_J = (e^{-V})^{I\bar{K}} (\mathcal{D}^{\alpha} e^V)_{\bar{K}J} , \quad B^{\dot{\alpha}\bar{I}} = (\bar{\mathcal{D}}^{\dot{\alpha}} e^V)_{\bar{J}K} (e^{-V})^{K\bar{I}} . \quad (2.59)$$

The derivative  $\nabla^{\alpha}$  ( $\bar{\nabla}^{\dot{\alpha}}$ ) has no connection part including the  $\bar{I}$  ( $I$ ) indices; for example

$$(\nabla^{\alpha} \xi)_{K\bar{J}}^{I\bar{I}} = \mathcal{D}^{\alpha} \xi_{K\bar{J}}^{I\bar{I}} + A_S^{\alpha I} \xi_{K\bar{J}}^{S\bar{I}} + A_S^{\alpha \bar{J}} \xi_{K\bar{I}}^{S\bar{I}} - A_K^{\alpha S} \xi_{S\bar{J}}^{I\bar{I}} . \quad (2.60)$$



The chiral field strength associated with the background gauge fields is given by

$$(W^\alpha)_J^I = -\frac{1}{4}(\bar{\mathcal{D}}^2 - 4R)(A^\alpha)_J^I, \quad (\bar{W}^{\dot{\alpha}})_{\bar{J}}^{\bar{I}} = -\frac{1}{4}(\mathcal{D}^2 - 4\bar{R})(B^{\dot{\alpha}})_{\bar{J}}^{\bar{I}}. \quad (2.61)$$

Sometimes it is convenient to express  $(W^\alpha)_J^I$  in terms of adjoint indices

$$(W^\alpha)_J^I = (W^\alpha)^A (T_A)_J^I. \quad (2.62)$$

When writing in components we find the following useful result

$$\int d^2\theta \varphi^3 \text{Tr}[W^\alpha W_\alpha] \Big|_{\theta^2} + c.c. \supset -4\text{Tr}[F^{\mu\nu} F_{\mu\nu}], \quad (2.63)$$

where  $F^{\mu\nu}$  is the ‘‘canonical’’ field strength, see footnote 3.

A useful property satisfied by the covariant derivative is  $\nabla^\alpha (e^V)_{\bar{J}K} = \bar{\nabla}^{\dot{\alpha}} (e^V)^{K\bar{J}} = 0$ . One of the consequences of this identity is that for any arbitrary covariant function of the sources  $Y = Y(\lambda, \bar{\lambda}, e^V)$  we have

$$\nabla^\alpha Y = \partial_I Y \nabla^\alpha \lambda^I, \quad \bar{\nabla}^{\dot{\alpha}} Y = \partial_{\bar{I}} Y \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}}. \quad (2.64)$$

### 2.8.2 Lie derivative

In order to express the super-Weyl variation of covariant functions in parameter space in a compact form, it is convenient to define the following operators which we refer to as the Lie derivatives along the RG flow:

$$\begin{aligned} \mathcal{L}(Y_I) &= \beta^J \partial_J Y_I + \gamma_I^J Y_J & \tilde{\mathcal{L}}(Y_{\bar{I}}) &= \beta^{\bar{J}} \partial_{\bar{J}} Y_{\bar{I}} + \tilde{\gamma}_{\bar{I}}^{\bar{J}} Y_{\bar{J}} \\ \bar{\mathcal{L}}(Y_I) &= \bar{\beta}^{\bar{J}} \partial_{\bar{J}} Y_I + \tilde{\gamma}_I^{\bar{J}} Y_{\bar{J}} & \bar{\tilde{\mathcal{L}}}(Y_{\bar{I}}) &= \bar{\beta}^{\bar{J}} \partial_{\bar{J}} Y_{\bar{I}} + \bar{\tilde{\gamma}}_{\bar{I}}^{\bar{J}} Y_{\bar{J}} \end{aligned} \quad (2.65)$$

where  $Y_I$  is some arbitrary function of the sources, and the anomalous super-Weyl weight matrices  $\gamma$  and  $\tilde{\gamma}$  are defined in eq. (2.37). The generalization to tensors with more indices is given in the appendix. The Lie derivatives appear in the variation of functional involving the covariant derivative of couplings:

$$\begin{aligned} \Delta_\sigma^{\text{SW}}(Y_I \nabla^\alpha \lambda^I) &= \left(\frac{1}{2}\sigma - \bar{\sigma}\right) Y_I \nabla^\alpha \lambda^I + \mathcal{D}^\alpha \sigma Y_I \beta^I + \sigma \mathcal{L}(Y_I) \nabla^\alpha \lambda^I + \bar{\sigma} \bar{\mathcal{L}}(Y_I) \nabla^\alpha \lambda^I \\ \Delta_\sigma^{\text{SW}}(Y_{\bar{I}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}}) &= \left(\frac{1}{2}\bar{\sigma} - \sigma\right) Y_{\bar{I}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}} + \bar{\mathcal{D}}^{\dot{\alpha}} \bar{\sigma} Y_{\bar{I}} \bar{\beta}^{\bar{I}} + \sigma \bar{\mathcal{L}}(Y_{\bar{I}}) \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}} + \bar{\sigma} \bar{\tilde{\mathcal{L}}}(Y_{\bar{I}}) \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}} \end{aligned} \quad (2.66)$$

(The first term in each line corresponds to the classical super-Weyl variations of the covariant derivative).

### 2.8.3 Functions transforming covariantly under super-Weyl symmetry

The transformation of higher order covariant derivatives has a more complicated form, e.g.:

$$\begin{aligned} \Delta_\sigma^{\text{SW}}(Y_I \nabla^2 \lambda^I) &= (\sigma - 2\bar{\sigma}) Y_I \nabla^2 \lambda^I + \mathcal{D}^2 \sigma Y_K \beta^K + 2\mathcal{D}^\alpha \sigma Y_K (\delta_I^K + \partial_I \beta^K) \nabla_\alpha \lambda^I \\ &\quad + \sigma \mathcal{L}(Y_I) \nabla^2 \lambda^I + \sigma Y_K \partial_{IJ} \beta^K \nabla^\alpha \lambda^I \nabla_\alpha \lambda^J \\ &\quad + \bar{\sigma} \bar{\mathcal{L}}(Y_I) \nabla^2 \lambda^I + \bar{\sigma} Y_K \partial_{IJ} ((e^{-V} \bar{\Gamma} e^V)_L^K \lambda^L) \nabla^\alpha \lambda^I \nabla_\alpha \lambda^J \end{aligned} \quad (2.67)$$

As it was shown in the non-SUSY case [5], it is sometimes convenient to use the following functions of the sources

$$\begin{aligned}\Lambda^I &= (U^{-1})^I_K (\nabla^2 \lambda^K + 4\beta^K \bar{R}) \\ \Pi^{IJ} &= \nabla^\alpha \lambda^I \nabla_\alpha \lambda^J - \frac{1}{2}(\beta^I \Lambda^J + \beta^J \Lambda^I)\end{aligned}\tag{2.68}$$

with  $U_I^J = \delta_I^J + \partial_I \beta^J$ , in which some of the the derivatives of  $\sigma$  appearing in the SW variation cancel, leading to the following SW transformations:

$$\begin{aligned}\Delta_\sigma^{\text{SW}}(Y_I \Lambda^I) &= (\sigma - 2\bar{\sigma}) Y_I \Lambda^I + 2Y_I (\mathcal{D}^\alpha \sigma) \nabla_\alpha \lambda^I + \sigma \mathcal{L}(Y_I) \Lambda^I + \bar{\sigma} \bar{\mathcal{L}}(Y_I) \Lambda^I \\ &\quad + \sigma Y_K \gamma_{IJ}^K \Pi^{IJ} + \bar{\sigma} Y_K \gamma_{IJ}^K \Pi^{IJ} \\ \Delta_\sigma^{\text{SW}}(Y_{IJ} \Pi^{IJ}) &= (\sigma - 2\bar{\sigma}) Y_{IJ} \Pi^{IJ} + \sigma (\mathcal{L}(Y_{IJ}) - Y_{KL} \gamma_{IJ}^{KL}) \Pi^{IJ} + \bar{\sigma} (\bar{\mathcal{L}}(Y_{IJ}) - Y_{KL} \gamma_{IJ}^{KL}) \Pi^{IJ}\end{aligned}\tag{2.69}$$

The explicit expressions for the  $\gamma_{IJ}^K$  and  $\gamma_{IJ}^{KL}$  tensors are given in eq. (B.6).

### 3 Consistency conditions for the generalized super-Weyl anomaly

#### 3.1 The main idea

As explained above, the superspace local RG (SLRG) equation is an anomalous Ward identity for the super-Weyl (SW) symmetry off-criticality

$$\Delta_\sigma^{\text{SW}} \mathcal{W} = \mathcal{A}_\sigma^{\text{SW}}.\tag{3.1}$$

Now that we have defined and discussed the generator of the super-Weyl symmetry  $\Delta_\sigma^{\text{SW}}$ , it is time to move on to the right hand side of the equation and construct the anomaly. The anomaly encodes contact terms in the SW Ward identities for correlation functions, or equivalently, as explained in [5], it contains information about the log divergences in the effective action  $\mathcal{W}$  and the corresponding counterterms (see eq. (2.2)). The anomaly is parameterized by a set of functions which we refer to as the anomaly coefficients. These coefficients satisfy a non-trivial set of differential equations, due to the Wess-Zumino condition

$$\int d^8 z E^{-1} \delta_{\sigma_1, \sigma_2}^{\text{WZ}} \equiv \Delta_{\sigma_2}^{\text{SW}} \mathcal{A}_{\sigma_1}^{\text{SW}} - \Delta_{\sigma_1}^{\text{SW}} \mathcal{A}_{\sigma_2}^{\text{SW}} = 0.\tag{3.2}$$

The derivation of this set of equation is the main goal of this paper.

The procedure we follow consists of the following steps:

1. Write  $\mathcal{A}_\sigma^{\text{SW}}$  as the most general scalar constructed from the sources, invariant under the global symmetries, with the correct classical SW weight (in section 3.3 we show that in the presence of chiral anomalies it is also necessary to add some special terms which are not covariant under the global symmetries). In the supersymmetric case it is a non-trivial step to verify that one has the complete basis of terms in the anomaly.
2. The generating functional  $\mathcal{W}$  is defined up to local terms, therefore the anomaly function is not-unique. It is a crucial step to check how the different anomaly coefficients are modified under such changes of scheme, and identify the components of the anomaly which cannot be eliminated.

3. Compute the SW variation of the anomaly, and write eq. (3.2) in the following form

$$\delta_{\sigma_1, \sigma_2}^{WZ} = \sigma_{[1} \mathcal{D}^\alpha \sigma_{2]} \sum_a f_a^1 \mathcal{S}_\alpha^a + \sigma_{[1} \bar{\sigma}_{2]} \sum_a f_a^2 \mathcal{S}^a + \nabla^\alpha \sigma_{[1} \bar{\nabla}^{\dot{\alpha}} \bar{\sigma}_{2]} \sum_a f_a^3 \mathcal{S}_{\alpha\dot{\alpha}}^a + \nabla^2 \sigma_{[1} \bar{\nabla}^2 \bar{\sigma}_{2]} f^4 \quad (3.3)$$

where  $\mathcal{S}^a$  are an independent set of functions of space-time derivatives of the sources, (e.g.  $R\nabla_\alpha \lambda^i$ ) and  $f_a$  are functions of the anomaly coefficients and their derivatives with respect to the sources. The equations for the vanishing of each of the functions  $f_a$  are the Wess-Zumino consistency conditions.

4. In the supersymmetric case there is an additional subtlety, not appearing in the non-supersymmetric case, as a certain combination of terms in  $\delta_{\sigma_1, \sigma_2}^{WZ}$  can be written as a total derivative, and can therefore be removed from (or added to) the consistency conditions. This ambiguity in the consistency conditions is parameterized using a chiral superfield  $\Upsilon_A$ , and an unconstrained superfield  $\Omega_I$

$$\delta_{\sigma_1, \sigma_2}^{WZ} \sim \delta_{\sigma_1, \sigma_2}^{WZ} + F_1(\Upsilon_A) + F_2(\Omega_I) \quad (3.4)$$

where the following functions can be written as total derivatives

$$\begin{aligned} F_1(\Upsilon_A) &= \sigma_{[1} \mathcal{D}^\alpha \sigma_{2]} \Upsilon_A W_\alpha^A, \\ &= -\frac{1}{4} \bar{\mathcal{D}}^2 (\sigma_{[1} \mathcal{D}^\alpha \sigma_{2]} \Upsilon_A A_\alpha^A) \\ F_2(\Omega_I) &= \sigma_{[1} \mathcal{D}^\alpha \sigma_{2]} \left( \partial_{\bar{J}\bar{K}} \Omega_I \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{K}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \nabla_\alpha \lambda^I + \partial_{\bar{J}} \Omega_I \bar{\nabla}^2 \bar{\lambda}^{\bar{J}} \nabla_\alpha \lambda^I \right. \\ &\quad \left. + 4i \partial_{\bar{J}} \Omega_I \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \nabla_{\alpha\dot{\alpha}} \lambda^I - 4(W_\alpha \lambda)^I \Omega_I \right) \\ &= \bar{\mathcal{D}}^{\dot{\alpha}} (\sigma_{[1} \mathcal{D}^\alpha \sigma_{2]} \bar{\mathcal{D}}_{\dot{\alpha}} (\Omega_I \nabla_\alpha \lambda^I)) - \bar{\mathcal{D}}_{\dot{\alpha}} (\bar{\mathcal{D}}^{\dot{\alpha}} (\sigma_{[1} \mathcal{D}^\alpha \sigma_{2]})) (\Omega_I \nabla_\alpha \lambda^I). \end{aligned} \quad (3.5)$$

### 3.2 The super-Weyl anomaly

The anomaly can be found by writing the most general scalar function of the sources, allowed by symmetry and dimensional analysis,<sup>7</sup> and then imposing the consistency condition. In a supergravity background without the background fields  $\lambda$  and  $V$  one finds [31]

$$\begin{aligned} \mathcal{A}_\sigma^{\text{SW}}|_{\lambda=0, V=0} &= \int d^6 z \varphi^3 \sigma \kappa W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} + \text{c.c.} \\ &\quad - \int d^8 z E^{-1} (\sigma (2a(G^2 + 2R\bar{R}) + bR\bar{R}) + D^2 \sigma dR) + \text{c.c.} \end{aligned} \quad (3.6)$$

where  $W_{\alpha\beta\gamma}$ ,  $G_{\alpha\dot{\alpha}}$  and  $R$  are the supergravity multiplets (we use the notations of [22]; see [32] for the expression in Wess and Bagger conventions). Also,  $G^2 = G^a G_a = -\frac{1}{2} G^{\alpha\dot{\alpha}} G_{\alpha\dot{\alpha}}$ , and the coefficients  $\kappa$ ,  $a$ ,  $b$  and  $d$  are model dependent numbers<sup>8</sup> (the non-standard choice of notations for  $\kappa$  instead of  $2(c-a)$  will be justified below). In fact, one

<sup>7</sup>By dimensional analysis we mean that the anomaly terms must be classically invariant under a global rescaling.

<sup>8</sup>Our normalization is such that  $T_\mu^\mu = aE_4 - cW^2$ , e.g. for a chiral multiplet  $a = \frac{1}{12 \times 64\pi^2}$ ,  $c = \frac{1}{6 \times 64\pi^2}$ .

can use the Wess-Zumino consistency condition (3.2) to show that  $b$  must vanish at the fixed point. Moreover, as we show below, by adding a local term to  $\mathcal{W}$  one can set  $d$  to zero.

In the presence of the background sources we can write many more terms. We find that a relatively convenient basis is the following:

$$\begin{aligned}
\mathcal{A}_\sigma^{\text{SW}} &= \mathcal{A}_\sigma^{\text{SW}}|_{\lambda=0, V=0} + \mathcal{A}_\sigma^{\text{GW}} + \int d^6 z \varphi^3 \sigma \kappa_{AB} W^{A\alpha} W_\alpha^B + \text{c.c.} \\
&\quad - \int d^8 z E^{-1} \sigma \left( \chi_{IJ} G^{\alpha\dot{\alpha}} \nabla_\alpha \lambda^I \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \right. \\
&\quad + \xi_{[IJ]\bar{K}}^1 \nabla_{\alpha\dot{\alpha}} \lambda^I \nabla^\alpha \lambda^{\bar{J}} \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{K}} + \xi_{[I\bar{J}]K}^2 \nabla_{\alpha\dot{\alpha}} \bar{\lambda}^{\bar{I}} \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \nabla^\alpha \lambda^K \\
&\quad + \zeta_{AI}^1 W^{A\alpha} \nabla_\alpha \lambda^I + \zeta_{A\bar{I}}^2 \bar{W}^{\bar{A}} \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{I}} \\
&\quad + \epsilon_{I\bar{J}}^1 \Lambda^I \bar{\Lambda}^{\bar{J}} + \epsilon_{I\bar{J}\bar{K}}^2 \Pi^{IJ} \bar{\Lambda}^{\bar{K}} + \epsilon_{I\bar{J}K}^3 \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{I}} \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \Lambda^K + \epsilon_{I\bar{J}\bar{K}\bar{L}}^4 \Pi^{IJ} \bar{\Pi}^{\bar{K}\bar{L}} \\
&\quad + \eta_I^1 R \nabla^2 \lambda^I + \eta_{\bar{I}}^2 \bar{R} \bar{\nabla}^2 \bar{\lambda}^{\bar{I}} + \eta_{IJ}^3 R \nabla^\alpha \lambda^I \nabla_\alpha \lambda^J + \eta_{\bar{I}\bar{J}}^4 \bar{R} \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{I}} \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \Big) + \text{c.c.} \\
&\quad - \int d^8 z E^{-1} \mathcal{D}^\alpha \sigma \left( w_I^1 G_{\alpha\dot{\alpha}} \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{I}} + w_I^2 R \nabla_\alpha \lambda^I + u_{I\bar{J}}^1 \nabla_\alpha \lambda^I \Lambda^{\bar{J}} \right. \\
&\quad \left. + u_{I\bar{J}\bar{K}}^2 \nabla_\alpha \lambda^I \nabla_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{K}} \right) + \text{c.c.} \\
&\quad - \int d^8 z E^{-1} \mathcal{D}^2 \sigma \left( v_{\bar{I}}^1 \bar{\Lambda}^{\bar{I}} + v_{\bar{I}\bar{J}}^2 \bar{\Pi}^{\bar{I}\bar{J}} \right) + \text{c.c.} \tag{3.7}
\end{aligned}$$

$\Lambda$  and  $\Pi$  are the functions defined in (2.68), and  $\nabla_{\alpha\dot{\alpha}} = \frac{i}{2} \{ \nabla_\alpha, \bar{\nabla}_{\dot{\alpha}} \}$ . The term  $\mathcal{A}_\sigma^{\text{GW}}$  is a non-covariant anomaly term required by the consistency with chiral anomalies and will be discussed in the next section. Let us make some comments regarding this basis:

- In the presence of the background sources, the anomaly coefficients are generalized to be covariant functions of the sources (the covariance is implied by the fact that Weyl symmetry commutes with the global symmetries).
- There are terms which seem to be missing from this basis (such as  $\nabla_{\alpha\dot{\alpha}} \lambda \nabla^{\alpha\dot{\alpha}} \lambda$ ), but we have verified that all these terms can be integrated by parts and absorbed in the terms appearing in this formula. The approach we were following was to start with a basis where the derivatives are acting only on the sources (and not on  $\sigma$ ), and then integrating by parts when necessary.
- The matching between the superfield anomaly coefficients and the anomaly coefficients in the LRG anomalies is sometimes non-trivial. Each of the superWeyl anomalies, written in components, contains several anomaly terms with space-time derivatives of  $\sigma$  and the anomaly coefficients themselves.
- We chose to write some of the anomalies in the basis of the functions  $\Lambda$  and  $\Pi$ , in order to simplify the consistency conditions. It is useful, however, to identify the linear combination of anomaly coefficients which correspond to the  $\nabla^2 \lambda^I \bar{\nabla}^2 \bar{\lambda}^{\bar{J}}$  anomaly:

$$g_{IJ} \equiv (U^{-1})_I^K (\bar{U}^{-1})_{\bar{J}}^{\bar{L}} (\epsilon_{K\bar{L}}^1 - \epsilon_{KM\bar{L}}^2 \beta^M + \epsilon_{KM\bar{L}\bar{N}}^4 \beta^M \bar{\beta}^{\bar{N}}) \tag{3.8}$$

At the fixed point, the hermitean part of this matrix (specializing to components corresponding to exactly marginal primary operators), is proportional to the Zamolodchikov metric [33]

$$\mathcal{G}_{I\bar{J}} \equiv \langle \mathcal{O}_I(0) \bar{\mathcal{O}}_{\bar{J}}(z) \rangle (z^2)^3 \tag{3.9}$$

The relation between the metric and the super-Weyl anomaly coefficients is clarified in appendix D.

- We have not included in the anomaly terms such as

$$\int d^8z E^{-1} \sigma (\alpha R^2 + \alpha_I \bar{R} \nabla^2 \lambda^I + \alpha_{IJ} \nabla^2 \lambda^I \nabla^2 \lambda^J), \tag{3.10}$$

whose global Weyl variation is proportional to  $(\sigma - \bar{\sigma})$ . We checked by direct calculation that the consistency condition for these terms decouple from the ones for the other terms (which include the interesting ones which are related to the central charge  $a$ ). In this paper we will not discuss these terms and we will leave this as a topic for further investigation.

- There was a previous attempt [34] to write the Wess-Zumino consistency conditions for the anomaly in eq. (3.7), which missed many of the crucial ingredients, such as the background gauge fields.

### 3.3 The super-Weyl anomaly in the presence of chiral U(1) anomalies

In addition to the WZ consistency condition (3.2), the SW anomaly must satisfy another constraint

$$\Delta_{\Lambda}^G \mathcal{A}_{\sigma}^{\text{SW}} - \Delta_{\sigma}^{\text{SW}} \mathcal{A}_{\Lambda}^G = 0 \tag{3.11}$$

which is the WZ condition associated with the fact that the super-Weyl symmetry commutes with the global symmetries. This constraint was first discussed in the non-SUSY case in [21]. If the global symmetries are anomaly free ( $\mathcal{A}_{\Lambda}^G = 0$ ), then this constraint is satisfied if  $\mathcal{A}_{\sigma}^{\text{SW}}$  is invariant under the internal symmetries. If the theory does contain chiral anomalies, there is another consistent possibility in which the chiral anomaly  $\mathcal{A}_{\Lambda}^G$  is SW invariant (which is the case of the chiral-gravitational anomaly).

In general, however, the chiral anomaly is not SW invariant, due to the SW transformation of the background gauge fields defined by (2.47). This implies that the condition (3.11) can be satisfied only if we add terms to the Weyl anomaly which are not singlets of the anomalous flavor symmetry. An important observation is that  $\mathcal{A}_{\Lambda}^G$  is the *consistent* chiral anomaly, and not the *covariant* one. While in the case of anomalies involving only abelian currents, the consistent and covariant anomalies are identical (up to an overall numerical factor), in the non-abelian case the consistent anomaly does not have a closed local form, and is given only in terms of an integral over an auxiliary parameter (see e.g. [35–40]). In order to simplify the analysis here, we will focus on theories with only U(1) currents

involved in the chiral anomalies. We leave the computation of the Weyl anomaly in the presence of anomalous global non-abelian currents for future work.

The consistent abelian anomaly is given by

$$\mathcal{A}_\Lambda^G = \int d^6 z \varphi^3 i\Lambda^A \left( k_A W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} + k_{ABC} W^{\alpha B} W_\alpha^C \right) + c.c., \quad (3.12)$$

where  $W^{\alpha\beta\gamma}$  is the Weyl tensor,  $W^{\alpha A}$  is the field strength associated with the gauge field  $V^A$ , and the coefficients are given by

$$k_A \equiv -\frac{1}{192\pi^2} \text{Tr} [T_A] \quad k_{ABC} \equiv -\frac{1}{192\pi^2} \frac{1}{2} \text{Tr} [T_A \{T_B, T_C\}] . \quad (3.13)$$

$k_{ABC}$  is  $\frac{1}{3}$  times the coefficient of the covariant anomaly. The traces in (3.13) are taken on generators acting on the elementary fermionic field content of the theory. In the absence of holomorphic singlet functions, as discussed in section 2.2, the expressions for  $k_A$  and  $k_{ABC}$  given here are the only tensors with the right index structure that one can write. This can be viewed as a quick derivation of the Adler-Bardeen theorem [41] for supersymmetric theories which satisfy this condition.

The solution to the Wess-Zumino consistency condition (3.11) is found using the following non-trivial identity:

$$\Delta_\sigma^{\text{SW}} \mathcal{A}_\Lambda^G = -\Delta_\Lambda^G \left( \int d^8 z E^{-1} (\sigma + \bar{\sigma}) \Gamma^A X_A \right), \quad (3.14)$$

where  $X_A$  is a non-covariant function of the background gauge fields:

$$X_A = 2k_{ABC} \left( \nabla^\alpha (V^B W_\alpha^C) + \bar{\nabla}_{\dot{\alpha}} (V^B \bar{W}^{\dot{\alpha} C}) - \frac{1}{2} V^B (\nabla^\alpha W_\alpha^C + \bar{\nabla}_{\dot{\alpha}} \bar{W}^{\dot{\alpha} C}) \right). \quad (3.15)$$

$X^A$  is the superfield analog of the function found in [42], which gives the difference between the consistent and covariant abelian currents. This identity suggests that eq. (3.11) is satisfied only if we add the following term to the Weyl anomaly

$$\mathcal{A}_\sigma^W \supset \mathcal{A}_\sigma^{GW} \equiv - \int d^8 z E^{-1} (\sigma + \bar{\sigma}) \Gamma^A X_A . \quad (3.16)$$

This is similar to the solution found in [21], although we have to stress again, that we checked its validity in the SLRG equation only for anomalies involving abelian symmetries.

The anomaly term found here is not covariant, and can be rewritten as

$$\mathcal{A}_\sigma^{GW} = 2k_{ABC} \int d^8 z E^{-1} \sigma \left( \Gamma^C \nabla^\alpha V^A W_\alpha^B - \partial_i \Gamma^C V^A \bar{W}_{\dot{\alpha}}^B \bar{\nabla}^{\dot{\alpha} i} \right) + c.c. \quad (3.17)$$

The first term has the same form as the consistent anomaly (3.12), but since the function  $\Gamma$  is not holomorphic, this anomaly is not covariant.

A simple interpretation of the new anomaly and the new functions can be found by considering the Ward identity in the presence of the background sources

$$[\mathcal{T}] = -b^I [\mathcal{O}_I] - \frac{1}{4} \Gamma^A \bar{\nabla}^2 [J_A] - \frac{1}{4} \Gamma^A \bar{\nabla}^2 X_A + \dots \quad (3.18)$$

We see that the new anomaly can be absorbed by replacing the consistent current  $J_A$  with the covariant one  $\tilde{J}_A$

$$[\tilde{J}_A] \equiv [J_A] + X_A . \quad (3.19)$$

### 3.3.1 Chiral anomalies, ambiguity and Weyl anomaly coefficients

The appearance of the chiral anomalies implies that the ambiguity discussed in section 2.5, associated with a redefinition of the SW generator

$$\Delta_\sigma^{\text{SW}} \rightarrow (\Delta_\sigma^{\text{SW}})' = \Delta_\sigma^{\text{SW}} + \Delta_{\Lambda=ia\sigma}^G, \quad (3.20)$$

is valid only if we also shift the anomaly

$$\mathcal{A}_\sigma^{\text{SW}} \rightarrow (\mathcal{A}_\sigma^{\text{SW}})' = \mathcal{A}_\sigma^{\text{SW}} + \mathcal{A}_{\Lambda=ia\sigma}^G. \quad (3.21)$$

Under this transformation, the SW anomaly coefficient  $\kappa$  and  $\kappa_{AB}$  are shifted as<sup>9</sup>

$$\begin{aligned} \kappa &\rightarrow \kappa' = \kappa - k_A a^A \\ \kappa_{AB} &\rightarrow \kappa'_{AB} = \kappa_{AB} - 3k_{ABC} a^C \end{aligned} \quad (3.22)$$

We can therefore define non-ambiguous functions

$$\begin{aligned} \tau_{AB} &= \kappa_{AB} + 3k_{ABC} \Gamma^C \\ 2(c - a) &= \kappa + k_A \Gamma^A. \end{aligned} \quad (3.23)$$

These functions appear in the SW Ward identity after the currents are eliminated using a flavor Ward identity as in eq. (2.24) (now the covariant Ward identity has to be used):

$$\begin{aligned} [\mathcal{T}] &= -b^I [\mathcal{O}_I] - \frac{1}{4} \Gamma^A \bar{\nabla}^2 [\tilde{J}_A] + \kappa W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} + \kappa_{AB} W^{\alpha A} W_\alpha^B \\ &= -\beta^I [\mathcal{O}_I] + 2(c - a) W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} + \tau_{AB} W^{\alpha A} W_\alpha^B + \dots \end{aligned} \quad (3.24)$$

The expressions for  $\tau_{AB}$  and  $c - a$  are generalization of results appearing in [43], [44] and [45].

At the fixed point, the imaginary part of the  $\theta^2$  component of this equation gives the anomalous Ward identity of the  $R$ -symmetry. Using results from [43, 44], we find

$$\begin{aligned} 2(c - a)|_{f.p} &= -\frac{1}{128\pi^2} \text{Tr}[T_{R,f}] \\ \tau_{AB}|_{f.p} &= -\frac{3}{128\pi^2} \text{Tr}[T_{R,f} T_A T_B]. \end{aligned} \quad (3.25)$$

where  $T_{R,f}$  are the  $R$ -charges of the fermions and the trace is on the microscopic field content space.

The  $R$ -charge can be written in terms of  $\Gamma$  at the fixed point

$$T_{R,f} = \frac{2}{3} \Gamma^A T_A|_{f.p} - \frac{1}{3}. \quad (3.26)$$

---

<sup>9</sup>In order to find the shift in  $\kappa$  one has to first combine the  $\kappa$  anomaly and the first term in eq. (3.17) into one non-holomorphic anomaly coefficient  $\kappa_{AB} + 2d_{ABC} \Gamma^C$ , and take into account the transformation of  $\Gamma$ .

Combining this with eq. (3.23) and (3.25) we find that

$$\begin{aligned}\kappa|_{f.p} &= \frac{1}{384\pi^2} Tr[1] \\ \kappa_{AB}|_{f.p} &= \frac{1}{128\pi^2} Tr[T_A T_B],\end{aligned}\tag{3.27}$$

(assuming we use the parameterization in which  $T_R$  coincides with the superconformal  $R$  symmetry). In theories with no chiral singlets, these expressions are valid off-criticality as well, since there is no chiral function of the sources with the necessary index structure. This property will be important in the discussion appearing in section 4.1.2. Finally, we comment that this result agrees with an interpretation of  $\kappa_{AB}$  as the holomorphic  $\beta$  function for a weakly gauged global symmetry.

### 3.4 Scheme dependence of the anomaly

An important aspect of the anomaly which is relevant to our discussion is its dependence on the choice of scheme. By choice of scheme we refer here to the freedom to add local terms to the generating functional  $\mathcal{W}$ . The super-Weyl variation of these local terms modifies the anomaly coefficients. For example, the anomaly coefficients  $v^1, v^2$  and  $u^2$ , and the real part of  $d$  and  $u^1$  can be eliminated by adding the following local term

$$\delta\mathcal{W} = - \int d^8z E^{-1} \left( 4\bar{R} \left( dR + v_{\bar{I}}^1 \bar{\Lambda}^{\bar{I}} + v_{\bar{I}\bar{J}}^2 \bar{\Pi}^{\bar{I}\bar{J}} \right) - \frac{1}{2} \Lambda^I \left( u_{I\bar{J}}^1 \Lambda^{\bar{J}} + u_{I\bar{J}\bar{K}}^2 \nabla_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{K}} \right) \right) + c.c.\tag{3.28}$$

(Notice that other anomaly coefficients are modified in this process). In the following we assume that we work in such a scheme.

One can consider the effect of adding other counter terms. An important example is

$$\delta\mathcal{W} = - \int d^8z E^{-1} 2A(G^2 + 2R\bar{R})\tag{3.29}$$

with some arbitrary function  $A$ . Under this action, the anomaly coefficients are modified as

$$a \rightarrow a + \beta^I \partial_I A \quad w_{\bar{I}}^1 \rightarrow w_{\bar{I}}^1 - \partial_{\bar{I}} A \quad w_{\bar{I}}^2 \rightarrow w_{\bar{I}}^2 + \partial_{\bar{I}} A\tag{3.30}$$

We will later see, by inspecting the consistency conditions, that  $w_{\bar{I}}^1 = -\bar{w}_{\bar{I}}^2 \equiv w_{\bar{I}}$ . Using this result, we can define a function, which we denote by  $\tilde{a}$ , which is invariant under this transformation

$$\tilde{a} \equiv a + \bar{w}_I \beta^I.\tag{3.31}$$

This function will play an important role in our discussion. A useful observation is that  $\tilde{a}$  is sensitive to the choice of scheme related to the following counter term:

$$\delta\mathcal{W} = - \int d^8z E^{-1} C_{I\bar{J}} G^{\alpha\dot{\alpha}} \nabla_{\alpha} \lambda^I \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}}.\tag{3.32}$$

Indeed, under the addition of this term  $(a, \bar{w}_I, \tilde{a})$  transform as

$$a \rightarrow a, \quad \bar{w}_I \rightarrow C_{I\bar{J}} \bar{\beta}^{\bar{J}}, \quad \tilde{a} \rightarrow \tilde{a} + C_{I\bar{J}} \beta^I \bar{\beta}^{\bar{J}}.\tag{3.33}$$

We conclude that any contribution to  $\tilde{a}$  which is of order  $\beta\bar{\beta}$  can be eliminated by a choice of scheme.



### 3.5 Consistency conditions

We are now in the position to give the consistency conditions, which are derived from the constraint discussed in section 3.1. We will write the various constraints  $f^{1,2,3,4} = 0$ , associated with the various terms  $(\mathcal{S}^a, \mathcal{S}_\alpha^a, \mathcal{S}_{\alpha\dot{\alpha}}^a)$  as defined in eq. (3.3). All the equations are written in the scheme described in section 3.4 in which we eliminate as many terms as possible. In certain cases we write the equation already taking into account constraints obtained from previous equations.

#### 3.5.1 Constraints setting anomaly coefficients to zero

Some of the constraints enforce the vanishing of certain anomaly coefficients:

$(\sigma_{[1}\bar{\sigma}_{2]})\bar{\nabla}^2\nabla^2\lambda^I$	$\eta_I^1 = 0$	(3.34)
---	----------------	--------

$(\sigma_{[1}\mathcal{D}^\alpha\sigma_{2]})\nabla_\alpha\bar{\nabla}^2\bar{\lambda}^{\bar{I}}$	$\eta_{\bar{I}}^2 = 0$	(3.35)
--	------------------------	--------

$(\sigma_{[1}\bar{\sigma}_{2]})\nabla_{\alpha\dot{\alpha}}\lambda^I\nabla^{\alpha\dot{\alpha}}\lambda^J$	$\eta_{IJ}^3 = 0$	(3.36)
--	-------------------	--------

$(\sigma_{[1}\mathcal{D}^\alpha\sigma_{2]})\nabla^{\dot{\alpha}}\bar{\lambda}^{\bar{I}}\nabla_{\alpha\dot{\alpha}}\bar{\lambda}^{\bar{J}}$	$\eta_{\bar{I}\bar{J}}^4 = 0$	(3.37)
--	-------------------------------	--------

$(\sigma_{[1}\mathcal{D}^\alpha\sigma_{2]})\mathcal{D}_\alpha R$	$b = 0$	(3.38)
--	---------	--------

$(\nabla^2\sigma_{[1}\bar{\nabla}^2\bar{\sigma}_{2]})$	$d - \bar{d} = 0$	(3.39)
--	-------------------	--------

Another useful constraint we use at this stage is

$(\mathcal{D}^\alpha\sigma_{[1}\bar{\mathcal{D}}^{\dot{\alpha}}\bar{\sigma}_{2]})\nabla_{\alpha\dot{\alpha}}\lambda^I$	$w_I^2 = -\bar{w}_I^1$	(3.40)
--	------------------------	--------

Using this information and the choice of scheme described above, the anomaly now takes the form

$$\begin{aligned}
 \mathcal{A}_\sigma^{\text{SW}} = & \int d^6z \varphi^3 \sigma (\kappa(W_{\alpha\beta\gamma})^2) + \text{c.c.} \\
 & - \int d^8z E^{-1} \sigma \left( 2a(G^2 + 2R\bar{R}) + \chi_{I\bar{J}} G^{\alpha\dot{\alpha}} \nabla_\alpha \lambda^I \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \right. \\
 & + \xi_{[IJ]\bar{K}}^1 \nabla_{\alpha\dot{\alpha}} \lambda^I \nabla^{\alpha\dot{\alpha}} \lambda^J \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{K}} + \xi_{[\bar{I}\bar{J}]K}^2 \nabla_{\alpha\dot{\alpha}} \bar{\lambda}^{\bar{I}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \nabla^\alpha \lambda^K \\
 & + \zeta_{A\bar{I}}^1 W^{A\alpha} \nabla_\alpha \lambda^{\bar{I}} + (\zeta_{A\bar{I}}^2 + 2k_{ABC} \partial_{\bar{I}} \Gamma^B V^C) \bar{W}_{\dot{\alpha}}^A \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}} - (\kappa_{AB} + 2k_{ABC} \Gamma^C) W^{A\alpha} A_\alpha^B \\
 & + \epsilon_{I\bar{J}}^1 \Lambda^I \bar{\Lambda}^{\bar{J}} + \epsilon_{IJ\bar{K}}^2 \Pi^{IJ} \bar{\Lambda}^{\bar{K}} + \epsilon_{\bar{I}\bar{J}K}^3 \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{I}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \Lambda^K + \epsilon_{I\bar{J}\bar{K}\bar{L}}^4 \Pi^{IJ} \bar{\Pi}^{\bar{K}\bar{L}} \Big) + \text{c.c.} \\
 & - \int d^8z E^{-1} \mathcal{D}^\alpha \sigma \left( w_{\bar{I}} G_{\alpha\dot{\alpha}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}} + \bar{w}_I R \nabla_\alpha \lambda^I + u_{[I\bar{J}]} \nabla_\alpha \lambda^I \Lambda^{\bar{J}} \right) + \text{c.c.} \tag{3.41}
 \end{aligned}$$

where we use the notations  $w_{\bar{I}} \equiv w_{\bar{I}}^1 = -\bar{w}_{\bar{I}}^2$  and  $u_{[I\bar{J}]} \equiv \frac{1}{2}(u_{I\bar{J}}^1 - \bar{u}_{\bar{J}I}^1)$ .

#### 3.5.2 Constraints which can be used to algebraically eliminate anomaly coefficients

The following constraints can be used to write some anomaly coefficients as an algebraic function of the others:

$(\sigma_{[1}\mathcal{D}^\alpha\sigma_{2]})\nabla_\alpha\lambda^I\bar{\Lambda}^{\bar{J}}$	$8\epsilon_{I\bar{J}}^1 = \bar{U}_{\bar{J}}^{\bar{K}}\left(-\chi_{I\bar{K}} + 4\mathcal{L}(u_{[I\bar{L}]})\bar{U}^{-1})_{\bar{K}}^{\bar{L}}\right)$	(3.42)
$(\sigma_{[1}\bar{\sigma}_{2]})\nabla_{\alpha\dot{\alpha}}\lambda^{(I}\nabla^\alpha\lambda^{J)}\bar{\nabla}^{\dot{\alpha}}\bar{\lambda}^{\bar{K}}$	$8\epsilon_{(IJ)\bar{K}}^2 = -\partial_{(J}\chi_{I)\bar{K}} + \mathcal{I}_{(IJ)\bar{K}}^0 + \mathcal{I}_{(I\bar{K}A}^1\partial_J)\Gamma^A$	(3.43)
$(\sigma_{[1}\mathcal{D}^\alpha\sigma_{2]})\nabla_\alpha\lambda^K\bar{\nabla}_{\dot{\alpha}}\bar{\lambda}^{\bar{I}}\bar{\nabla}^{\dot{\alpha}}\bar{\lambda}^{\bar{J}}$	$8\epsilon_{(\bar{I}\bar{J})K}^3 = -\partial_{(J}\chi_{K\bar{I})} + \bar{\mathcal{I}}_{(\bar{I}\bar{J})K}^0 + \mathcal{I}_{K(\bar{I}A}^1\partial_{\bar{J})}\Gamma^A$	(3.44)
$(\mathcal{D}^\alpha\sigma_{[1}\bar{\mathcal{D}}^{\dot{\alpha}}\bar{\sigma}_{2]})\nabla_\alpha\lambda^I\bar{\nabla}_{\dot{\alpha}}\bar{\lambda}^{\bar{J}}$	$8u_{[I\bar{J}]} = \partial_{\bar{J}}\bar{w}_I - \partial_I w_{\bar{J}} + i\xi_{[\bar{K}\bar{J}]I}^2\bar{\beta}^{\bar{K}} + i\bar{\xi}_{[KI]\bar{J}}^2\beta^K$	(3.45)

where:

$$\begin{aligned}
 \mathcal{I}_{(IJ)\bar{K}}^0 &\equiv 4u_{[\bar{K}L]}\gamma_{IJ}^L + \bar{\zeta}_{A(I}\partial_{J)\bar{K}}\Gamma^A \\
 \mathcal{I}_{I\bar{J}A}^1 &\equiv 2i\xi_{[KI]\bar{J}}^1(T_A\lambda)^K + 2i\xi_{[\bar{K}\bar{J}]I}^2(e^V T_A e^{-V}\bar{\lambda})^{\bar{K}} \\
 &\quad - \partial_{\bar{J}}\zeta_{IA}^1 - \partial_I\zeta_{\bar{J}\bar{K}}^2(e^V T_A e^{-V})_{\bar{L}}^{\bar{K}}.
 \end{aligned}
 \tag{3.46}$$

and the symmetrization in eqs. (3.43), (3.44) is just on the  $(I, J)$  and  $(\bar{I}, \bar{J})$  indices. The symbol  $\gamma_{IJ}^K$  defined in (B.6) and  $\zeta_{\bar{J}\bar{K}}^{2\bar{L}}$  is defined such that  $\zeta_{\bar{J}A}^2\bar{W}_{\dot{\alpha}}^A = \zeta_{\bar{J}\bar{K}}^{2\bar{L}}(\bar{W}_{\dot{\alpha}})_{\bar{L}}^{\bar{K}}$ .

We are now left with the following independent anomaly coefficients:

$$a, \kappa, \kappa_{AB}, \chi_{I\bar{J}}, w_{\bar{I}}, \xi_{I\bar{J}\bar{K}}^1, \xi_{\bar{I}\bar{J}K}^2, \zeta_{AI}^1, \zeta_{A\bar{I}}^2, \epsilon_{I\bar{J}\bar{K}\bar{L}}^4
 \tag{3.47}$$

### 3.5.3 Reality constraints

Some of the consistency condition take the form of reality constraints on some combinations of the anomaly coefficients:

$(\mathcal{D}^\alpha\sigma_{[1}\bar{\mathcal{D}}^{\dot{\alpha}}\bar{\sigma}_{2]})G_{\alpha\dot{\alpha}}$	$\tilde{a} = \bar{a}$	(3.48)
$(\sigma_{[1}\bar{\sigma}_{2]})G^{\alpha\dot{\alpha}}G_{\alpha\dot{\alpha}}$	$\bar{\mathcal{L}}(a) = \mathcal{L}(\bar{a})$	(3.49)
$(\sigma_{[1}\bar{\sigma}_{2]})G_{\alpha\dot{\alpha}}\nabla^\alpha\lambda^i\bar{\nabla}^{\dot{\alpha}}\bar{\lambda}^{\bar{j}}$	$\bar{\mathcal{L}}(\chi_{I\bar{J}}) = \mathcal{L}(\bar{\chi}_{\bar{J}I})$	(3.50)
$(\sigma_{[1}\bar{\sigma}_{2]})\Lambda^I\bar{\Lambda}^{\bar{J}}$	$\bar{\mathcal{L}}(g_{I\bar{J}}) = \mathcal{L}(\bar{g}_{\bar{J}I})$	(3.51)
$(\sigma_{[1}\bar{\sigma}_{2]})\nabla^\alpha W_\alpha^A$	$(e^V)_{\bar{J}K}(\bar{w}_I\lambda^K + \beta^L\bar{\zeta}_{LI}^{2K}) = h.c.$	(3.52)
$(\sigma_{[1}\bar{\sigma}_{2]})\nabla^{\alpha\dot{\alpha}}\lambda^I\nabla_{\alpha\dot{\alpha}}\bar{\lambda}^{\bar{J}}$	$\chi_{I\bar{J}} - \partial_I w_{\bar{J}} - 2i\xi_{[\bar{J}\bar{K}]I}^2\bar{\beta}^{\bar{K}} + \zeta_{A\bar{J}}^2\partial_I\bar{\Gamma}^A = h.c.$	(3.53)
$(\sigma_{[1}\bar{\sigma}_{2]})\nabla^\alpha\lambda^I\nabla_\alpha\lambda^J\bar{\nabla}_{\dot{\alpha}}\bar{\lambda}^{\bar{K}}\bar{\nabla}^{\dot{\alpha}}\bar{\lambda}^{\bar{L}}$	$\bar{\mathcal{L}}(\epsilon_{I\bar{J}\bar{K}\bar{L}}^4) - \epsilon_{MN\bar{K}\bar{L}}^4\gamma_{IJ}^{MN} - \epsilon_{I\bar{J}M\bar{N}}^4\bar{\gamma}_{\bar{K}\bar{L}}^{\bar{M}\bar{N}} + \epsilon_{I\bar{J}M}^2\bar{\gamma}_{\bar{K}\bar{L}}^{\bar{M}}$ $+ \epsilon_{I\bar{J}M}^3\gamma_{\bar{K}\bar{L}}^M - \frac{1}{4}\mathcal{I}_{I\bar{K}A}^1\partial_{\bar{J}\bar{L}}\Gamma^A = h.c.$	(3.54)

where the quantities  $\tilde{a}$  and  $g_{I\bar{J}}$  are defined in eq. (3.31) and eq. (3.8).

### 3.5.4 Other constraints

The following consistency conditions will be further discussed in section 4; they give constraints on the  $a$ -function and on the conformal manifold:

$(\sigma_{[1}\mathcal{D}^\alpha\sigma_{2]})\bar{\nabla}^{\dot{\alpha}}\lambda^{\bar{I}}G_{\alpha\dot{\alpha}}$	$\partial_{\bar{I}}a = \beta^J\chi_{J\bar{I}} - \mathcal{L}(w_{\bar{I}})$	(3.55)
--	---	--------

$(\sigma_{[1}\mathcal{D}^\alpha\sigma_{2]})W_\alpha^A$	$-2\kappa_{AB}\Gamma^B - 3k_{ABC}\Gamma^B\Gamma^C + \zeta_{AI}^1\beta^I + w_{\bar{I}}(e^V T_A e^{-V}\bar{\lambda})^{\bar{I}}$ $= \Upsilon_A + \Omega_I(T_A\lambda)^I$	(3.56)
--	--	--------

$(\sigma_{[1}\mathcal{D}^\alpha\sigma_{2]})\nabla_{\alpha\dot{\alpha}}\lambda^{\bar{I}}\bar{\nabla}^{\dot{\alpha}}\bar{\lambda}^{\bar{J}}$	$\chi_{I\bar{J}} - \partial_I w_{\bar{J}} - 2i\xi_{[IK]\bar{J}}^1\beta^K + \zeta_{AI}^1\partial_{\bar{J}}\Gamma^A = -\partial_{\bar{J}}\Omega_I$	(3.57)
--	--	--------

$(\sigma_{[1}\mathcal{D}^\alpha\sigma_{2]})\nabla_{\alpha\dot{\alpha}}\bar{\lambda}^{\bar{I}}\bar{\nabla}^{\dot{\alpha}}\bar{\lambda}^{\bar{J}}$	$\partial_{\bar{I}}w_{\bar{J}} - \partial_{\bar{J}}w_{\bar{I}} = 2i\xi_{[\bar{I}\bar{J}]K}^2\beta^K + (\zeta_{A\bar{J}}^2\partial_{\bar{I}}\bar{\Gamma}^A - \zeta_{A\bar{I}}^2\partial_{\bar{J}}\bar{\Gamma}^A)$	(3.58)
--	--	--------

Combining (3.55) with other constraints eqs. (3.48), (3.52), (3.53), (3.58), one finds a similar equation

$$\partial_I a = \bar{\beta}^{\bar{J}}\chi_{I\bar{J}} - \mathcal{L}(\bar{w}_I). \quad (3.59)$$

The same equation is found in a completely different way from the  $(\sigma_{[1}\bar{\sigma}_{2]})G_{\alpha\dot{\alpha}}\nabla^{\alpha\dot{\alpha}}\lambda^I$  consistency condition; this is a non-trivial cross-check of our calculations. An interesting property of eq. (3.56) is that it seems to be sensitive to the ambiguity discussed in eqs. (2.44), (2.45). However, the invariance of the equation can be restored by assigning  $\Upsilon_A$  appropriate transformation properties:

$$\Upsilon'_A = \Upsilon_A - 2\kappa_{AB}a^B - 3\kappa_{ABC}a^B a^C. \quad (3.60)$$

Finally, we list the set of remaining constraints, which we will not use in the discussion of the physical implication:

$(\sigma_{[1}\bar{\sigma}_{2]})\nabla_{\alpha\dot{\alpha}}\bar{\lambda}^{[\bar{I}}\bar{\nabla}^{\dot{\alpha}}\bar{\lambda}^{\bar{J}]}\nabla^\alpha\lambda^K$	$\partial_{[J}\chi_{I]\bar{K}} = i\mathcal{L}(\bar{\xi}_{[IJ]\bar{K}}^2) + \mathcal{I}_{IJ\bar{K}\bar{L}}^2\bar{\beta}^{\bar{L}} + \mathcal{I}_{[J\bar{K}A}^1\partial_I\Gamma^A - \bar{\xi}_{A[I}\partial_{J]\bar{K}}\Gamma^A$	(3.61)
--	--	--------

$(\sigma_{[1}\bar{\sigma}_{2]})W^{\alpha A}\nabla_\alpha\lambda^I$	$\chi_{I\bar{J}}(e^V T_A e^{-V}\bar{\lambda})^{\bar{J}} = \mathcal{L}(\bar{\xi}_{AI}^2) + \mathcal{I}_{IJA}^1\bar{\beta}^{\bar{J}} - 2\tau_{AB}\partial_I\Gamma^B + \mathcal{I}_{AI}^3$	(3.62)
--	---	--------

$(\sigma_{[1}\bar{\sigma}_{2]})\nabla^\alpha\lambda^I\nabla_\alpha\bar{\nabla}^2\bar{\lambda}^{\bar{J}}$	$8(\epsilon_{IJ}^1 - \beta^K\epsilon_{IKJ}^2 + \bar{\beta}^{\bar{K}}\epsilon_{\bar{J}\bar{K}I}^3)$ $= U_I^K (-\chi_{K\bar{J}} + 4\mathcal{L}(u_{[L\bar{J}]}(U^{-1})^L_K))$	(3.63)
--	---	--------

$(\sigma_{[1}\bar{\sigma}_{2]})\nabla^\alpha\lambda^I\nabla_\alpha\lambda^J\bar{\Lambda}^{\bar{K}}$	$0 = \gamma_{IJ}^L(\epsilon_{L\bar{K}}^1 - \bar{\epsilon}_{\bar{K}L}^1) + \bar{\mathcal{L}}(\epsilon_{IJ\bar{K}}^2) - \epsilon_{LM\bar{K}}^2\gamma_{IJ}^{LM}$ $- \mathcal{L}(\bar{\epsilon}_{IJ\bar{K}}^3) + \bar{\beta}^{\bar{L}}\bar{\epsilon}_{\bar{K}LM}^3\gamma_{IJ}^M$ $- \frac{1}{4}\mathcal{I}_{I\bar{K}A}^1\partial_{J\bar{L}}\bar{\Gamma}^A\bar{\beta}^{\bar{L}} + \frac{1}{4}\bar{\mathcal{I}}_{\bar{K}IA}^1\partial_{L\bar{J}}\Gamma^A\bar{\beta}^{\bar{L}}$	(3.64)
---	--	--------

where  $\mathcal{I}^1$  and  $\tau_{AB}$  were introduced in eq. (3.46) and (3.23), and we also defined

$$\begin{aligned} \mathcal{I}_{IJ\bar{K}\bar{L}}^2 &\equiv i\partial_{\bar{L}}\xi_{[IJ]\bar{K}}^1 - i\partial_{\bar{K}}\xi_{[IJ]\bar{L}}^1 + i\partial_J\xi_{[\bar{K}\bar{L}]I}^2 - i\partial_I\xi_{[\bar{K}\bar{L}]J}^2 \\ \mathcal{I}_{AI}^3 &\equiv -\partial_I\bar{\Gamma}^B \left( \zeta_{BI}^2(e^V T_A e^{-V}\bar{\lambda})^{\bar{I}} + \zeta_{\bar{K}\bar{N}}^{2\bar{M}}(e^{-V} T_A e^V)^{\bar{N}}_{\bar{M}}(\bar{T}_B\bar{\lambda})^{\bar{K}} \right) \\ &\quad - \partial_I\Gamma^B \left( \zeta_{BK}^1(T_A\lambda)^K + \zeta_{AK}^1(T_B\lambda)^K \right). \end{aligned} \quad (3.65)$$

The Lie derivative of objects containing a global symmetry index such as  $\mathcal{L}(\bar{\zeta}_{AI}^2)$  is defined in eq. (B.3).

### 3.5.5 Comments on the consistency conditions

We have presented the set of consistency conditions for the SLRG anomaly. In the original formulation of the LRG equation, there are just three consistency condition which can not be used to eliminate some anomaly coefficients as an algebraic function of the others (see eqs. (2.66) and (2.68) of [5]); these three equations are similar in form to eqs. (3.55), (3.62), and (3.52); in the superspace formalism we find many more consistency conditions. This is to be expected, as the anomaly discussed here is related to the super-Weyl symmetry, which is a larger symmetry than the one considered in the LRG equation. Moreover, one has to bare in mind that the SLRG anomaly coefficients are not in one-to one correspondence with the ones of the LRG equation. Another crucial difference is appearance of the unknown functions  $\Upsilon$  and  $\Omega$ , which makes the equations less constraining.

Finally, we mention that the equations associated with the following terms in  $\delta_{\sigma_1, \sigma_2}^{WZ}$  are vanishing by imposing the previous constraints, thus providing a highly non-trivial consistency check for our set of equations:

$$(\sigma_{[1}\mathcal{D}^\alpha\sigma_{2]})R\nabla_\alpha\lambda^i, (\sigma_{[1}\bar{\sigma}_{2]})\bar{R}R, (\sigma_{[1}\bar{\sigma}_{2]})R\nabla^2\lambda^i, (\sigma_{[1}\bar{\sigma}_{2]})R\nabla^\alpha\lambda^i\nabla_\alpha\lambda^j, (\sigma_{[1}\bar{\sigma}_{2]})\bar{D}^2\bar{R},$$

$$(\sigma_{[1}\bar{\sigma}_{2]})\nabla^\alpha\lambda^i\nabla_\alpha R, (\sigma_{[1}\bar{\sigma}_{2]})\nabla^{\alpha\dot{\alpha}}\lambda^i G_{\alpha\dot{\alpha}}.$$

## 4 General implications

### 4.1 $a$ -function and $a$ -maximization

The anomaly coefficient  $a$  at the fixed point is related to the irreversibility of RG flow (the  $a$ -theorem) [2, 3, 46–48] and gives a constraint on the number of degrees of freedom which can emerge in the IR from a given UV description. In the case of supersymmetric theories  $a$  can be used to determine the superconformal R-charge at criticality and the anomalous dimension of chiral primaries using  $a$ -maximization [13]. In the context of the local RG equation, the function  $\tilde{a}$  is a continuation of the anomaly coefficient  $a$  off criticality. As was shown in [2, 3], away from conformal fixed point the quantity  $\tilde{a}$  is ambiguous up to terms which are quadratic in the beta functions (see eq. (3.33)). Moreover, it is shown that at leading order  $\tilde{a}$  is monotonically decreasing along the RG flow; this feature, if valid non-perturbatively (perhaps just in some specific scheme), would correspond to a strong version of the  $a$ -theorem, in analogy with the two-dimensional case [33].

In this section we will study the constraints on  $\tilde{a}$  in the superspace formulation of the local RG equation. This quantity was already defined in eq. (3.31) and was shown to be real by the consistency condition eq. (3.48). We will also discuss  $a$ -maximization, and its generalization off-criticality, proving a conjectured exact formula for this quantity.

In the superspace formulation,  $\tilde{a}$  satisfies

$$\partial_I\tilde{a} = \bar{\beta}^{\bar{J}}\chi_{I\bar{J}} - \beta^J(\partial_J\bar{w}_I - \partial_I\bar{w}_J), \tag{4.1}$$

which follows combining eqs. (3.59), (3.48). This equation has a direct analog also in the generic non-supersymmetric case and gives non-trivial relations between  $\beta$ -functions (see, e.g. [6]); it is mostly famous for providing a proof for the irreversibility of the RG flows in perturbation theory [2, 3]. Indeed, by multiplying by  $\beta$ -functions, and using the reality condition (3.48) we find

$$\mu \frac{d}{d\mu} \tilde{a} \equiv \left( \beta^I \partial_I + \bar{\beta}^{\bar{J}} \partial_{\bar{J}} \right) \tilde{a} = \beta^I \bar{\beta}^{\bar{J}} (\chi_{I\bar{J}} + \bar{\chi}_{\bar{J}I}) \quad (4.2)$$

At leading order in  $\beta$  and  $\partial\beta$  we can use eqs. (3.9), (3.42) to show that

$$\chi_{I\bar{J}} + \bar{\chi}_{\bar{J}I} \propto \mathcal{G}_{I\bar{J}} + O(\beta, \partial\beta) \quad (4.3)$$

where  $\mathcal{G}$  is the positive definite matrix defined in 3.9. We conclude that the r.h.s. of (4.2) is positive definite, and  $\tilde{a}$  is therefore a function of the couplings which changes monotonously along the RG flow (at leading order), thus excluding the possibility of cyclic flows [30, 49, 50]. As was demonstrated in [5], there exists a scheme in which  $\tilde{a}$  coincides with a quantity related to the on-shell dilaton amplitude, which shares the same property.

The quantity  $\Xi_{JI} = \partial_J \bar{w}_I - \partial_I \bar{w}_J$  was found to vanish in some appropriate scheme<sup>10</sup> in every example, to all orders checked. If one could prove, in addition to the positivity of  $\chi_{I\bar{J}}$ , that  $\Xi_{JI}$  vanishes one would establish the gradient flow property for  $\tilde{a}$  (see eq. (4.1)). The consistency condition in eq. (3.58) can be used to re-express  $\Xi_{JI}$  in term of  $\xi^2$  and  $\zeta^2$  anomaly coefficients, but does not give us any argument supporting the conjecture that  $\Xi_{JI} = 0$  outside the fixed point.

#### 4.1.1 *a*-maximization

In this section we assume the theory can be formulated in term of some elementary field description, which corresponds to assuming that the theory is asymptotically free and can be thought as in the UV as a deformation of a free field theory. Also, in this section the lower-case indices  $i, j$  are running on the elementary field content of the theory.

The idea behind the “*a*-maximization” prescription is the following: the superconformal  $R$  symmetry is a useful tool in the study of RG flows, because the  $R$  charge of a field determines also its scaling dimension

$$R_{f,i} = \frac{2\gamma_i - 1}{3} \quad (4.4)$$

where  $R_{f,i}$  is the  $R$ -charge of the fermion in the superfield  $\Phi_i$  and  $\gamma_i$  is the corresponding anomalous dimension, which is assumed to be diagonalized. We can write (after setting the couplings to their background values, and diagonalizing the matrix  $\Gamma^A T_A$ ):

$$\gamma_i^j = \Gamma^A (T_A)_i^j, \quad (4.5)$$

where the matrix  $(T_A)_i^j$  is acting on the elementary field content.

If in a theory there are some unbroken not- $R$   $U(1)_A$  symmetries, the candidate  $R$ -symmetry is ambiguous, because we can generate another  $R$ -symmetry by shifting the

---

<sup>10</sup>Due to the transformation law in eq. (3.33), the relation  $\Xi_{JI} = 0$  can not be valid in all the schemes.

original one by some linear combination of the  $U(1)_{AS}$ . The correct superconformal R-symmetry can be fixed using the property that at the fixed point the anomaly coefficient of each  $U(1)_A \times \text{Gravity}^2$  and  $U(1)_A \times U(1)_R^2$  are equal<sup>11</sup> [13]; this is equivalent to  $a$ -extremization, where the central charge  $a$  is calculated from the fermionic R-charges using [43, 44].

Concretely, the following equation which must be satisfied by each of the unbroken  $U(1)$  symmetries:

$$9\text{Tr}[R_f^2 T_A] - \text{Tr}[T_A] = 0, \tag{4.6}$$

where again the trace is in the space of the elementary fields. Writing this equation using (4.4) we find the constraint

$$\text{Tr}[T_A T_B] \Gamma^B - \text{Tr}[T_A T_B T_C] \Gamma^B \Gamma^C = 0. \tag{4.7}$$

Let us compare this equation with the consistency condition (3.56), evaluated at the fixed point, for a  $U(1)$  symmetry which is unbroken by the background spurion:<sup>12</sup>

$$2\kappa_{AB} \Gamma^B + 3k_{ABC} \Gamma^B \Gamma^C = -\Upsilon_A. \tag{4.8}$$

Using eqs. (3.13), (3.27) we find that the  $a$ -maximization equation corresponds to this consistency condition when  $\Upsilon_A$  vanishes. Indeed, under the assumption that there are no chiral singlet functions,  $\Upsilon_A$  is a constant and it can always be set to zero using the reparameterization described in section 2.5, see eq. (3.60). In an expansion nearby a free fixed point, the parameterization for which  $\Upsilon_A = 0$  coincides with the one usually used in perturbative calculation (in which the eigenvalues of  $\Gamma$  are the anomalous dimensions of elementary fields).

We stress that eq. (3.56) generalizes  $a$ -maximization in two ways, because it is valid also outside the conformal fixed point and for symmetries which are explicitly broken by some coupling. Setting  $\Upsilon_A = 0$  in eq. (3.56) we get:

$$2\kappa_{AB} \Gamma^B + 3k_{ABC} \Gamma^B \Gamma^C - \zeta_{AI}^1 \beta^I - w_{\bar{I}} (e^V T_A e^{-V \bar{\lambda}})^{\bar{I}} = -\Omega_I (T_A \lambda)^I. \tag{4.9}$$

$\Omega_I$  is not an anomaly coefficient, but some arbitrary function which appears also (and only) in eq. (3.57):

$$\chi_{I\bar{J}} - \partial_I w_{\bar{J}} - 2i \zeta_{[IK] \bar{J}}^1 \beta^K + \zeta_{AI}^1 \partial_{\bar{J}} \Gamma^A = -\partial_{\bar{J}} \Omega_I. \tag{4.10}$$

This equation can be used to relate  $\chi$  to other anomaly coefficients (this is possible just modulo the function  $\Omega_I$ , which is not directly an anomaly coefficient). These two equations have no analog in the case without supersymmetry.

---

<sup>11</sup>There is a potential loophole due to possible IR accidental symmetries which may mix with the  $U(1)_R$ .

<sup>12</sup>Recall that we are considering chiral anomalies with abelian symmetries only, therefore all the indices in this equation correspond to  $U(1)$  symmetries.

### 4.1.2 $\tilde{a}$ off-criticality

We can now start with the differential equation for  $a$  eq. (3.55) and use eq. (3.57) to eliminate  $\chi$  and combine with (3.56). We finally find the following relation:

$$\partial_{\tilde{I}} \left( \tilde{a} + \Upsilon_A \Gamma^A + \kappa_{AB} \Gamma^A \Gamma^B + k_{ABC} \Gamma^A \Gamma^B \Gamma^C + \beta^I \tilde{\Omega}_I \right) = 0 \quad (4.11)$$

where we introduced the notation

$$\tilde{\Omega}_I \equiv \Omega_I - \bar{w}_I . \quad (4.12)$$

This result implies that the expression in parentheses is a function of the chiral sources, which is invariant under all global symmetries of the theory. In the absence of chiral singlets, as discussed in section 2.2, such a function must be a constant. Working in the parameterization with vanishing  $\Upsilon$ , we conclude that  $\tilde{a}$  has the following form

$$\tilde{a} = -\kappa_{AB} \Gamma^A \Gamma^B - k_{ABC} \Gamma^A \Gamma^B \Gamma^C - \beta^I \tilde{\Omega}_I + \text{const.} \quad (4.13)$$

Under the same assumption, the chiral functions  $\kappa_{AB}$  and  $k_{ABC}$  must be constant, and eqs. (3.13), (3.27) can be used off-criticality. We find the expression for  $\tilde{a}$  in terms of the anomalous dimensions matrices, as advertised in the introduction:

$$\tilde{a} = -\frac{1}{128\pi^2} \text{Tr}[\gamma^2] + \frac{1}{192\pi^2} \text{Tr}[\gamma^3] - \beta^I \tilde{\Omega}_I + \text{const.} \quad (4.14)$$

At the conformal fixed point,  $\tilde{a}$  coincides with the central charge  $a$  and eq. (4.14) coincides with the expression found in [43, 44]. Out of criticality, eq. (4.14) was conjectured and tested in [6, 10, 11, 14–16, 51].<sup>13</sup> Moreover, the same line of argument lead to an interesting conjecture regarding the structure of anomalous dimensions in renormalizable supersymmetric field theories (see eq. (4.17) of [11] which is an extension of an earlier equation appearing in [6]). Our equation (3.56) looks like a generalization of this conjectured formula. The advantage of our method is that these results are derived as consistency conditions, rather than conjectured, and are valid off-criticality by definition. The downside is that in the presence of chiral singlets, the most we can get is eq. (4.11).

As a last comment, we recall that any dependence of  $\tilde{\Omega}$  on the  $\beta$ -function can be removed by a choice of scheme (see discussion in section 3.4).

## 4.2 The conformal manifold

When we perturb a conformal field theory with a marginal operator, the deformation preserve conformality just at zero order because in general the deforming operator dimension itself gets corrections: we can further classify such perturbations into marginally relevant, marginally irrelevant and exactly marginal. One may expect that the case of exactly marginal deformations is fine-tuned in absence of special symmetries; indeed the only known examples of such deformations in  $d = 4$  are realized in supersymmetric theories [17, 19, 20, 52, 53].

---

<sup>13</sup>In [14–16] the term proportional to  $\beta$  in (4.14) was introduced as a Lagrange multiplier, whose purpose was to ensure the vanishing of the  $\beta$  function at the fixed point.

In general one can have multiple exactly marginal deformation which parameterize a continuous family of conformal field theories. Assuming that there are no singularities, one can locally think of this family of CFTs as a submanifold of the space of all the couplings of the theory. Such objects are referred as Conformal Manifolds (CM), and appear whenever the number of independent conditions, which are necessary for the vanishing of all the  $\beta$ -functions is less than the number of coupling constants in the theory [17]. Nearby a given  $\mathcal{N} = 1$  SCFT, the conformal manifold can be built as the quotient of all the marginal deformation of the theory divided by the complexified symmetry group [20]. This explains why the existence of a CM is a rather common property in the case of supersymmetric theories.

In this section we will apply the superspace formulation of the local RG equation for the study of supersymmetric CM. Along the CM all the  $\beta$ -functions  $\beta^I = 0$ ; also, we can always choose the coordinates in the space of the couplings in such a way that nearby the origin  $\lambda^I = 0$  the conformal manifold directions are tangent to some group of coordinates which we denote with  $\hat{I}$ . Along these directions

$$\partial_{\hat{I}}\beta^J = \partial_{\hat{I}}\beta^J = 0. \tag{4.15}$$

Moreover we should identify any two points in  $\lambda^I$  space, which are related by a global symmetry transformation

$$\lambda^I \sim \lambda^I + (T_A\lambda)^I. \tag{4.16}$$

Deformation by descendant operators are non-physical as they do not modify the action. According to the Ward identities of the flavor symmetries, the vectors  $(T_A\lambda)^I$  correspond to operators which are descendants of currents. The CM direction are then characterized by the following property

$$\partial_{\hat{I}}\bar{\Gamma}^A = \partial_{\hat{I}}\Gamma^A = 0, \tag{4.17}$$

which is the constraint on primary operators  $\mathcal{O}_{\hat{I}}$  and  $\bar{\mathcal{O}}_{\hat{I}}$  derived in section 2.4. The properties in (4.15) and (4.17) have the following interesting implication: if we consider covariant functions which have legs just in the conformal manifold  $\hat{I}, \hat{J}$ , the Lie derivatives  $\mathcal{L}, \bar{\mathcal{L}}$  defined in section 2.8.2 are vanishing.

We can now write the consistency conditions discussed in section 3.5, specializing to the conformal manifold (namely, imposing equations (4.15) and (4.17) for indices along the CM).

- Eqs. (3.48), (3.55) and (3.59) we find that  $a$  is real and constant along the manifold

$$a - \bar{a} = \partial_{\hat{I}}a = \partial_{\hat{I}}a = 0. \tag{4.18}$$

- Specializing to components tangent to the manifold, eqs. (3.42) and (3.8) tell us that

$$-\frac{\chi_{\hat{I}\hat{J}}}{8} = \epsilon_{\hat{I}\hat{J}}^1 = g_{\hat{I}\hat{J}}, \tag{4.19}$$



from which we find that the hermitean part of  $\chi_{\hat{I}\hat{J}}$  is proportional to the Zamolodchikov metric

$$\chi_{\hat{I}\hat{J}} + \bar{\chi}_{\hat{J}\hat{I}} \propto \mathcal{G}_{\hat{I}\hat{J}}. \tag{4.20}$$

- Eq. (3.58) tells us that along the manifold

$$\partial_{\hat{I}} \bar{w}_{\hat{J}} = \partial_{\hat{J}} \bar{w}_{\hat{I}} \tag{4.21}$$

and therefore locally we can write

$$\bar{w}_{\hat{J}} = \partial_{\hat{J}} \omega. \tag{4.22}$$

- Eqs. (3.53), (3.57) give

$$\bar{\chi}_{\hat{J}\hat{I}} - \partial_{\hat{J}} \bar{w}_{\hat{I}} = \chi_{\hat{I}\hat{J}} - \partial_{\hat{I}} w_{\hat{J}} = -\partial_{\hat{J}} \Omega_{\hat{I}}, \tag{4.23}$$

therefore  $\Omega_{\hat{I}}$  is constrained by the relation  $\partial_{\hat{J}} \Omega_{\hat{I}} = \partial_{\hat{I}} \bar{\Omega}_{\hat{J}}$ . We finally find

$$\mathcal{G}_{\hat{I}\hat{J}} \propto \chi_{\hat{I}\hat{J}} + \bar{\chi}_{\hat{J}\hat{I}} = \partial_{\hat{J}} (-2\Omega_{\hat{I}} + \partial_{\hat{I}} (\omega + \bar{\omega})) \tag{4.24}$$

and conclude that the Zamolodchikov metric on the conformal manifold is Kähler, because it satisfies

$$\partial_{\hat{K}} \mathcal{G}_{\hat{I}\hat{J}} - \partial_{\hat{I}} \mathcal{G}_{\hat{K}\hat{J}} = 0, \quad \partial_{\hat{K}} \mathcal{G}_{\hat{I}\hat{J}} - \partial_{\hat{J}} \mathcal{G}_{\hat{I}\hat{K}} = 0. \tag{4.25}$$

An earlier proof of this result was found in [18].

## 5 Conclusions

In this work we presented the superspace formulation of the local RG equation, and derived the Wess-Zumino consistency conditions associated with it. We found that many of the results in the literature regarding supersymmetric RG flows, which were derived using a variety of methods, appear naturally within this framework. However, we believe that there is much room for further exploration. Here are just some of the ideas for future research directions:

- We found physical interpretation for only a few of the consistency conditions. The choice of basis for the consistency conditions given here is not necessarily the optimal one, and perhaps there are alternative representations of the equations which makes the physical meaning more visible.
- Our work here was restricted to theories with chiral anomalies involving only U(1) symmetries. The solution we found for the consistency condition relating the chiral and Weyl anomalies proved instrumental in the derivation of the results of section 4.1, but perhaps a treatment of non-abelian global anomalies may uncover more interesting structure.

- The curved background was defined using the old-minimal formulation of supergravity. This required the usage of the chiral compensator and the Ferrara-Zumino multiplet. However, in order to have a better understanding of the constraints on RG flow related to  $R$ -symmetry, it could prove useful to use the so called  $\mathcal{R}$  multiplet, whose lowest component is the  $R$ -current (see e.g. [54]). Examples for the usage of this multiplet are given in [55, 56]. This adjustment might be non-trivial, as it requires switching to the new-minimal formulation of supergravity, in which some of the identities used in our computation are no longer valid.
- Another possible extension of this work is to consider theories with extended supersymmetry, and to check whether the consistency conditions reveal more structure in the RG flow of such theories.
- It would be interesting to extend this analysis in different dimensions. The case of  $d = 6$ , where an  $a$ -theorem is still lacking [57–60], is especially interesting; it could be that supersymmetry adds essential ingredients.

## Acknowledgments

We are grateful to R. Rattazzi for many useful comments and to H. Osborn for the hospitality and helpful discussions. We would also like to thank L. Bonora for a useful discussion. This work was supported by the Swiss National Science Foundation under grant 200020-150060.

## A Formulas and conventions

### A.1 Classical superWeyl variations

The calculations are done in superspace in old minimal supergravity; the notations and the conventions of [22] are used. Other useful references are: [34, 54, 61–64]. Super-Weyl transformation will be parameterized by the chiral superfield  $\sigma$ . We denote with  $E$  the determinant of the supervielbein and with  $\varphi$  the conformal compensator; the gravitational superfields background is described in terms of the chiral fields  $W_{\alpha\beta\gamma}$  and  $R$  (whose components contain the Weyl tensor and the scalar curvature) and of the real  $G_{\alpha\dot{\alpha}}$  (which contains the traceless Ricci tensor). Under a super Weyl transformations, these fields transform as:

$$\begin{aligned}
 \delta_\sigma \varphi &= \sigma \varphi \\
 \delta_\sigma E^{-1} &= (\sigma + \bar{\sigma}) E^{-1} \\
 \delta_\sigma R &= (\bar{\sigma} - 2\sigma) R - \frac{1}{4} \bar{\mathcal{D}}^2 \bar{\sigma} \\
 \delta_\sigma G_{\alpha\dot{\alpha}} &= -\frac{1}{2} (\sigma + \bar{\sigma}) G_{\alpha\dot{\alpha}} - \mathcal{D}_{\alpha\dot{\alpha}} (i\sigma - i\bar{\sigma}) \\
 \delta_\sigma W_{\alpha\beta\gamma} &= -\frac{3}{2} \sigma W_{\alpha\beta\gamma}
 \end{aligned} \tag{A.1}$$

The Weyl variation of SUSY derivatives (specializing to derivatives acting on scalars):

$$\begin{aligned}
 \delta_\sigma \mathcal{D}_\alpha &= \frac{1}{2} (\sigma - 2\bar{\sigma}) \mathcal{D}_\alpha \\
 \delta_\sigma \mathcal{D}_{\alpha\dot{\alpha}} &= -\frac{1}{2} (\sigma + \bar{\sigma}) \mathcal{D}_{\alpha\dot{\alpha}} - \frac{i}{2} ((\mathcal{D}_\alpha \sigma) \bar{\mathcal{D}}_{\dot{\alpha}} + (\bar{\mathcal{D}}_{\dot{\alpha}} \bar{\sigma}) \mathcal{D}_\alpha) \\
 \delta_\sigma \mathcal{D}^2 &= (\sigma - 2\bar{\sigma}) \mathcal{D}^2 + 2(\mathcal{D}^\alpha \sigma) \mathcal{D}_\alpha
 \end{aligned} \tag{A.2}$$

where  $\mathcal{D}_{\alpha\dot{\alpha}} = \frac{i}{2} \{\mathcal{D}_\alpha, \bar{\mathcal{D}}_{\dot{\alpha}}\}$ .

## A.2 Covariant derivatives

Covariant derivatives acting on chiral fields

$$\begin{aligned}
 A_J^{\alpha I} &= (e^{-V})^{I\bar{K}} (\mathcal{D}^\alpha e^V)_{\bar{K}J} \\
 \bar{\nabla}^\alpha e^V &= 0 \\
 \bar{\nabla}^\alpha \lambda^I &= \mathcal{D}^\alpha \lambda^I + (A^\alpha)_J^I \lambda^J \\
 \bar{\nabla}^{\alpha\dot{\alpha}} \lambda^I &= \mathcal{D}^{\alpha\dot{\alpha}} \lambda^I + \frac{i}{2} \bar{\mathcal{D}}^{\dot{\alpha}} A_J^{\alpha I} \lambda^J \\
 \bar{\nabla}^2 \lambda^I &= \mathcal{D}^2 \lambda^I + 2A_J^{\alpha I} \mathcal{D}_\alpha \lambda^J + (A_J^{\alpha I} A_{\alpha K}^J + \mathcal{D}^\alpha A_{\alpha K}^I) \lambda^K
 \end{aligned} \tag{A.3}$$

Covariant derivatives acting on anti-chiral fields

$$\begin{aligned}
 B_J^{\dot{\alpha}\bar{I}} &= (\bar{\mathcal{D}}^{\dot{\alpha}} e^V)_{\bar{J}K} (e^{-V})^{K\bar{L}} \\
 \bar{\nabla}^{\dot{\alpha}} e^V &= 0 \\
 \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}} &= \bar{\mathcal{D}}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}} + (B^{\dot{\alpha}})_J^{\bar{I}} \bar{\lambda}^{\bar{J}} \\
 \bar{\nabla}^{\alpha\dot{\alpha}} \bar{\lambda}^{\bar{I}} &= \mathcal{D}^{\alpha\dot{\alpha}} \bar{\lambda}^{\bar{I}} + \frac{i}{2} \mathcal{D}^\alpha B_J^{\dot{\alpha}\bar{I}} \bar{\lambda}^{\bar{J}} \\
 \bar{\nabla}^2 \bar{\lambda}^{\bar{I}} &= \bar{\mathcal{D}}^2 \bar{\lambda}^{\bar{I}} + 2B_{\dot{\alpha}\bar{J}}^{\bar{I}} \bar{\mathcal{D}}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} + (B_{\dot{\alpha}\bar{J}}^{\bar{I}} B_{\bar{K}}^{\dot{\alpha}\bar{J}} + \bar{\mathcal{D}}_{\dot{\alpha}} B_{\bar{K}}^{\dot{\alpha}\bar{I}}) \bar{\lambda}^{\bar{K}}
 \end{aligned} \tag{A.4}$$

## A.3 Useful identities

$$\begin{aligned}
 \bar{\mathcal{D}}^{\dot{\alpha}} G_{\alpha\dot{\alpha}} &= \mathcal{D}_\alpha R \\
 \bar{\nabla}_{\dot{\alpha}} W_\alpha^\pm &= 0 \\
 \bar{\nabla}^\alpha W_\alpha^\pm &= \bar{\nabla}_{\dot{\alpha}} (\bar{W}^\pm)^{\dot{\alpha}} \\
 \bar{\nabla}_\alpha \bar{\nabla}^2 \lambda^I &= 4\bar{R} \bar{\nabla}_\alpha \lambda^I \\
 \bar{\nabla}^2 \bar{\nabla}_\alpha \lambda^I &= 4R \bar{\nabla}_\alpha \lambda^I - 4(W_\alpha \lambda)^I \\
 \bar{\nabla}_{\dot{\alpha}} \bar{\nabla}^2 \lambda^I &= 4(G_{\alpha\dot{\alpha}} + i \nabla_{\alpha\dot{\alpha}}) \bar{\nabla}^\alpha \lambda^I - 4(\bar{W}_{\dot{\alpha}}^+ \lambda)^I \\
 \bar{\nabla}^\alpha \bar{\nabla}_{\dot{\alpha}} \bar{\nabla}_\alpha \lambda^I &= -\frac{1}{2} \bar{\nabla}_{\dot{\alpha}} \bar{\nabla}^2 \lambda^I - 2G_{\alpha\dot{\alpha}} \bar{\nabla}^\alpha \lambda^I + 2(\bar{W}_{\dot{\alpha}}^+ \lambda)^I \\
 \bar{\nabla}^2 \bar{\nabla}^2 \lambda^I &= -8i G_{\alpha\dot{\alpha}} \bar{\nabla}^{\alpha\dot{\alpha}} \lambda^I - 8 \nabla_{\alpha\dot{\alpha}} \bar{\nabla}^{\alpha\dot{\alpha}} \lambda^I \\
 &\quad + 4\mathcal{D}^\alpha R \bar{\nabla}_\alpha \lambda^I + 8R \bar{\nabla}^2 \lambda^I - 8(W^\alpha)_K^I \bar{\nabla}_\alpha \lambda^K - 4(\bar{\nabla}^\alpha W_\alpha)_K^I \lambda^K
 \end{aligned} \tag{A.5}$$

where we used the notations

$$\begin{aligned}
 (W_\alpha^+)_J^I &\equiv (W_\alpha)_J^I & (\bar{W}_{\dot{\alpha}}^+)_J^I &\equiv (e^{-V})^{I\bar{L}} (\bar{W}_{\dot{\alpha}})_L^{\bar{K}} (e^V)_{\bar{K}J}, \\
 (\bar{W}_\alpha^-)_J^I &\equiv (\bar{W}_{\dot{\alpha}}^-)_J^I & (W_\alpha^-)_J^I &\equiv (e^V)_{\bar{J}K} (W_\alpha)_L^K (e^{-V})^{LI}.
 \end{aligned} \tag{A.6}$$

## B Useful definitions and results

### B.1 Notations

$$\begin{aligned}
 \beta^I &\equiv b^I + \Gamma^A (T_A \lambda)^I \\
 U_I^J &\equiv \delta_I^J + \partial_I \beta^J \\
 \Lambda^I &\equiv (U^{-1})_K^I (\nabla^2 \lambda^K + 4\beta^K \bar{R}) \\
 \Pi^{IJ} &\equiv \nabla^\alpha \lambda^I \bar{\nabla}_\alpha \lambda^J - \frac{1}{2} (\beta^I \Lambda^J + \beta^J \Lambda^I)
 \end{aligned} \tag{B.1}$$

## B.2 Lie derivatives in parameter space

The Lie derivatives of a covariant function  $Y_{IJ\dots}^K$ , where the indices  $I, J, \dots$  run over the marginal operators, is given by

$$\begin{aligned}\mathcal{L}(Y_{IJ\dots}^K) &\equiv \beta^L \partial_L Y_{IJ\dots}^K + \gamma_I^L Y_{LJ\dots}^K + \tilde{\gamma}_{\bar{J}}^{\bar{L}} Y_{I\bar{L}\dots}^K + \dots - \gamma_L^K Y_{IJ\dots}^L - \dots \\ \bar{\mathcal{L}}(Y_{IJ\dots}^K) &\equiv \bar{\beta}^{\bar{L}} \partial_{\bar{L}} Y_{IJ\dots}^K + \tilde{\gamma}_I^{\bar{L}} Y_{LJ\dots}^K + \tilde{\gamma}_{\bar{J}}^{\bar{L}} Y_{I\bar{L}\dots}^K + \dots - \tilde{\gamma}_L^K Y_{IJ\dots}^L - \dots\end{aligned}\quad (\text{B.2})$$

For tensors which multiply  $W_\alpha^A$  we find

$$\begin{aligned}\mathcal{L}(Y_{A\dots}) &\equiv \beta^I \partial_I Y_{A\dots}^K + (\gamma_A^B + if_{AC}^B \Gamma^C) Y_{B\dots} + \dots \\ \bar{\mathcal{L}}(Y_{A\dots}) &\equiv \bar{\beta}^{\bar{I}} \partial_{\bar{I}} Y_{A\dots} + \gamma_A^B Y_{B\dots} + \dots\end{aligned}\quad (\text{B.3})$$

For tensors which multiply  $\bar{W}_\alpha^A$  we find

$$\begin{aligned}\mathcal{L}(Y_{A\dots}) &\equiv \beta^I \partial_I Y_{A\dots}^K + \bar{\gamma}_A^B Y_{B\dots} + \dots \\ \bar{\mathcal{L}}(Y_{A\dots}) &\equiv \bar{\beta}^{\bar{I}} \partial_{\bar{I}} Y_{A\dots} + (\bar{\gamma}_A^B + if_{AC}^B \Gamma^C) Y_{B\dots} + \dots\end{aligned}\quad (\text{B.4})$$

The anomalous dimension matrices are given by

$$\begin{aligned}\gamma_I^J &\equiv \partial_I \beta^J \\ \tilde{\gamma}_I^J &\equiv \partial_I \Gamma^A (T_A \lambda)^J \\ \gamma_A^B &= \partial_I \Gamma^B (T_A \lambda)^I\end{aligned}\quad (\text{B.5})$$

It is also useful to introduce the following notations

$$\begin{aligned}\gamma_{JK}^I &= (U^{-1})_L^I \partial_{KJ} \beta^L \\ \gamma_{KL}^{IJ} &= \gamma_{KL}^{(I} \beta^{J)}\end{aligned}\quad (\text{B.6})$$

Using the covariance of the function  $Y$  and the consistency condition (2.56) one can show that this Lie derivative has the following properties:

$$\begin{aligned}\beta^I \mathcal{L}(Y_{I\dots}) &= \mathcal{L}(\beta^I Y_{I\dots}) \\ (T_A \lambda)^I \mathcal{L}(Y_{I\dots}) &= \mathcal{L}((T_A \lambda)^I Y_{I\dots})\end{aligned}\quad (\text{B.7})$$

$$\begin{aligned}[\mathcal{L}, \bar{\mathcal{L}}] &= 0 \\ [\mathcal{L}, \partial_I] Y_{J\dots} &= -\partial_{IJ} \beta^K Y_{K\dots} + \dots \\ [\mathcal{L}, \partial_{\bar{I}}] Y_{J\dots} &= -\partial_{\bar{I}J} \Gamma^A (T_A \lambda)^K Y_{K\dots} + \dots \\ [\bar{\mathcal{L}}, \partial_I] Y_{\bar{J}\dots} &= -\partial_{I\bar{J}} \Gamma^A (\bar{T}_A \bar{\lambda})^{\bar{L}} Y_{\bar{L}\dots} + \dots \\ [\bar{\mathcal{L}}, \partial_{\bar{I}}] Y_{\bar{J}\dots} &= -\partial_{\bar{I}\bar{J}} \Gamma^A (\bar{T}_A \bar{\lambda})^{\bar{L}} Y_{\bar{L}\dots} + \dots\end{aligned}\quad (\text{B.8})$$

$$\begin{aligned}\mathcal{L}(Y_I) - Y_K \gamma_{IJ}^K \beta^J &= U_I^J \mathcal{L}(Y_K (U^{-1})_J^K) \\ \bar{\mathcal{L}}(Y_I) - Y_K \tilde{\gamma}_{IJ}^K \beta^J &= U_I^J \bar{\mathcal{L}}(Y_K (U^{-1})_J^K)\end{aligned}\quad (\text{B.9})$$

## B.3 Super-Weyl variations of functions

$$\begin{aligned}\Delta_\sigma^{\text{SW}}((e^V)_{IJ}) &= (\sigma + \bar{\sigma}) \Gamma^A (e^V T_A)_{IJ} \\ \Delta_\sigma^{\text{SW}}((A_\alpha)_J^I) &= \frac{1}{2}(\sigma - 2\bar{\sigma})(A_\alpha)_J^I + \mathcal{D}_\alpha \sigma \Gamma_J^I + (\sigma + \bar{\sigma}) \partial_K \Gamma_J^I \nabla_\alpha \lambda^J \\ \Delta_\sigma^{\text{SW}}(Y_I \nabla^\alpha \lambda^I) &= \frac{1}{2}(\sigma - 2\bar{\sigma}) Y_I \nabla^\alpha \lambda^I + \mathcal{D}^\alpha \sigma Y_I \beta^I \\ &\quad + \sigma \mathcal{L}(Y_I) \nabla^\alpha \lambda^I + \bar{\sigma} \bar{\mathcal{L}}(Y_I) \nabla^\alpha \lambda^I \\ \Delta_\sigma^{\text{SW}}(Y_{\bar{I}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}}) &= \frac{1}{2}(\bar{\sigma} - 2\sigma) Y_{\bar{I}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}} + \bar{\mathcal{D}}^{\dot{\alpha}} \bar{\sigma} Y_{\bar{K}} \bar{\beta}^{\bar{K}} \\ &\quad + \sigma \mathcal{L}(Y_{\bar{I}}) \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}} + \bar{\sigma} \bar{\mathcal{L}}(Y_{\bar{I}}) \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{I}}\end{aligned}$$

$$\begin{aligned}
 \Delta_\sigma^{\text{SW}}(Y_I \nabla^2 \lambda^I) &= (\sigma - 2\bar{\sigma}) Y_I \nabla^2 \lambda^I && + \mathcal{D}^2 \sigma Y_K \beta^K + 2\mathcal{D}^\alpha \sigma Y_K U_I^K \nabla_\alpha \lambda^I \\
 &+ \sigma \mathcal{L}(Y_I) \nabla^2 \lambda^I && + \sigma Y_K \partial_{IJ} \beta^K \nabla^\alpha \lambda^I \nabla_\alpha \lambda^J \\
 &+ \bar{\sigma} \bar{\mathcal{L}}(Y_I) \nabla^2 \lambda^I && + \bar{\sigma} Y_K \partial_{IJ} (\Gamma_L^K \lambda^L) \nabla^\alpha \lambda^I \nabla_\alpha \lambda^J \\
 \Delta_\sigma^{\text{SW}}(Y_I \Lambda^I) &= (\sigma - 2\bar{\sigma}) Y_I \Lambda^I && + 2Y_I (\mathcal{D}^\alpha \sigma) \nabla_\alpha \lambda^I \\
 &+ \sigma \mathcal{L}(Y_I) \Lambda^I && + \bar{\sigma} \bar{\mathcal{L}}(Y_I) \Lambda^I \\
 &+ \sigma Y_I \gamma_{KL}^I \Pi^{KL} && + \bar{\sigma} Y_I \gamma_{KL}^I \Pi^{KL} \\
 \Delta_\sigma^{\text{SW}}(Y_{IJ} \Pi^{IJ}) &= (\sigma - 2\bar{\sigma}) Y_{IJ} \Pi^{IJ} && + \sigma (\mathcal{L}(Y_{IJ}) - Y_{KL} \gamma_{IJ}^{KL}) \Pi^{IJ} \\
 &+ \bar{\sigma} (\bar{\mathcal{L}}(Y_{IJ}) - Y_{KL} \gamma_{IJ}^{KL}) \Pi^{IJ} \\
 \Delta_\sigma^{\text{SW}}(Y_I \nabla^{\alpha\dot{\alpha}} \lambda^I) &= -\frac{1}{2}(\sigma + \bar{\sigma}) Y_I \nabla^{\alpha\dot{\alpha}} \lambda^I && + \mathcal{D}^{\alpha\dot{\alpha}} \sigma Y_I \beta^I \\
 &-\frac{i}{2} Y_I \partial_{\bar{J}} \beta^I \mathcal{D}^\alpha \sigma \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} && + \frac{i}{2} Y_I (-\delta_J^I + \partial_J \Gamma_K^I \lambda^K) \bar{\mathcal{D}}^{\dot{\alpha}} \bar{\sigma} \nabla^\alpha \lambda^J \\
 &+ \sigma \mathcal{L}(Y_I) \nabla^{\alpha\dot{\alpha}} \lambda^I && - \frac{i}{2} \sigma Y_I \partial_{J\bar{K}} \beta^I \nabla^\alpha \lambda^J \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{K}} \\
 &+ \bar{\sigma} \bar{\mathcal{L}}(Y_I) \nabla^{\alpha\dot{\alpha}} \lambda^I && - \frac{i}{2} \bar{\sigma} Y_I \partial_{J\bar{K}} \Gamma_L^I \lambda^L \nabla^\alpha \lambda^J \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{K}} \\
 \Delta_\sigma^{\text{SW}}(Y_A W^{\alpha A}) &= -\frac{3}{2} \sigma Y_A W^{\alpha A} && + i Y_A \mathcal{D}^{\alpha\dot{\alpha}} \sigma \partial_{\bar{I}} \Gamma^A \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{I}} - \frac{1}{4} Y_A \mathcal{D}^\alpha \sigma \left( \partial_{\bar{I}} \Gamma^A \bar{\nabla}^2 \bar{\lambda}^{\bar{I}} + \partial_{\bar{I}\bar{J}} \Gamma^A \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{I}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \right) \\
 &+ i Y_A \mathcal{D}^{\alpha\dot{\alpha}} \sigma \partial_{\bar{I}} \Gamma^A \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{I}} && - \frac{1}{4} Y_A \bar{\mathcal{D}}^2 \bar{\sigma} \partial_{\bar{I}} \Gamma^A \nabla^\alpha \lambda^{\bar{I}} + Y_A \bar{\mathcal{D}}_{\dot{\alpha}} \bar{\sigma} \left( i \partial_{\bar{I}} \Gamma^A \nabla^{\alpha\dot{\alpha}} \lambda^{\bar{I}} + \frac{1}{2} \partial_{\bar{I}\bar{J}} \Gamma^A \nabla^\alpha \lambda^{\bar{I}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \right) \\
 &- \frac{1}{4} Y_A \bar{\mathcal{D}}^2 \bar{\sigma} \partial_{\bar{I}} \Gamma^A \nabla^\alpha \lambda^{\bar{I}} && + Y_A \partial_{\bar{I}\bar{J}} \Gamma^A \left( -\frac{1}{4} \nabla^\alpha \lambda^{\bar{I}} \bar{\nabla}^2 \bar{\lambda}^{\bar{J}} + i \nabla^{\alpha\dot{\alpha}} \lambda^{\bar{I}} \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \right) \\
 &+ (\sigma + \bar{\sigma}) \left( \mathcal{L}(Y_A) W^{\alpha A} + Y_A \partial_{\bar{I}\bar{J}} \Gamma^A \left( -\frac{1}{4} \nabla^\alpha \lambda^{\bar{I}} \bar{\nabla}^2 \bar{\lambda}^{\bar{J}} + i \nabla^{\alpha\dot{\alpha}} \lambda^{\bar{I}} \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \right) \right. \\
 &\left. - \frac{1}{4} Y_A \partial_{\bar{I}\bar{J}\bar{K}} \Gamma^A \nabla^\alpha \lambda^{\bar{I}} \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{K}} \right)
 \end{aligned} \tag{B.10}$$

where we used the constraint that  $\Gamma$  is hermitean in the sense of eq. (2.34):  $\bar{\Gamma} e^V = e^V \Gamma$ .

#### B.4 Integrations by parts

The following identities (valid up to total derivatives) are useful in order to perform some integration by parts which are very common in the task of writing the consistency conditions ( $A$  is an arbitrary covariant function of the sources).

$$\begin{aligned}
 \sigma_{[1} \mathcal{D}^2 \sigma_{2]} A &= -\sigma_{[1} \mathcal{D}^\alpha \sigma_{2]} \mathcal{D}_\alpha A \\
 \sigma_{[1} \bar{\mathcal{D}}^2 \bar{\sigma}_{2]} A &= \sigma_{[1} \bar{\sigma}_{2]} \bar{\mathcal{D}}^2 A \\
 \sigma_{[1} \mathcal{D}^{\alpha\dot{\alpha}} \sigma_{2]} A_{\alpha\dot{\alpha}} &= \frac{i}{2} \sigma_{[1} \mathcal{D}^\alpha \sigma_{2]} \bar{\mathcal{D}}^{\dot{\alpha}} A_{\alpha\dot{\alpha}} \\
 \sigma_{[1} \mathcal{D}^{\alpha\dot{\alpha}} \bar{\sigma}_{2]} A_{\alpha\dot{\alpha}} &= -\frac{i}{2} \mathcal{D}^\alpha \sigma_{[1} \bar{\mathcal{D}}^{\dot{\alpha}} \bar{\sigma}_{2]} A_{\alpha\dot{\alpha}} - \frac{i}{2} \sigma_{[1} \bar{\sigma}_{2]} \bar{\mathcal{D}}^{\dot{\alpha}} \mathcal{D}^\alpha A_{\alpha\dot{\alpha}} \\
 \mathcal{D}^\alpha \sigma_{[1} \mathcal{D}_{\alpha\dot{\alpha}} \sigma_{2]} A^{\dot{\alpha}} &= \sigma_{[1} \mathcal{D}^\alpha \sigma_{2]} \left( \frac{i}{2} \bar{\mathcal{D}}_{\dot{\alpha}} \mathcal{D}_\alpha A^{\dot{\alpha}} + \frac{i}{4} \mathcal{D}_\alpha \bar{\mathcal{D}}_{\dot{\alpha}} A^{\dot{\alpha}} + i G_{\alpha\dot{\alpha}} A^{\dot{\alpha}} \right) \\
 \mathcal{D}^\alpha \sigma_{[1} \mathcal{D}_{\alpha\dot{\alpha}} \bar{\sigma}_{2]} A^{\dot{\alpha}} &= \sigma_{[1} \bar{\sigma}_{2]} (2i \bar{\mathcal{D}}_{\dot{\alpha}} (\bar{R} A^{\dot{\alpha}}) - \frac{i}{2} \bar{\mathcal{D}}_{\dot{\alpha}} \mathcal{D}^2 A^{\dot{\alpha}}) + \mathcal{D}^\alpha \sigma_{[1} \bar{\mathcal{D}}^{\dot{\alpha}} \bar{\sigma}_{2]} \frac{i}{2} \mathcal{D}_\alpha A^{\dot{\alpha}} \\
 \mathcal{D}^\alpha \sigma_{[1} \bar{\mathcal{D}}^2 \bar{\sigma}_{2]} A_\alpha &= \sigma_{[1} \bar{\sigma}_{2]} (\mathcal{D}^\alpha \bar{\mathcal{D}}^2 A_\alpha - \mathcal{D}^\alpha (4R A_\alpha)) + \mathcal{D}^\alpha \sigma_{[1} \bar{\mathcal{D}}^{\dot{\alpha}} \bar{\sigma}_{2]} 2\bar{\mathcal{D}}_{\dot{\alpha}} A_\alpha
 \end{aligned} \tag{B.11}$$

#### B.5 Computation of the consistency conditions

As an example for the type of computations necessary for deriving the Wess-Zumino consistency conditions, we compute the contribution of the  $\chi_{I\bar{J}}$  anomaly.

$$\begin{aligned}
 \Delta_{\sigma_2}^{\text{SW}} \left( \frac{1}{2} E^{-1} \sigma_1 \chi_{I\bar{J}} \nabla_\alpha \lambda^I \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} G^{\alpha\dot{\alpha}} \right) &= \frac{1}{2} E^{-1} \left( \sigma_1 \chi_{I\bar{J}} \mathcal{D}^{\alpha\dot{\alpha}} (i\bar{\sigma}_2 - i\sigma_2) \nabla_\alpha \lambda^I \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \right. \\
 &+ \beta^I \chi_{I\bar{J}} \mathcal{D}_\alpha \sigma_2 \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} G^{\alpha\dot{\alpha}} - \beta^{\bar{J}} \chi_{I\bar{J}} \mathcal{D}_{\dot{\alpha}} \bar{\sigma}_2 \nabla_\alpha \lambda^I G^{\alpha\dot{\alpha}} \\
 &\left. + \bar{\sigma}_2 \bar{\mathcal{L}}(\chi_{I\bar{J}}) \nabla_\alpha \lambda^I \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} G^{\alpha\dot{\alpha}} + \sigma_2(\dots) \right) \tag{B.12}
 \end{aligned}$$

The contribution to the consistency condition (see eq. (3.3)) is

$$\begin{aligned}
 \delta_{\sigma_1, \sigma_2}^{WZ} \supset & (\sigma_{[1} \bar{\sigma}_{2]}) \left( \left( \bar{\mathcal{L}}(\chi_{I\bar{J}}) - \partial_{\bar{J}}(\bar{\beta}^{\bar{k}} \chi_{I\bar{K}}) - \chi_{I\bar{J}} \right) \nabla_{\alpha} \lambda^I \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} G^{\alpha\dot{\alpha}} \right. \\
 & - 2i \bar{\beta}^{\bar{K}} \chi_{I\bar{K}} G^{\alpha\dot{\alpha}} \nabla_{\alpha\dot{\alpha}} \lambda^I - \bar{\beta}^{\bar{k}} \chi_{I\bar{K}} \mathcal{D}^{\alpha} R \nabla_{\alpha} \lambda^I \\
 & - \frac{1}{2} \chi_{I\bar{J}} \nabla^2 \lambda^I \bar{\nabla}^2 \bar{\lambda}^{\bar{J}} - 2 \chi_{I\bar{J}} \nabla^{\alpha\dot{\alpha}} \lambda^I \nabla_{\alpha\dot{\alpha}} \bar{\lambda}^{\bar{J}} - \frac{1}{2} \chi_{I\bar{J}} \bar{\nabla}_{\dot{\alpha}} \nabla^2 \lambda^I \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} - \frac{1}{4} \chi_{I\bar{J}} \nabla^{\alpha} \lambda^I \nabla_{\alpha} \bar{\nabla}^2 \bar{\lambda}^{\bar{J}} \\
 & + \chi_{I\bar{J}} \nabla^{\alpha} \lambda^I (W_{\alpha}^{-})_{\bar{K}}^{\bar{J}} \bar{\lambda}^{\bar{K}} - \frac{1}{2} \partial_{\bar{k}} \chi_{I\bar{J}} \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{K}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \nabla^2 \lambda^I + i \partial_{\bar{K}} \chi_{I\bar{J}} \nabla_{\alpha} \lambda^I \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{K}} \nabla^{\alpha\dot{\alpha}} \bar{\lambda}^{\bar{J}} \\
 & - \frac{1}{2} \partial_K \chi_{I\bar{J}} \nabla^{\alpha} \lambda^K \nabla_{\alpha} \lambda^I \bar{\nabla}^2 \bar{\lambda}^{\bar{J}} - i (\partial_K \chi_{I\bar{J}} + \partial_I \chi_{K\bar{J}}) \nabla_{\alpha} \lambda^I \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \nabla^{\alpha\dot{\alpha}} \lambda^K \\
 & - \frac{1}{2} \partial_{K\bar{L}} \chi_{I\bar{J}} \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{L}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \nabla^{\alpha} \lambda^K \nabla_{\alpha} \lambda^I \\
 & + (\sigma_{[1} \mathcal{D}^{\alpha} \sigma_{2]}) \left( \beta^I \chi_{I\bar{J}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} G_{\alpha\dot{\alpha}} + \frac{1}{2} \chi_{I\bar{J}} \nabla_{\alpha} \lambda^I \bar{\nabla}^2 \bar{\lambda}^{\bar{J}} + i \chi_{I\bar{J}} \nabla_{\alpha\dot{\alpha}} \lambda^I \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \right. \\
 & \left. + \frac{1}{2} \partial_{\bar{K}} \chi_{I\bar{J}} \nabla_{\alpha} \lambda^I \bar{\nabla}_{\dot{\alpha}} \bar{\lambda}^{\bar{K}} \bar{\nabla}^{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \right) \\
 & + (\mathcal{D}^{\alpha} \sigma_{[1} \bar{\mathcal{D}}^{\dot{\alpha}} \bar{\sigma}_{2]}) \left( \frac{1}{2} \chi_{I\bar{J}} \mathcal{D}_{\alpha} \lambda^I \bar{\mathcal{D}}_{\dot{\alpha}} \bar{\lambda}^{\bar{J}} \right)
 \end{aligned} \tag{B.13}$$

## C Basic examples

### C.1 The Wess-Zumino model

In the Wess-Zumino model each classically marginal coupling  $\lambda^I$  is a trilinear in the elementary chiral superfields  $\Phi_i$ , which are in the fundamental representation of the  $U(N)$  symmetry group. Each coupling index  $I$  corresponds to a symmetric 3-tensor product of fundamentals, e.g.:  $I = (ijk)$ . one finds that the holomorphic  $\beta$ -function vanishes to all orders<sup>14</sup>

$$b^I = 0, \tag{C.1}$$

and the anomalous dimension matrix  $\Gamma^A(T_A)_j^i = \Gamma_j^i$  can be computed perturbatively; the leading and next-to-leading contributions are (see, e.g. [6]):

$$\begin{aligned}
 (\Gamma_j^i)_1 &= \frac{1}{2(2\pi)^2} \lambda^{ikl} \bar{\lambda}^{\bar{i}\bar{k}\bar{l}} (e^V)_{\bar{k}k} (e^V)_{\bar{l}l} (e^V)_{\bar{i}j} \\
 (\Gamma_j^i)_2 &= -\frac{1}{2(4\pi)^4} \left( (e^V)_{\bar{j}j} (e^V)_{\bar{k}k} (e^V)_{\bar{q}r} \bar{\lambda}^{\bar{j}\bar{k}\bar{q}} \right) \left( \lambda^{rst} \bar{\lambda}^{\bar{r}\bar{s}\bar{t}} (e^V)_{\bar{s}s} (e^V)_{\bar{t}t} (e^V)_{\bar{u}u} \right) \lambda^{uki}.
 \end{aligned} \tag{C.2}$$

Using the results of section 2.6 we read the corresponding terms in the LRG equation

$$\begin{aligned}
 \beta^I &= \Gamma_J^I \lambda^J & \bar{\beta}^{\bar{I}} &= (\beta^I)^* \\
 \rho_I^A &= -\partial_I \Gamma^A & \rho_{\bar{I}}^A &= \partial_{\bar{I}} \Gamma^A,
 \end{aligned} \tag{C.3}$$

where

$$\Gamma_J^I = \Gamma^A(T_A)_J^I = \Gamma_j^i \delta_m^k \delta_n^l + \delta_j^i \Gamma_m^k \delta_n^l + \delta_j^i \delta_m^k \Gamma_n^l. \tag{C.4}$$

<sup>14</sup>If one is interested just in composite marginal operators, one may use the ambiguity discussed in section 2.5 to shift  $\beta$  and  $\gamma$ . We use in the following discussion the canonical choice  $b^I = 0$ .

## C.2 Gauge theories

In the case of a gauge theory with a simple gauge group, the global symmetry group is the product  $\mathcal{G} = \prod_i U(N_i)$ , where the index  $i$  runs on the representations of the matter fields and each  $U(N_i)$  corresponds to a group of  $N_i$  fields in the same gauge representation. The symmetry group  $\mathcal{G}$  is in general broken by the chiral coupling spurions.

The gauge coupling itself can be promoted to a chiral superfield:  $S = \frac{4\pi}{g_h^2} - i\frac{\Theta}{2\pi}$ . We call  $U(1)_K$  the symmetry which rotates with the same phase all the matter fields charged under the gauge group. Under  $U(1)_K$  the superfield  $S$  transform in a way specified by Konishi anomaly [65, 66]:

$$S \rightarrow S + \frac{i\Lambda}{\pi} \sum_i N_i t(r_i) \tag{C.5}$$

where the index  $i$  run on the matter fields gauge representation and  $t(r_i)$  is the its Dynkin index. For our purpose it is convenient to consider the combination  $\lambda_G = e^{-S}$  which transforms linearly under  $U(1)_K$  and vanishes in the limit of the free theory:

$$\lambda_G \rightarrow \lambda_G e^{-\frac{i\Lambda}{\pi} \sum_i N_i t(r_i)}. \tag{C.6}$$

The gauge coupling  $\lambda_G$  does not transform under all the other global symmetries in  $\mathcal{G}$  which are orthogonal to  $U(1)_K$ .

Inside each  $U(N_i)$  factor, one can diagonalize the matrix  $\Gamma_i = \Gamma^{A_i} T_{A_i}$ , where the label  $A_i$  runs on the  $U(N_i)$  generators. We denote with  $\gamma^{(a,i)}$  the  $N_i$  eigenvalues of the matrix  $\Gamma_i$ , with  $a = 1 \dots N_i$ ;  $\gamma^{(a,i)}$  are the anomalous dimension of the elementary fields in the gauge representation  $i$ .

The action of the global symmetry background gauge field on the chiral coupling  $\lambda^G$  can be written as

$$\Gamma^{A_i}(T_{A_i} \lambda^G) = -\frac{1}{\pi} t(r_i) \sum_a \gamma^{(a,i)} \lambda^G. \tag{C.7}$$

The holomorphic  $\beta$ -function for  $\lambda_G$  is given by the one-loop result:

$$b^{\lambda_G} = - \left( \frac{(3t(G) - \sum_i N_i t(r_i))}{2\pi} \right) \lambda_G \tag{C.8}$$

This  $\beta$ -function  $b^{\lambda_G}$  can be non-zero at the conformal fixed points. Using eq. (2.25), we find the physical  $\beta$ -function:

$$\beta^{\lambda_G} = -\frac{1}{\pi} \left( \frac{3}{2} t(G) - \sum_i t(r_i) \left( \frac{N_i}{2} - \sum_a \gamma^{(a,i)} \right) \right) \lambda_G. \tag{C.9}$$

This coincides with the numerator of the NSVZ  $\beta$ -function [8, 9] and vanishes at the fixed point. The connection between space-time dependent couplings and the denominator of NSVZ has been studied in [45].

## D The super-Weyl anomaly and the Zamolodchikov metric

In this appendix we explain the relation between the Zamolodchikov metric and the super-Weyl anomaly coefficient  $g_{I\bar{J}}$  defined in (3.8)

$$\mathcal{A}_\sigma^{\text{SW}} \supset \int d^8z E^{-1} \sigma g_{I\bar{J}} \nabla^2 \lambda^I \bar{\nabla}^2 \bar{\lambda}^{\bar{I}} . \quad (\text{D.1})$$

The following is a variation of the argument appearing in [28], which discusses the  $c$  anomaly coefficient.

The Zamolodchikov metric  $\mathcal{G}_{I\bar{J}}$  is defined as the matrix controlling the two point function of marginal operators at the fixed point. It is convenient to use the following expression

$$\langle O_I(z_1) \bar{O}_{\bar{J}}(z_2) \rangle = \mathcal{G}_{I\bar{J}} \bar{\mathcal{D}}^2 \mathcal{D}^2 \square \left( \frac{1}{(x_1 - x_2)^6} \delta^{(4)}(\theta_1 - \theta_2) \right) . \quad (\text{D.2})$$

Notice that the singularity at  $x_1 = x_2$  has to be regulated. Using the method of differential regularization [28], we replace  $\frac{1}{x^6}$  with

$$\mathcal{R} \left( \frac{1}{x^6} \right) = -\frac{1}{32} \square^2 \left( \frac{1}{x^2} \ln(\mu^2 x^2) \right) \quad (\text{D.3})$$

where  $\mu$  is an arbitrary renormalization scale. The dependence of the regulated correlator on the scale  $\mu$  is thus given by

$$\mu \frac{\partial}{\partial \mu} \langle O_I(z_1) \bar{O}_{\bar{J}}(z_2) \rangle = \frac{\pi^2}{4} \mathcal{G}_{I\bar{J}} \bar{\mathcal{D}}^2 \mathcal{D}^2 \square \delta^{(8)}(z_1 - z_2) \quad (\text{D.4})$$

On the other hand, the dependence on  $\mu$  can be found by noticing that the generating functional  $\mathcal{W}$  is invariant under a change of renormalization scale combined with a global rescaling of the chiral compensator

$$\mu \frac{\partial}{\partial \mu} \mathcal{W} = \left( \int d^6z \varphi \frac{\delta}{\delta \varphi} + c.c. \right) \mathcal{W} . \quad (\text{D.5})$$

Using the SLRG equation at the fixed point we can now write

$$\begin{aligned} \mu \frac{\partial}{\partial \mu} \langle O_I(z_1) \bar{O}_{\bar{J}}(z_2) \rangle &= -\frac{\delta}{\delta \lambda^I(z_1)} \frac{\delta}{\delta \lambda^{\bar{I}}(z_2)} \mathcal{A}_{\sigma=1}^{\text{SW}} \Big|_{\lambda=0} \\ &= -\frac{1}{8} (g_{I\bar{J}} + \bar{g}_{\bar{J}I}) \bar{\mathcal{D}}^2 \mathcal{D}^2 \square \delta^{(8)}(z_1 - z_2) . \end{aligned} \quad (\text{D.6})$$

Comparing with (D.4) we conclude that

$$\mathcal{G}_{I\bar{J}} \propto g_{I\bar{J}} + \bar{g}_{\bar{J}I} . \quad (\text{D.7})$$

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.



## References

- [1] H. Osborn, *Derivation of a Four-dimensional  $c$  Theorem*, *Phys. Lett. B* **222** (1989) 97 [[INSPIRE](#)].
- [2] I. Jack and H. Osborn, *Analogs for the  $c$  Theorem for Four-dimensional Renormalizable Field Theories*, *Nucl. Phys. B* **343** (1990) 647 [[INSPIRE](#)].
- [3] H. Osborn, *Weyl consistency conditions and a local renormalization group equation for general renormalizable field theories*, *Nucl. Phys. B* **363** (1991) 486 [[INSPIRE](#)].
- [4] Y. Nakayama, *Scale invariance vs conformal invariance*, *Phys. Rept.* **569** (2015) 1 [[arXiv:1302.0884](#)] [[INSPIRE](#)].
- [5] F. Baume, B. Keren-Zur, R. Rattazzi and L. Vitale, *The local Callan-Symanzik equation: structure and applications*, *JHEP* **08** (2014) 152 [[arXiv:1401.5983](#)] [[INSPIRE](#)].
- [6] I. Jack and H. Osborn, *Constraints on RG Flow for Four Dimensional Quantum Field Theories*, *Nucl. Phys. B* **883** (2014) 425 [[arXiv:1312.0428](#)] [[INSPIRE](#)].
- [7] N. Seiberg, *Naturalness versus supersymmetric nonrenormalization theorems*, *Phys. Lett. B* **318** (1993) 469 [[hep-ph/9309335](#)] [[INSPIRE](#)].
- [8] V.A. Novikov, M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Exact Gell-Mann-Low Function of Supersymmetric Yang-Mills Theories from Instanton Calculus*, *Nucl. Phys. B* **229** (1983) 381 [[INSPIRE](#)].
- [9] M.A. Shifman and A.I. Vainshtein, *Solution of the Anomaly Puzzle in SUSY Gauge Theories and the Wilson Operator Expansion*, *Nucl. Phys. B* **277** (1986) 456 [*Sov. Phys. JETP* **64** (1986) 428] [*Zh. Eksp. Teor. Fiz.* **91** (1986) 723] [[INSPIRE](#)].
- [10] D.Z. Freedman and H. Osborn, *Constructing a  $c$  function for SUSY gauge theories*, *Phys. Lett. B* **432** (1998) 353 [[hep-th/9804101](#)] [[INSPIRE](#)].
- [11] I. Jack and C. Poole, *The  $a$ -function for gauge theories*, *JHEP* **01** (2015) 138 [[arXiv:1411.1301](#)] [[INSPIRE](#)].
- [12] N. Arkani-Hamed, G.F. Giudice, M.A. Luty and R. Rattazzi, *Supersymmetry breaking loops from analytic continuation into superspace*, *Phys. Rev. D* **58** (1998) 115005 [[hep-ph/9803290](#)] [[INSPIRE](#)].
- [13] K.A. Intriligator and B. Wecht, *The exact superconformal  $R$  symmetry maximizes  $a$* , *Nucl. Phys. B* **667** (2003) 183 [[hep-th/0304128](#)] [[INSPIRE](#)].
- [14] D. Kutasov and A. Schwimmer, *Lagrange multipliers and couplings in supersymmetric field theory*, *Nucl. Phys. B* **702** (2004) 369 [[hep-th/0409029](#)] [[INSPIRE](#)].
- [15] E. Barnes, K.A. Intriligator, B. Wecht and J. Wright, *Evidence for the strongest version of the 4d  $a$ -theorem, via  $a$ -maximization along RG flows*, *Nucl. Phys. B* **702** (2004) 131 [[hep-th/0408156](#)] [[INSPIRE](#)].
- [16] D. Kutasov, *New results on the 'a theorem' in four-dimensional supersymmetric field theory*, [hep-th/0312098](#) [[INSPIRE](#)].
- [17] R.G. Leigh and M.J. Strassler, *Exactly marginal operators and duality in four-dimensional  $N = 1$  supersymmetric gauge theory*, *Nucl. Phys. B* **447** (1995) 95 [[hep-th/9503121](#)] [[INSPIRE](#)].

- [18] V. Asnin, *On metric geometry of conformal moduli spaces of four-dimensional superconformal theories*, *JHEP* **09** (2010) 012 [[arXiv:0912.2529](#)] [[INSPIRE](#)].
- [19] B. Kol, *On conformal deformations*, *JHEP* **09** (2002) 046 [[hep-th/0205141](#)] [[INSPIRE](#)].
- [20] D. Green, Z. Komargodski, N. Seiberg, Y. Tachikawa and B. Wecht, *Exactly marginal deformations and global symmetries*, *JHEP* **06** (2010) 106 [[arXiv:1005.3546](#)] [[INSPIRE](#)].
- [21] B. Keren-Zur, *The local RG equation and chiral anomalies*, *JHEP* **09** (2014) 011 [[arXiv:1406.0869](#)] [[INSPIRE](#)].
- [22] I.L. Buchbinder and S.M. Kuzenko, *Ideas and methods of supersymmetry and supergravity: A Walk through superspace*, IOP Publishing, Bristol U.K. (1995).
- [23] J. Gasser and H. Leutwyler, *Chiral Perturbation Theory to One Loop*, *Annals Phys.* **158** (1984) 142 [[INSPIRE](#)].
- [24] L. Bonora, P. Cotta-Ramusino and C. Reina, *Conformal Anomaly and Cohomology*, *Phys. Lett. B* **126** (1983) 305 [[INSPIRE](#)].
- [25] S. Ferrara and B. Zumino, *Transformation Properties of the Supercurrent*, *Nucl. Phys. B* **87** (1975) 207 [[INSPIRE](#)].
- [26] Z. Komargodski and N. Seiberg, *Comments on Supercurrent Multiplets, Supersymmetric Field Theories and Supergravity*, *JHEP* **07** (2010) 017 [[arXiv:1002.2228](#)] [[INSPIRE](#)].
- [27] Y. Nakayama, *Supercurrent, Supervirial and Superimprovement*, *Phys. Rev. D* **87** (2013) 085005 [[arXiv:1208.4726](#)] [[INSPIRE](#)].
- [28] H. Osborn,  *$N = 1$  superconformal symmetry in four-dimensional quantum field theory*, *Annals Phys.* **272** (1999) 243 [[hep-th/9808041](#)] [[INSPIRE](#)].
- [29] J. Erdmenger, C. Rupp and K. Sibold, *Conformal transformation properties of the supercurrent in four-dimensional supersymmetric theories*, *Nucl. Phys. B* **530** (1998) 501 [[hep-th/9804053](#)] [[INSPIRE](#)].
- [30] J.-F. Fortin, B. Grinstein, C.W. Murphy and A. Stergiou, *On Limit Cycles in Supersymmetric Theories*, *Phys. Lett. B* **719** (2013) 170 [[arXiv:1210.2718](#)] [[INSPIRE](#)].
- [31] L. Bonora, P. Pasti and M. Tonin, *Cohomologies and Anomalies in Supersymmetric Theories*, *Nucl. Phys. B* **252** (1985) 458 [[INSPIRE](#)].
- [32] L. Bonora and S. Giaccari, *Weyl transformations and trace anomalies in  $N = 1$ ,  $D = 4$  supergravities*, *JHEP* **08** (2013) 116 [[arXiv:1305.7116](#)] [[INSPIRE](#)].
- [33] A.B. Zamolodchikov, *Irreversibility of the Flux of the Renormalization Group in a 2D Field Theory*, *JETP Lett.* **43** (1986) 730 [*Pisma Zh. Eksp. Teor. Fiz.* **43** (1986) 565] [[INSPIRE](#)].
- [34] J. Grosse, *Quantum Field Theories Coupled to Supergravity: AdS/CFT and Local Couplings*, *Fortsch. Phys.* **56** (2008) 183 [[arXiv:0711.0444](#)] [[INSPIRE](#)].
- [35] E. Guadagnini, K. Konishi and M. Mintchev, *Nonabelian Chiral Anomalies in Supersymmetric Gauge Theories*, *Phys. Lett. B* **157** (1985) 37 [[INSPIRE](#)].
- [36] I.N. McArthur and H. Osborn, *Supersymmetric Chiral Effective Action and Nonabelian Anomalies*, *Nucl. Phys. B* **268** (1986) 573 [[INSPIRE](#)].
- [37] M. Marinkovic, *Wess-Zumino Effective Action for Supersymmetric Yang-Mills Theories*, *Nucl. Phys. B* **366** (1991) 74 [[INSPIRE](#)].

- [38] Y. Ohshima, K. Okuyama, H. Suzuki and H. Yasuta, *Remark on the consistent gauge anomaly in supersymmetric theories*, *Phys. Lett. B* **457** (1999) 291 [[hep-th/9904096](#)] [[INSPIRE](#)].
- [39] S.J. Gates Jr., M.T. Grisaru and S. Penati, *Holomorphy, minimal homotopy and the 4-D,  $N = 1$  supersymmetric Bardeen-Gross-Jackiw anomaly*, *Phys. Lett. B* **481** (2000) 397 [[hep-th/0002045](#)] [[INSPIRE](#)].
- [40] S.J. Gates Jr., M.T. Grisaru, M.E. Knutt, S. Penati and H. Suzuki, *Supersymmetric gauge anomaly with general homotopic paths*, *Nucl. Phys. B* **596** (2001) 315 [[hep-th/0009192](#)] [[INSPIRE](#)].
- [41] S.L. Adler and W.A. Bardeen, *Absence of higher order corrections in the anomalous axial vector divergence equation*, *Phys. Rev.* **182** (1969) 1517 [[INSPIRE](#)].
- [42] W.A. Bardeen and B. Zumino, *Consistent and Covariant Anomalies in Gauge and Gravitational Theories*, *Nucl. Phys. B* **244** (1984) 421 [[INSPIRE](#)].
- [43] D. Anselmi, D.Z. Freedman, M.T. Grisaru and A.A. Johansen, *Nonperturbative formulas for central functions of supersymmetric gauge theories*, *Nucl. Phys. B* **526** (1998) 543 [[hep-th/9708042](#)] [[INSPIRE](#)].
- [44] D. Anselmi, J. Erlich, D.Z. Freedman and A.A. Johansen, *Positivity constraints on anomalies in supersymmetric gauge theories*, *Phys. Rev. D* **57** (1998) 7570 [[hep-th/9711035](#)] [[INSPIRE](#)].
- [45] J. Babington and J. Erdmenger, *Space-time dependent couplings in  $N = 1$  SUSY gauge theories: Anomalies and central functions*, *JHEP* **06** (2005) 004 [[hep-th/0502214](#)] [[INSPIRE](#)].
- [46] J.L. Cardy, *Is There a  $c$  Theorem in Four-Dimensions?*, *Phys. Lett. B* **215** (1988) 749 [[INSPIRE](#)].
- [47] Z. Komargodski and A. Schwimmer, *On Renormalization Group Flows in Four Dimensions*, *JHEP* **12** (2011) 099 [[arXiv:1107.3987](#)] [[INSPIRE](#)].
- [48] Z. Komargodski, *The Constraints of Conformal Symmetry on RG Flows*, *JHEP* **07** (2012) 069 [[arXiv:1112.4538](#)] [[INSPIRE](#)].
- [49] J.-F. Fortin, B. Grinstein and A. Stergiou, *Limit Cycles and Conformal Invariance*, *JHEP* **01** (2013) 184 [[arXiv:1208.3674](#)] [[INSPIRE](#)].
- [50] M.A. Luty, J. Polchinski and R. Rattazzi, *The  $a$ -theorem and the Asymptotics of 4D Quantum Field Theory*, *JHEP* **01** (2013) 152 [[arXiv:1204.5221](#)] [[INSPIRE](#)].
- [51] D. Erkal and D. Kutasov,  *$a$ -Maximization, Global Symmetries and RG Flows*, [[arXiv:1007.2176](#)] [[INSPIRE](#)].
- [52] O. Aharony, B. Kol and S. Yankielowicz, *On exactly marginal deformations of  $N = 4$  SYM and type IIB supergravity on  $AdS_5 \times S^5$* , *JHEP* **06** (2002) 039 [[hep-th/0205090](#)] [[INSPIRE](#)].
- [53] B. Kol, *On Conformal Deformations II*, [[arXiv:1005.4408](#)] [[INSPIRE](#)].
- [54] S.J. Gates, M.T. Grisaru, M. Roček and W. Siegel, *Superspace Or One Thousand and One Lessons in Supersymmetry*, *Front. Phys.* **58** (1983) 1 [[hep-th/0108200](#)] [[INSPIRE](#)].
- [55] S. Abel, M. Buican and Z. Komargodski, *Mapping Anomalous Currents in Supersymmetric Dualities*, *Phys. Rev. D* **84** (2011) 045005 [[arXiv:1105.2885](#)] [[INSPIRE](#)].
- [56] M. Buican, *A Conjectured Bound on Accidental Symmetries*, *Phys. Rev. D* **85** (2012) 025020 [[arXiv:1109.3279](#)] [[INSPIRE](#)].

- [57] H. Elvang et al., *On renormalization group flows and the a-theorem in 6d*, *JHEP* **10** (2012) 011 [[arXiv:1205.3994](#)] [[INSPIRE](#)].
- [58] B. Grinstein, A. Stergiou and D. Stone, *Consequences of Weyl Consistency Conditions*, *JHEP* **11** (2013) 195 [[arXiv:1308.1096](#)] [[INSPIRE](#)].
- [59] B. Grinstein, D. Stone, A. Stergiou and M. Zhong, *Challenge to the a Theorem in Six Dimensions*, *Phys. Rev. Lett.* **113** (2014) 231602 [[arXiv:1406.3626](#)] [[INSPIRE](#)].
- [60] H. Osborn and A. Stergiou, *Structures on the Conformal Manifold in Six Dimensional Theories*, *JHEP* **04** (2015) 157 [[arXiv:1501.01308](#)] [[INSPIRE](#)].
- [61] J. Wess and J. Bagger, *Supersymmetry and supergravity*, Princeton University Press, Princeton U.S.A. (1992).
- [62] P.S. Howe and R.W. Tucker, *Scale Invariance in Superspace*, *Phys. Lett. B* **80** (1978) 138 [[INSPIRE](#)].
- [63] V. Kaplunovsky and J. Louis, *Field dependent gauge couplings in locally supersymmetric effective quantum field theories*, *Nucl. Phys. B* **422** (1994) 57 [[hep-th/9402005](#)] [[INSPIRE](#)].
- [64] A. Schwimmer and S. Theisen, *Spontaneous Breaking of Conformal Invariance and Trace Anomaly Matching*, *Nucl. Phys. B* **847** (2011) 590 [[arXiv:1011.0696](#)] [[INSPIRE](#)].
- [65] T.E. Clark, O. Piguet and K. Sibold, *The Absence of Radiative Corrections to the Axial Current Anomaly in Supersymmetric QED*, *Nucl. Phys. B* **159** (1979) 1 [[INSPIRE](#)].
- [66] K. Konishi, *Anomalous Supersymmetry Transformation of Some Composite Operators in SQCD*, *Phys. Lett. B* **135** (1984) 439 [[INSPIRE](#)].